University of Rajshahi	Rajshahi-6205	Bangladesh.
RUCL Institutional Repository		http://rulrepository.ru.ac.bd
Institute of Biological Sciences (IBSc)		PhD Thesis

2021-05

Sustainable Improvement of Wheat (Triticum Aestivum L.) Yield Through Soil Amendments with Probiotic and Organic Manures

Hossain, Md. Anwar

University of Rajshahi

http://rulrepository.ru.ac.bd/handle/123456789/1043 Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

SUSTAINABLE IMPROVEMENT OF WHEAT (*TRITICUM AESTIVUM* L.) YIELD THROUGH SOIL AMENDMENTS WITH PROBIOTIC AND ORGANIC MANURES



THESIS SUBMITTED FOR THE DEGREE OF DOCTOR OF PHILOSOPHY IN THE INSTITUTE OF BIOLOGICAL SCIENCES UNIVERSITY OF RAJSHAHI BANGLADESH

BY

MD. ANWAR HOSSAIN

B. Sc. Ag., MS (AGRONOMY)

PLANT BIOTECHNOLOGY AND GENETIC ENGINEERING LABORATORY INSTITUTE OF BIOLOGICAL SCIENCES UNIVERSITY OF RAJSHAHI RAJSHAHI-6205 BANGLADESH

MAY 2021



DECLARATION

I hereby declare that the research work embodied in this thesis entitled "SUSTAINABLE IMPROVEMENT OF WHEAT (*TRITICUM AESTIVUM* L.) YIELD THROUGH SOIL AMENDMENTS WITH PROBIOTIC AND ORGANIC MANURES" has been carried out by me for the degree of Doctor of Philosophy under the guidance of Professor Dr. S. M. Shahinul Islam (Principal Supervisor), Institute of Biological Sciences, University of Rajshahi and Professor Dr. Md. Mahmodol Hasan (Co-supervisor), Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh.

I also declare that the result presented in this dissertation is my own investigation and any part of this thesis work has not been submitted to elsewhere for any degree/diploma or for similar purpose.

May 2021

Md. Anwar Hossain PhD Fellow Roll No. P-532 Session: 2016-2017 Institute of Biological Sciences University of Rajshahi Rajshahi-6205 Bangladesh



CERTIFICATE

This is to certify that **Md. Anwar Hossain** worked under our supervision as a PhD Fellow, Roll No: P-532, Session: 2016-2017, Institute of Biological Sciences, University of Rajshahi, Bangladesh. It is our great pleasure to forward his thesis entitled **"SUSTAINABLE IMPROVEMENT OF WHEAT (***TRITICUM AESTIVUM* **L.) YIELD THROUGH SOIL AMENDMENTS WITH PROBIOTIC AND ORGANIC MANURES"** which is a genuine record of research carried out at Plant Biotechnology and Genetic Engineering Laboratory, Institute of Biological Sciences and its experimental field and in the Plant Pathology Laboratory, Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh.

This work is original and has not been submitted so far in part or in full, for the award of any degree or diploma by any other institute in home or abroad. Md. Anwar Hossain has fulfilled all the requirements for submission of the thesis for the award of the degree of **Doctor of Philosophy.**

Principal Supervisor

Co-Supervisor

Dr. S. M. Shahinul Islam Professor Institute of Biological Sciences University of Rajshahi Rajshahi-6205, Bangladesh Phone: +88-01715-209907 E-mail: shahinul68@gmail.com

Dr. Md. Mahmodol Hasan Professor Dept. of Agronomy and Agricultural Extn. University of Rajshahi Rajshahi-6205, Bangladesh Phone: +88-01739-824746 E-mail: mahmudul742001@yahoo.com

CONTENTS

Page No.

Declaration	i
Certificate	ii
Contents	iii-x
Acknowledgements	xi
Abstract	xii-xiii
List of Figures	xiv-xv
List of Tables	xvi-xviii
List of Appendices	xix
List of Abbreviations	xx-xxiii
Chapter I: Introduction	1-7
Chapter II: Review of Literature	8-41
2.1 Fertility situation of soil in Bangladesh	8
2.1.1 Status of organic matter	9
2.1.2 Selected essential nutrient status	10
2.2 Organic amendments and soil properties	11
2.2.1 Non-leguminous crop residue incorporation and soil properties	11
2.2.2 Leguminous residue incorporation and soil properties	13
2.2.3 Cow dung incorporation and soil properties	16
2.2.4 Compost incorporation and soil properties	17
2.2.5 Vermicompost incorporation and soil properties	19

2.2.6 Poultry manure incorporation and soil properties	20
2.2.7 Green manure incorporation and soil properties	21
2.3 Probiotic inoculation and soil amendments	23
2.3.1 <i>Trichoderma</i> spp. inoculation for soil	23
2.4 Chemical fertilizer and soil properties	24
2.5 Organic amendments for wheat growth and yield	26
2.5.1 Non-leguminous crop residue incorporation	26
2.5.2 Leguminous crop residue incorporation	27
2.5.3 Cow dung incorporation	29
2.5.4 Compost incorporation	31
2.5.5 Vermicompost incorporation	32
2.5.6 Poultry manure incorporation	33
2.5.7 Green manure incorporation	36
2.6 Probiotic inoculation for wheat growth and yield	37
2.6.1 <i>Trichoderma</i> spp. inoculation for wheat	37
2.7 Chemical fertilizer for wheat growth and yield	40
Chapter III: Materials and Methods	42-60
3.1 Description of the experimental site	42
3.1.1 Location	42
3.1.2 Soil properties	42
3.1.2.1 Physical properties	42
3.1.2.2 Chemical properties	42
3.1.3 Climatic condition	43
3.2 Cropping season	43

3.3 Cropping history of the experimental plot	43
3.4 Experiment	43
3.4.1 Factors and treatments	43
3.4.2 Experimental design	44
3.4.3 Description of wheat varieties	46
3.4.4 Green manuring	46
3.4.4.1 Land preparation	46
3.4.4.2 Seed sowing	47
3.4.4.3 Harvesting	47
3.4.5 Establishment and management of wheat crop	47
3.4.5.1 Land preparation	47
3.4.5.2 Treatment application	47
3.4.5.3 Collection of seed	48
3.4.5.4 Seed sowing	48
3.4.5.5 Weeding	48
3.4.5.6 Irrigation	48
3.4.5.7 Harvesting and processing	48
3.4.6 Soil sample collection and analysis	50
3.4.6.1 Soil sample collection	50
3.4.6.2 Physical properties analysis	50
3.4.6.2.1 Moisture content	50
3.4.6.2.2 Particle density	50
3.4.6.2.3 Bulk density and porosity	51
3.4.6.2.4 Textural class	51

3.4.6.2.5 Chemical analysis	52
3.4.6.2.6 Soil pH	52
3.4.6.2.7 Organic matter	53
3.4.6.2.8 Total nitrogen	53
3.4.6.2.9 Available phosphorus	53
3.4.6.2.10 Exchangeable potassium	53
3.4.6.2.11 Available sulphur	53
3.4.6.2.12 Available zinc	53
3.4.6.2.13 Electrical conductivity	54
3.4.7 Crop characters studied	54
3.4.7.1 Seedling infection	54
3.4.7.2 Seedling blight	54
3.4.7.3 Leaf chlorophyll content	54
3.4.7.4 Growth parameters of wheat	55
3.4.7.4.1 Total dry matter	55
3.4.7.4.2 Leaf area index and crop growth rate	55
3.4.7.5 Yield and yield contributing characters of wheat	56
3.4.7.5.1 Plant height	56
3.4.7.5.2 Total plant/m ²	56
3.4.7.5.3 Total tiller/plant	56
3.4.7.5.4 Effective tiller/plant	56
3.4.7.5.5 Awn length	56
3.4.7.5.6 Spike length	57
3.4.7.5.7 Spikelet/spike	57

3.4.7.5.8 Fertile spikelet/spike	57
3.4.7.5.9 Grains/spike	57
3.4.7.5.10 Deformed grains/spike	57
3.4.7.5.11 Grain weight/spike	57
3.4.7.5.12 1000-grain weight	57
3.4.7.5.13 Grain yield	58
3.4.7.5.14 Straw yield	58
3.4.7.5.15 Harvest index	58
3.4.8 Seed quality parameters of wheat	58
3.4.8.1 Germination	58
3.4.8.1.1 Germination media	58
3.4.8.1.2 Determination of germination	59
3.4.8.2 Vigor index	59
3.4.8.3 Total soluble protein content of wheat seed	59
3.4.9 Statistical analysis	60
Chapter IV: Results	61-144
4.1 Soil properties	61
4.1.1 Physical properties	61
4.1.1.1 Soil moisture	61
4.1.1.2 Bulk density	63
4.1.1.3 Particle density	64
4.1.1.4 Soil porosity	65
4.1.1.5 Soil sand particle	66
4.1.1.6 Soil silt particle	67

	4.1.1.7 Soil clay particle	68
	4.1.1.8 Soil textural class	69
	4.1.2 Chemical properties	70
	4.1.2.1 Soil pH	70
	4.1.2.2 Soil organic matter	71
	4.1.2.3 Total nitrogen	72
	4.1.2.4 Exchangeable potassium	73
	4.1.2.5 Available phosphorus	74
	4.1.2.6 Available sulphur	75
	4.1.2.7 Available zinc	76
	4.1.2.8 Electrical conductivity	77
	4.1.2.9 Carbon nitrogen ratio	78
4.2	Wheat	79
	4.2.1 Seedling infection	79
	4.2.2 Seedling blight	79
	4.2.3 Leaf chlorophyll content	80
4.3	Growth attributes	85
	4.3.1 Total dry matter	85
	4.3.2 Leaf area index	92
	4.3.3 Crop growth rate	98
4.4	Yield and yield contributing characters	105
	4.4.1 Plant height	105
	4.4.2 Total plant/m ²	105

4.4.3 Total tiller/plant	106
4.4.4 Effective tiller/plant	107
4.4.5 Awn length	113
4.4.6 Spike length	113
4.4.7 Spikelet/spike	114
4.4.8 Fertile spikelet/spike	115
4.4.9 Grains/spike	121
4.4.10 Deformed grains/spike	121
4.4.11 Grain weight/spike	122
4.4.12 1000-grain weight	123
4.4.13 Grain yield	129
4.4.14 Straw yield	130
4.4.15 Harvest index	134
4.5 Seed quality	138
4.5.1 Germination	138
4.5.2 Vigor index	139
4.5.3 Total soluble protein content	140
Chapter V: Discussion	145-157
5.1 Effects of soil amendments on soil properties	145
5.1.1 Physical properties	145
5.1.2 Chemical properties	147
5.2 Effects of soil amendments on growth and yield of wheat	150
5.2.1 Seedling survivability	150

5.2.2 Leaf chlorophyll content	151
5.2.3 Growth attributes	151
5.2.4 Yield and yield contributing attributes	152
5.2.5 Seed quality	156
Chapter VI: Summary	
Chapter VII: References	163-197
Appendices	198-200
List of Publications and Abstract	
Curriculum Vitae	

ACKNOWLEDGEMENTS

Firstly, I would like to show due respect to the **Almighty ALLAH** who enabled me to complete this research successfully.

It is a debt of gratitude and pleasant duty to express my devotion to my reverent research supervisors Professor Dr. S. M. Shahinul Islam, Institute of Biological Sciences, University of Rajshahi and Professor Dr. Md. Mahmodol Hasan, Department of Agronomy and Agriculture Extension, University of Rajshahi for their continuous encouragement, indefatigable patience, proper guidance, valuable suggestions and stimulating discussions throughout the research work and preparing this paper. I am grateful for their intellectual and valuable supervisory activities.

I am indebted to Dr. Md. Firoz Alam, Professor and Director, Institute of Biological Sciences, University of Rajshahi and others respectable teachers for their encouragement and help in need.

I sincerely acknowledge Professor Dr. Md. Belal Hossain and Professor Dr. Md. Ruhul Amin, Department of Agronomy and Agricultural Extension, University of Rajshahi for valuable suggestions during research as well as thesis presentation.

I convey my heartfelt gratitude to Md. Sazzad Hossain, Principal Scientific Officer, Plant Physiology Division, BARI, Gazipur, Md. Shakawat Hossain, Senior Scientific Officer, On Farm Research Division, BARI, Rajshahi, Md. Nur-E-Alam, Senior Plant Breeder, Regional Wheat Research Centre, Rajshahi and Md. Aminul Islam, Senior Scientific Officer, Regional Soil Resource Development Institute, Rajshahi for their technical support and sincere cooperation in analyzing relevant data.

My special thanks to Dr. Md. Jahangir Alam, Dr. Md. Nazrul Islam, Dr. Mirza Fida Hasan, Zannatul Fardous and Mahabua Akter, PhD Fellow and others research colleagues at the Institute of Biological Sciences, University of Rajshahi for their constant co-operation. I wish to extend my especial thanks to all my colleagues, friends and juniors.

I express my deepest gratitude to my parents who opened my eyes for acquiring knowledge, their blessings and continuous inspiration, which led to achieving this success in my life.

I express my deep thanks to my wife Mst. Noor-Un-Naher Sarkar, my daughter Noshin Arbab Nubah, brother and sisters for their encouragement and inspiration to complete this research work.

Finally, the Ministry of Science and Technology, Govt. of Bangladesh and Bangabandhu Science and Technology Fellowship Trust for their financial support and the kind co-operation of all staff of the Institute of Biological Sciences and Department of Agronomy and Agricultural Extension, University of Rajshahi is gratefully acknowledged.

The Author

ABSTRACT

The present research work was conducted at Plant Biotechnology and Genetic Engineering Laboratory and the experimental field under Institute of Biological Sciences (IBSc), Rajshahi University and Plant Pathology Laboratory, Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh during the period from July 2016 to June 2019 to study the sustainable improvement of wheat (Triticum aestivum L.) yield through soil amendments with probiotic and organic manures. The treatments used in the research consisted of two factors i.e. factor-A (wheat varieties) viz. BARI wheat-28 (V_1) , BARI wheat-29 (V_2) , BARI wheat-30 (V_3) and **factor-B** (soil amendments) viz. Rice straw + vermicompost + green manure (T_1) , Cow dung + vermicompost + green manure (T_2) , Compost + vermicompost + green manure (T_3) , Poultry manure + vermicompost + green manure (T_4) , Trichoderma harzianum + vermicompost + green manure (T_5) , Mung bean residue + vermicompost + green manure (T_6), Trichoderma viride + vermicompost + green manure (T_7) , Chemical fertilizer (T_8) and Control (T_0) . Sesbania rostrata was produced as green manure crop in this study. The experiment was designed as randomized complete block design (RCBD) having three replications. Initially the soil was analyzed and mainly featured with loam textured, bulk densities 1.27 g/cc, particle densities 2.65 g/cc, pH 8.10 and organic matter 1.20%. After organic amendments the value in respect of bulk density, particle density and sand particles which were found to be decreased though porosity, moisture content and silt particles were increased. The chemical characters of soil like pH and carbon nitrogen ratio were noted as negative changes with probiotic and organic manures amendments. Furthermore, the organic matter content, total N, exchangeable K, available P, available S, available Zn and electrical conductivity were found to be improved during the three sequential years of the study. Seedling infection (14 days after sowing) and seedling blight (21 days after sowing) was examined and recorded significant reduction by the application of probiotic and organic manures. The probiotic fungi T. harzianum (T₅) significantly reduced the seedling infection and seedling blight by 86% and 81% in comparison with control treatment.

Total dry matter content increased with increasing the age of the crop. The leaf area index and crop growth rate was lower at the early stage of crop growth and reached a maximum during 55 and 55-70 days after sowing and then declined. All the aforesaid growth parameters were improved by the application of probiotic and organic manures application and produced maximum value especially with poultry manure combination (T_4) over control. Studied yield contributing characters including plant height, total plant/m², total tiller/plant, effective tiller/plant, awn length, spike length, spikelet/spike, fertile spikelet/spike, grains/spike, deformed grains/spike, grain weight/spike and 1000-grain weight were influenced significantly due to application of organic amendments. As grain yield positively linked with the above mentioned yield contributing characters, thus grain yield was significantly enhanced with organic amendments. Besides this, the straw yield and harvest index was also recorded upturned with the similar amendments. Grain yield, straw yield and harvest index showed 73%, 27% and 22% greater response at the final research period under poultry manure combination (T₄) over control. Seed quality parameters such as germination, vigor index and total soluble protein content significantly affected by different organic amendments. Among the treatments, poultry manure combination (T_4) exhibited higher results in this regard, whereas lower value was recorded in control. However, with few exceptions, V₃ (BARI Wheat-30) and T₄ (Poultry manure + vermicompost + green manure) not only individually performed better results than others but also their interactivity (V_3T_4) exhibited better results under this study.

In connection with the above findings, it can be recommended that the wheat growers to adopt the most efficient soil amendment option poultry manure + vermicompost + green manure (T_4) to improve soil fertility and productivity by its positive changing ability and to enhance the growth, yield and grain quality of wheat in a sustainable manner.

LIST OF FIGURES

Figure No.	Title	Page No.
1	Different treatment materials used in the experiment	45
2	Scenario of crop production and data collection	49
3	Effect of variety on total dry matter content of wheat in 2016-2017	87
4	Effect of variety on total dry matter content of wheat in 2017-2018	87
5	Effect of variety on total dry matter content of wheat in 2018-2019	88
6	Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2016-2017	88
7	Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2017-2018	89
8	Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2018-2019	89
9	Effect of variety on leaf area index of wheat in 2016-2017	93
10	Effect of variety on leaf area index of wheat in 2017-2018	93
11	Effect of variety on leaf area index of wheat in 2018-2019	94
12	Effect of organic and probiotic soil amendments on leaf area index of wheat in 2016-2017	94
13	Effect of organic and probiotic soil amendments on leaf area index of wheat in 2017-2018	95
14	Effect of organic and probiotic soil amendments on leaf area index of wheat in 2018-2019	95
15	Effect of variety on crop growth rate of wheat in 2016-2017	100
16	Effect of variety on crop growth rate of wheat in 2017-2018	100

Figure No.	Title	Page No.
17	Effect of variety on crop growth rate of wheat in 2018-2019	101
18	Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2016-2017	101
19	Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2017-2018	102
20	Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2018-2019	102
21	Effect of variety on grain and straw yield of wheat in 2016-2017	131
22	Effect of variety on grain and straw yield of wheat in 2017-2018	131
23	Effect of variety on grain and straw yield of wheat in 2018-2019	132
24	Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2016-2017	132
25	Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2017-2018	133
26	Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2018-2019	133
27	Effect of variety on harvest index of wheat in 2016-2017, 2017-2018 and 2018-2019	135
28	Effect of organic and probiotic soil amendments on harvest index of wheat in 2016-2017, 2017-2018 and 2018-2019	135

LIST OF TABLES

Table No.	Title	Page No.
1	Changes of soil moisture content (%) as influenced by different soil amendments for three consecutive years of wheat production	62
2	Changes of soil bulk density (g/cc) as influenced by different soil amendments for three consecutive years of wheat production	63
3	Changes of soil particle density (g/cc) as influenced by different soil amendments for three consecutive years of wheat production	64
4	Changes of soil porosity (%) as influenced by different soil amendments for three consecutive years of wheat production	65
5	Changes of soil sand particle (%) as influenced by different soil amendments for three consecutive years of wheat production	66
6	Changes of soil silt particle (%) as influenced by different soil amendments for three consecutive years of wheat production	67
7	Changes of soil clay particle (%) as influenced by different soil amendments for three consecutive years of wheat production	68
8	Changes of soil textural class as influenced by different soil amendments for three consecutive years of wheat production	69
9	Changes of soil pH as influenced by different soil amendments for three consecutive years of wheat production	70
10	Changes of soil organic matter content (%) as influenced by different soil amendments for three consecutive years of wheat production	71
11	Changes of soil total nitrogen (%) as influenced by different soil amendments for three consecutive years of wheat production	72
12	Changes of soil exchangeable potassium (cmol (+)/kg) as influenced by different soil amendments for three consecutive years of wheat production	73

Table No.	Title	Page No.
13	Changes of soil available phosphorus $(\mu g/g)$ as influenced by different soil amendments for three consecutive years of wheat production	74
14	Changes of soil available sulphur $(\mu g/g)$ as influenced by different soil amendments for three consecutive years of wheat production	75
15	Changes of soil available zinc $(\mu g/g)$ as influenced by different soil amendments for three consecutive years of wheat production	76
16	Changes of soil electrical conductivity (μ s/cm) as influenced by different soil amendments for three consecutive years of wheat production	77
17	Changes of soil carbon nitrogen ratio (C:N) as influenced by different soil amendments for three consecutive years of wheat production	78
18	Effect of variety on seedling infection, seedling blight and leaf chlorophyll content of wheat	81
19	Effect of organic and probiotic soil amendments on seedling infection, seedling blight and leaf chlorophyll content of wheat	82
20	Interaction effect of variety and soil amendments on seedling infection, seedling blight and leaf chlorophyll content of wheat	83
21	Interaction effects variety and of soil amendments on total dry matter content of wheat (g/m^2)	90
22	Interaction effects of variety and soil amendments on leaf area index of wheat	96
23	Interaction effects of variety and soil amendments on crop growth rate of wheat $(g/m^2/d)$	103
24	Effect of variety on yield and yield contributing characters of wheat	109
25	Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat	110
26	Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat	111

Table No.	Title	Page No.
27	Effect of variety on yield and yield contributing characters of wheat	117
28	Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat	118
29	Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat	119
30	Effect of variety on yield and yield contributing characters of wheat	125
31	Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat	126
32	Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat	127
33	Interaction effects of variety and soil amendments on grain yield, straw yield and harvest index of wheat	136
34	Effect of variety on seed quality of wheat	141
35	Effect of organic and probiotic soil amendments on seed quality of wheat	142
36	Interaction effects of variety and soil amendments on seed quality of wheat	143

LIST OF APPENDICES

Table No.	Title	Page No.
1	Scenario of soil organic matter depletion during 20 years in different Agro-Ecological Zones of Bangladesh	198
2	Emergence of new nutrient deficiency with time in soil	199
3	Nutrient elements content in different organic matter	199
4	Monthly minimum, maximum and average air temperature, rainfall and relative humidity	200

LIST OF ABBREVIATIONS

The following abbreviations have been used through the text:

AEZ	:	Agro-Ecological Zone
ANOVA	:	Analysis of variance
BARI	:	Bangladesh Agricultural Research Institute
BBS	:	Bangladesh Bureau of Statistics
BD	:	Bulk density
C:N	:	Carbon nitrogen ratio
CARE	:	Cooperative for American Relief Everywhere
CBB	:	Coomassie Brilliant Blue
CDM	:	Cattle dung manure
CGR	:	Crop growth rate
cm	:	Centimeter
cm cmol (+)/kg	:	Centimeter Centimoles of positive charge per kilogram of soil
cmol (+)/kg	:	Centimoles of positive charge per kilogram of soil
cmol (+)/kg CRB	:	Centimoles of positive charge per kilogram of soil Crop residue burning
cmol (+)/kg CRB CRI	:	Centimoles of positive charge per kilogram of soil Crop residue burning Crown root initiation
cmol (+)/kg CRB CRI DAS	:	Centimoles of positive charge per kilogram of soil Crop residue burning Crown root initiation Days after sowing
cmol (+)/kg CRB CRI DAS DMRT	:	Centimoles of positive charge per kilogram of soil Crop residue burning Crown root initiation Days after sowing Duncan's Multiple Range Test
cmol (+)/kg CRB CRI DAS DMRT EC	:	Centimoles of positive charge per kilogram of soil Crop residue burning Crown root initiation Days after sowing Duncan's Multiple Range Test Electrical conductivity

FC	:	Field capacity
Fig.	:	Figure
FYM	:	Farmyard manure
FYMC	:	Composted farmyard manure
g	:	Gram
g/m ²	:	Gram per meter square
g/m ² /d	:	Gram per meter square per day
GDP	:	Gross domestic product
ha	:	Hectare
hr	:	Hour
i.e.	:	That is
IPNS	:	Integrated plant nutrient system
Κ	:	Potassium
Kg	:	Kilogram
Kharif	:	The wet season in Bangladesh (March to October)
LAI	:	Leaf area index
LS	:	Level of significance
LSD	:	Least significant difference
µg/g	:	Microgram per gram
m	:	Meter
m ²	:	Meter square
meq	:	Milliequivalent
Mg	:	Mega gram

Mg/g FW	:	Milligram per gram fresh weight
mm	:	Millimeter
MWHC	:	Maximum water holding capacity
Ν	:	Nitrogen
nm	:	Nanometer
NPK	:	Nitrogen, Phosphorus and Potassium
NS	:	Non-significant
OM	:	Organic matter
Р	:	Phosphorus
PD	:	Particle density
PGS	:	Participatory guarantee system
\mathbf{P}^{H}	:	Potentiality of hydrogen ion
ppm	:	Parts per million
Rabi	:	The dry season in Bangladesh (November to February)
RCBD	:	Randomized complete block design
S	:	Sulphur
SE	:	Standard error
SGM	:	Sesbania green manure
SOC	:	Soil organic carbon
SOM	:	Soil organic matter
SPAD	:	The soil plant analysis development
SRDI	:	Soil Resource Development Institute
t	:	Ton

TDM	:	Total dry matter
TMC	:	Trichoderma multiplied culture
UNDP	:	United Nations Development Programme
USDA	:	United States Department of Agriculture
viz.	:	Namely
WHC	:	Water holding capacity
Zn	:	Zinc
%	:	Percent
/	:	Per

Chapter I

Introduction

1. INTRODUCTION

Wheat (Triticum aestivum L.) has long been a highly popular cereal crop all over the world, as it grows in many countries in the world. It is the third most widely produced agricultural commodity next to rice and maize, and it is a staple food of billions of people all over the globe. Bangladesh possesses 31st position among 80 wheat growing countries over the world, whereas the European Union possesses 1st position. Near about 50% total world grain crop production is occupied by wheat (Banglapedia 2014). Wheat occupies in Bangladesh 4% of the total cropped area and 11% of the cropped area in Rabi season and contributes 7% to the total output of food cereals (BBS 2008). In 2017, wheat occupies 415339 ha areas with yield 1311473 t which is 2.72% less compared over the year 2016 (BBS 2017). In Bangladesh, the average yield of wheat 3.16 t/ha (BBS 2017) which is below the achievable yield of 4.5 t/ha of wheat (BARI 2011). Alam et al. (2013) stated two reasons for the yield gap as (i) biotic factors including poor quality seeds and seedlings, insects, diseases, weeds and rodents; and (ii) abiotic factors including soil, nutrients and water. However, many reasons for this yield gap remain unexplained. Wheat is a global commodity which is being traded about 150 ton annually (World Agricultural Outlook Board 2014). It is a rich source of calories (327 kcal/100g) well as multiple nutrients such as protein, dietary fiber and minerals, manganese, phosphorus, niacin and several vitamins (Shewry et al. 2002). The contribution of wheat is being increased significantly in different countries as Nigeria (1%-6.64%), India (11.85-20.41%) and China (12.20-17.83%) regarding total kcal (Shewry and Hey 2015).

Bangladesh is known as a rapidly increasing populated country and thus land under cultivation is decreasing day by day to provide their accommodation. To keep harmony with increased population, intensive cultivation of land causes land degradation which affects food security. To face the land limitation problem and provide food for increased population sustainable use of land is burning issue for agricultural development (Alam et al. 2020). It is widely agreed that long-term application of excess inorganic fertilizer or poor fertilization practice may result in soil acidification and nutrient depletion as well as,

health and environmental hazards (Liebig et al. 2002). For long-term application of unbalanced synthetic fertilizer, it is assumed that after a certain period of time the land will lose its fertility. Furthermore, synthetic fertilizers are chemicals in nature that are harmful to humans and animals. So, for achieving higher yield and to retain soil health sound, use of high yielding varieties along with organic amendments are felt to be important. The wheat-rice cropping system has emerged as the most productive cereal grain producing system in Bangladesh. This pattern is highly productive but very exhaustive. In Bangladesh, it has been observed that the nutrient balance for N is moderately negative, P is slightly negative and K is highly negative in soils (Rijpma and Jahiruddin 2004). The soil fertility, climatic conditions, characters of cultivar and yield considerably influence the nutrients requirements of the growing crops.

Soil organic matter is a positive factor for sustainable soil fertility and productivity. Organic matter undergoes mineralization with the release of substantial amounts of N, P, K, S and smaller amounts of micronutrients. Easily decomposable part of the soil organic matter undergoes mineralization quickly and becomes a part of soil humus, a small portion of which may remain in the soil. A good soil should have organic matter content of more than 2.5%. Most of the soils in Bangladesh have low to very low organic matter content, generally less than 1.5% (BARC 2012). The environmental conditions such as high temperature, high relative humidity and frequent tillage operation for high cropping intensity are favoring rapid decomposition that accelerates degradation of soil organic matter in Bangladesh (Ahsan and Karim 1986). Soil organic matter status strongly associated with soil productivity. Organic amendment improves the quality of soil, reduces compaction and crusting, and increases drainage and water-holding capacity. Plants become healthier, tolerant of drought, insects and diseases and more yields by providing organic matter. Soil organic matter serves as a reservoir for plant nutrients thus need less fertilizer (Cooperband 2002). Organic materials like crop residues, green manure, and animal manure and their spontaneous use shows great influence on soil productivity and N dynamics in the soil-plant system (Yadvinder-Singh et al. 2008). As a part of organic amendments of soil beneficial microbe populations are essential to creating the ideal soil environment for healthy plant growth. Probiotics like *Trichoderma* may be environmentally-friendly alternatives to the chemical fertilizers to increase the soil fertility and the crops productivity and yield without causing harmful environmental effects (Hajieghrari and Mohammadi 2016).

South Asian farmers need to manage 5-7 t/ha of rice residues and overcome the problems for planting wheat. Burning of rice straw before wheat planting results in huge losses of N (up to 80%), P (25%), K (21%) and S (4-60%), air pollution (CO₂ 13 t/ha) and soils depriving organic matter (SOM). This loss of SOM is one of the recognized threats to sustainability (Mandal et al. 2004). Residue management practices affect soil physical properties viz. soil moisture, temperature, aggregate formation and bulk density. Rice crop residues reduce P fixation; improving base retention and increasing the soil pH. Rice straw incorporation coupled with organic manure increases grain yield of wheat as well as improves soil physical condition. The soils having crop residue enhances the microbial activities than residue removal or burning. Besides the rice straw, grain legumes mung bean also offers considerable increased value in production when its plant residue is incorporated into soil. Incorporation of N-fixing legume crop residues into the soil may provide organic N for the subsequent benefit of a cereal crop (Rosales et al. 1998). The use of mung bean residues in the rice-wheat cropping system was equivalent to the use of 30-120 kg urea-N/ha (Sharma et al. 2000). However, crop residue might also affect other factors, such as the soil pH and soil organic content and also soil structure and plant growth (Ghassan et al. 2017). Thus, if residues are managed properly, then it can warrant the improvements in soil properties and the sustainability in crop productivity. The most economical and ecological means of conserving soil and water and sustaining crop production is returning crop residues to soil (Wilhelm et al. 2007).

In Bangladesh, cow dung is generally used for crop production, but small mechanization in the crop production field resulted in lessen use of livestock as well as cow dung. Then again, generally rural farmers are using cow dung as fuel for cooking purposes. Cow manure is packed with high levels of minerals and nutrients and is one of the best natural fertilizers to use in organic farming. Cow dung contains an average of 1.04% nitrogen,

0.15% potassium and 0.78% phosphorus and 32 other micronutrients. Manure applications result in increases in pH, water holding capacity and infiltration rates (Ram 2017). Soil conditioning ability of cow manure is outstanding which is not comparable to fertilizer.

Land is the major non-renewable resource and faces the biggest threat of degradation. Nutrients availability of soils is declining with time pace but which have been rich in the past (Zia et al. 1994). Use of compost can be beneficial to improve organic matter status of soil. Compost is a prosperous source of nutrients with high organic matter content. Physical and chemical properties of soil can be improved by using compost, which may ultimately increase crop yields (Sarwar et al. 2008). Besides soil health improvement, compost leads to managing large volumes of organic wastes in a comprehensive manner (Lasaridi and Stentiford 1999).

Vermicomposting is an effective means of composting the decomposable organic wastes using earthworms naturally present in the soil. Vermicompost is an excellent vehicle for carrying nutrients to soil and plants as it is more than a fertilizer (Sheela and Khimiya 2013). Vermicompost is a mixture of worm casts enriched with macro and micronutrients (N, P, K, Mn, Fe, Mo, B, Cu and Zn), some growth regulating substances (gibberellins and auxins) and useful to micro flora which naturally helps in improvement of soil fertility or soil quality (Dandotiya and Agrawal 2014). Vermicompost is associated with the improvement of soil physical properties like aggregate formation, bulk density and porosity having its excellent structure, porosity, aeration, drainage and moisture holding capacity (Dominguez et al. 1997). In fact, there must be a balance between soil degradation and restoration process to determine sustainability. To maintain these phenomena vermicompost may be a viable tool for soil amendment and signifies quality produce of crops.

Green manure is a once grown crop and usually incorporated to the soil before the next cash crop. It is grown for a short time and dug into soil so that organic matter in the crop can be transferred back into the soil. In Bangladesh, the aus rice is cultivated from March and April after the pre-monsoon rainfall and harvested between July and August (Shelley et al. 2016).

Thus after harvest of rice and cultivation of wheat there is a short fallow period of 40-70 days. This fallow period of about 2 months between rice harvest and wheat cultivation can be used effectively for raising a suitable legume crop. In this system crop productivity and maintaining soil fertility can be definitely improved by adding green manures. It has been noticed by researchers that green manures increase soil biological and enzymatic activities more than mineral fertilizers (Bolton et al. 1985, Kirchner et al. 1993, Abdallahi and N'Dayegamiye 2000). Green manures also showed remarkable improvement in soil structural stability and higher water-holding capacities (Muller et al. 1988, Kuo et al. 1997, Abdallahi and N'Dayegamiye 2000) and resulting in higher crop yields and nutrients availability.

Recently, the poultry industry possesses a remarkable stand and creates an opportunity to utilize poultry manure in organic farming in Bangladesh. Poultry manure is available fertilizer and can serve as a suitable alternative to chemical fertilizer. Over 53% increase of N level in the soil from 0.09 to 0.14% and exchangeable cations increase was noticed with manure application (Boateng et al. 2006). Mainly poultry manure provided to soil in order to include the organic amendment of the soil and the provision of nutrients to crops (Warren et al. 2006). As poultry manure contains high N, P, K and other essential nutrients thus it is an excellent organic fertilizer (Farhad et al. 2009). Poultry manure has been reported to supply P ready to plant than other organic sources (Garg and Bahla 2008). It shows more significant compared to farmyard manure and vermicompost (Behara and Pandey 2002) and its application was near about 1.5 times more effective than compost (Singh et al. 1983). Li et al. (2011) explained that poultry and livestock manure influence pore structure and aggregate stability where pore structure plays a very important role in soil structure formation, soil moisture and maintaining nutrients, and protection of microbial diversity, while aggregate stability has a positive impact on the seed germination, plant roots and shoots development. So poultry litter is a valuable, natural soil amendment that adds nutrients and organic matter to increase soil fertility.

Probiotics are live microorganisms which confer a beneficial health benefit on the host. Trichoderma are model organisms to demonstrate influence on plant health as applied bio-fertilizer (Berg 2009). Trichoderma is considered as a probiotic having the characteristic of fungi and considered as a potential and promising bio-control agent and growth promoter for many crops. Bio-fertilizer is regarded as an eco-friendly substitute of chemical fertilizer trends to enhance soil fertility and increase crop productivity and yield without creating harm to the environment (Mihov and Tringovska 2010, Hermosa et al. 2012). Trichoderma has been a common component of bio-fertilizer which is a common inhabitant in soil and common symbiont of plant root (Sharma et al. 2012, Vinale et al. 2012). T. harzianum can solubilize several plant nutrients (Altomare et al. 1999) and T. asperellum has been shown to enhance the availability of P and Fe to plants, with significant increases in dry weight, shoot length and leaf area (Yedidia et al. 2001). Application of *Trichoderma* as a stand-alone agent, or in combination with compost or manure can significantly boost yields in rice and vegetables. Nahar et al. (2012) stated that soil applications of tricho-compost and tricho-leachate significantly increased the seedling germination rate and reduced the incidence of soil-borne diseases and infestation of rootknot nematodes and also reduced the seedling mortalities by 40.9% to 64.5% in Gazipur and 53.3% to 62.1% in Bogra, Bangladesh. It was established that Trichoderma is a biological degrader and promotes plant disease defense, increasing the immunity of the plant. Hence, Trichoderma have tremendous opportunities for disease management of soil borne pathogens but also have the capacity to improve plant growth parameters, and soil health.

Among different agricultural inputs fertilizer is the most important one and nearly 50% of the modern agricultural production depends on this insert (Pradhan 1992). It is authentic that fairly good soil fertility and plant nutrients are important to farming, whether the practices are considered "conventional" or "sustainable" (Hue and Silva 2000). But in agriculture conventional practices like excess tilling, injudicious application of readily soluble inorganic fertilizers and pesticides formulations have potentially damaged the environment.

To minimize environmental damages and promote economic feasibility agricultural producers have shown interest switching from "conventional" to "sustainable" agriculture. Although sustainable farming is often a much broader concept than "organic" farming, sustainable farming also emphasizes use of organic materials as soil amendments and sources of plant nutrients. Hence to improve soil fertility and increase plant nutrients availability, efforts need to be made to increase soil organic matter content. Therefore, the present research work has been carried out on the following objectives:

- i. To study the effect of soil amendments on physical and chemical properties of soil.
- ii. To analyze the impact of soil amendments for better crop growth environment.
- iii. To determine the effect of soil amendments on growth and development of wheat plants.
- iv. To determine the effect of soil amendments on yield, yield attributes and grain quality of wheat.

Chapter II

Review of Literature

2. REVIEW OF LITERATURE

Producing more crops per unit area is the present need of Bangladesh in order to obtain maximum return from the total cultivable land as well as to meet up the demand of food for the increasing population. Due to unrest cultivation of land, it loses its natural inheritances which affects crop yield. Soil fertility is a measure of the ability of soil to sustain satisfactory crop growth in the long-term, and can be determined by physical, chemical and biological processes intrinsically linked to soil organic matter content and quality. Effective management of organic matter in the field should conserve soil with minimal adverse effects on the environment. The information generated from different research works by many investigators on various aspects of soil amendments have been reviewed in this chapter.

2.1 Fertility situation of soil in Bangladesh

Though Bangladesh is a small country, it has a wide variety of soils. The status of fertility in Bangladesh soils is extremely variable. Most of the soils are depleted and in urgent need of replenishment with manure and fertilizer if productivity is to be enhanced. Nutrient supply ability of soil is known as fertility. It is a combined activity of physical, chemical and biological composition of the soil environment. In Bangladesh, the fertility of soil is deteriorating day by day (BARC 2005).

Vast population of Bangladesh pressured the land resources to derive necessary food, fiber and fuel over the last 2-3 decades. Expansions of modern crop varieties increase the land use intensification. This creates noticeable injury to the land qualities for non judicious withdrawal of plant nutrients by growing crops without proper replenishment. Land degradation occurs when land qualities are affected negatively due to natural causes or human interference. This is very much true for Bangladesh. Thus, plant nutrient deficiency/unbalance, soil organic matter depletion, soil erosion, soil compaction, flash flood, water logging etc., have emerged as vital problems as the result of irrational land use (Karim and Iqbal 2001).

2.1.1 Status of organic matter

Organic matter is called the 'storehouse of plant nutrients' and 'life force of the soil'. Organic matter plays a great role in successful crop production as well as soil fertility and crop productivity. In Bangladesh, mostly agricultural soils content low organic matter. A good soil should have at least 2.5% organic matter, but in Bangladesh most of the soils have less than 1.5%, and some soils even less than 1% organic matter (BARC 2005). Islam et al. (1994) found that organic matter ranged from 0.6 to 1.7% in 29 soil series from different regions of the country. Islam (1990) reported that at least 2% organic matter should be present in the soil for successful crop production. But, he observed that 90% soils of Bangladesh contained 0.5-1.0% organic matter. On the other hand, Akter (2011) reported that organic matter content in soils of eight, out of ten soil series namely Silmondi, Tejgaon, Chandra, Khilgaon, Halma, Brahmapra, Jamalpur and Sherpur gradually increased since 1991 having the values of 0.57-3.02% in 1991, 0.9-2.29% in 2006 and 0.91-3.02% in 2011.

In Bangladesh, it is true that high cropping intensity causes depletion of soil organic matter which declines the productivity of soils. Lack of organic recycling through addition of crop residues, animal waste, and other organic manures are the main reasons of low organic matter content in Bangladesh soils. Mia and Karim (1995) mentioned that 81% of the total biomass fuel is consumed for domestic cooking. FAO and UNDP (1994) reported that decreased organic matter leads to the degradation of soil physical properties including water holding capacity and reduced nutrient retention capacity leading to the lower release of nutrients from mineralization of organic matter. Over the 20 years, due to intensive cultivation, average organic matter content of the top soil has decreased by 20-46% presented in Table 1 (Appendix).

A pilot survey in Jessore and Kushtia revealed that 73% of the farmers faced yield decline leading to reduced soil fertility due to the cultivation of continuous cereals or increased cropping pressure (Saunders 1991). Eighty-eight percent of wheat growing farmers applied farmyard manure (FYM), of which 30% applied pure cow dung, and most farmers who applied FYM did so each year, mostly after wheat harvest. In 1998, CARE (Cooperative for American Relief Everywhere) interviewed the farmers of different regions of

Bangladesh and recorded their expressions about organic matter. They said that soils become more hard and compact, losing structure and water holding capacity for intensive inorganic fertilizer use. Farmers said that inclusion of organic matter increased yield, reduced production cost, improves crop growth, increased water holding capacity and improves soil structure. They also reported that farmers realized the addition of cow dung and other organic manure as also leaving crop residues increased the organic matter content of the soils.

There is no separate policy for supporting organic farming in Bangladesh. The national food policy of 2006 is the main document on food security in Bangladesh. Recently, the cabinet approved the food safety act, 2013 to save the people from adulterated and contaminated food. During recent days, Bangladesh Agricultural Research Institute (BARI) initiated the Participatory Guarantee System (PGS) for authentications. It will add more value to the consumer and organic promotion as well. PROSHIKA and Unnyan Dhara created organic consumer groups and sold their products directly to the farmers (Musa et al. 2015).

2.1.2 Selected essential nutrient status

Our life and food security is closely related with proper soil management. So, it is an urgent issue to identify the main constraints related to physical, chemical and biological nature and integrating them for improvement in effectiveness of input use, crop yield and quality, soil fertility and overall sustainability. In Bangladesh, soil fertility deterioration is a main constraint for higher crop production. Use of low external sources of nutrients and continuous cropping are also defined as the constraints to restore the soil nutrients. Thus depletion of soil fertility is mainly due to exploitation of land without proper replenishment of plant nutrients.

Ali et al. (1997) reported that on an average, the total carbon content has decreased by 11%, the total N by 12%, pH decreased by 4% and the exchangeable acidity increased by 30%. The exchangeable K content in soil has decreased by 31% and available P showed a decline of about 9% over a period of 27 years (1967-1995) in Bangladesh.

As the time advances, new nutrient deficiency arises in soil. Six mineral elements such as N, P, K, S, Zn and B are commonly deficient in Bangladesh soils. In the early 1980s, S and Zn deficiencies in rice were observed. But another study by Ali et al. (2012) presented, change in organic carbon and total N contents showed an increase in most of the layers in different soils of Ganges River Floodplain during the period between 1960s and 2010. Magnesium is reported to be deficient in Old Himalayan Piedmont Plain and Tista Floodplain soils (OFRD 1998). The chronological deficiencies of nutrients with time are shown in Table 2 (Appendix).

2.2 Organic amendments and soil properties

2.2.1 Non-leguminous crop residue incorporation and soil properties

Cassman et al. (1998) mentioned that crop residues are a potential source of organic matter and it needs efficient management. If crop residues are managed efficiently, it can increase organic matter as well as soil N. Addition of organic matter in soil can improve soil health, meets the demand for N. Since one-third of total rice plant N is in the straw, the N fertilizer requirements may be replaced partially by returning straw to the field. Munawar et al. (1990) mentioned that crop residues placed on the soil surface reduce water loss by evaporation due to their mulching effect. Walters et al. (1992) and Nyborg et al. (1995) mentioned, respectively that a crop residue is a vital source of conserving and sustaining soil productivity. It is the primary substrate for replenishment of soil organic matter (SOM). Upon mineralization, crop residues supply essential plant nutrients. Additionally, improvement of the soil physical and biological conditions and prevents soil degradation by residue incorporation.

Patra et al. (1992), Sidhu et al. (1995) and Samra et al. (2003) reported that regular addition of sufficient amounts of organic materials, such as crop residues to the soil leads to the maintenance of microbial biomass and improvement of soil fertility. Liu and Shen (1992) and Meelu (1994) mentioned that removal or burning of crop residues deteriorated soil physical properties, while incorporation of crop residues into the soil under rice-based cropping systems improved soil aggregation. Crop residues also play an important role in maintaining soil physical conditions (Prasad and Power 1991).

Beri et al. (1995) conducted field experiments in India on the rice-wheat cropping system that resulted in both nutrient contents and their availability increased with the incorporation of crop residues compared with their removal or burning. In an 11 year field experiment on a loamy sand soil in the Punjab, the incorporation of residues of both crops in the rice-wheat cropping system increased the total P, available P and K contents in the soil over the removal of residues. In another study over a 5 year period on a silt loam soil in Himachal Pradesh, the incorporation of rice straw in wheat caused a slight increase in the availability of P, Mn and Zn and a marked increase in the availability of K (Verma and Bhagat 1992). Sen and Jana (1998) undertook a field trial and opined that 65-70% of the total moisture was extracted by rice from the upper 30 cm soil layer. Incorporation of organic residues in soil holds water fairly tightly, increasing water holding capacity, soil porosity and helps the root respiration. Kumar and Goh (2000) stated that crop residues are an important source of organic matter that can be returned to soil for nutrient recycling, and to improve soil physical, chemical and biological properties.

Basic et al. (2004) and Lal (2005) showed the positive side, retaining residues on the surface through conservation tillage is important in reducing P losses, as this practice alleviates runoff and erosion losses. Zeleka et al. (2004) found a significant decrease in bulk density, an increase in macro plus meso-porosity, and a decrease in penetration and shear resistance in systems where maize residues were incorporated annually over 3 years than where residues were removed. Kharub et al. (2004) observed in a field experiment and reported that N and K uptake by crop was higher when rice straw was incorporated in the field alone and was highest when incorporated along with green manure.

Bonde et al. (2004) observed that all the organic residues significantly lowered the bulk density over the control. Among different organic residues, FYM recorded greater value of availability of nitrogen, phosphorus and potassium in soil as compared to other treatments (wheat straw and press mud compost treatments) including control, in the cotton-soybean cropping system. Mandal et al. (2004) determined that despite some advantages like killing of deleterious pests and clearing the piles before wheat planting, burning results huge losses of N (up to 80%), P (25%), K (21%) and S (4-60%), air pollution (CO₂ 13 t/ha) depriving soils of organic matter (SOM). This loss of SOM is one of the recognized threats

to sustainability. Soil physical properties *viz.* soil moisture, temperature, aggregate formation, bulk density and hydraulic conductivity affected by residue management practice. High siliceous content of rice crop residues have the potential of transforming electrochemical properties of acidic soils that reduces P fixation, improving base retention and increasing the soil pH.

Lal (2005) mentioned that globally the total crop residue production is estimated at 3.8 billion tons per year, of which 74% are from cereals, 8% from legumes, 3% from oil crops, 5% from tubers and 10% from sugar crops. Besides C, crop residues contain all mineral nutrients, the content of which varies among crop species depending on the fertility of the soil. Saddiq and Al-Amir (2011) conducted a study in fall 2009 at the College of Agriculture, Babylon University to determine effects of two types of organic matter on the physical properties of clay soil. The results showed a significant increase in both soil saturated hydraulic conductivity and soil porosity with the addition of chicken waste and rice straw. The increases were more pronounced with chicken waste than with rice straw and the values of the bulk density decreased significantly with organic matter addition to the soil. Dhar et al. (2014) observed the lowest soil pH (7.70) with soil incorporation of wheat straw and green manure in alluvial soil due to the production of organic acids during decomposition. Electrical conductivity of the soils was not significantly influenced by the incorporation of organics. Shahrzad et al. (2014) reported an increase in pH and electrical conductivity of soil with incorporation of crop residues. Chaudhary et al. (2017) conducted a long-term fertilizer experiment in rice-wheat cropping system at Punjab Agricultural University, Punjab on sandy loam soil revealed that the incorporation of straw + NPK increased total soil porosity (46.3%) and decreased the bulk density (1.42 Mg/m^3) up to 0-15 cm when compared to 100% NPK treated plots (43.1%, 1.51 mg/m³, respectively).

2.2.2 Leguminous residue incorporation and soil properties

Sharma et al. (2000) studied the effect of *Sesbania aculeata* green manure and mung bean (*Phaseolus radiatus* L.) residue incorporation on rice-wheat cropping system. They reported the soil chemical properties showed that *Sesbania* green manure and mung bean residue incorporation increased soil organic C over summer fallow by 0.105-0.135%,

Kjeldahl N by 0.01% and available P by 5.0-5.5 kg/ha and positive effect was observed in the *Sesbania* green manure and mung bean residue incorporated plots but not in summer fallow plots. Additionally, they concluded through incorporation practice of *Sesbania* green manure and mung bean residue interact positively in building-up of soil N. Sharma and Prasad (2008) stated that the return of rice and wheat residue can recycle up to 20-30% N absorbed by the crop. Native and applied N can temporarily immobilize for their wide C:N ratio. They opined that combining application of wheat straw with *Sesbania* green manure or mung bean residue increased cereal grain yield and agronomic N efficiency and generally improved negative apparent N balances.

Sing et al. (2008) carried out an experiment having sorghum and mung bean-lentil cropping system, for three successive years to assess the effect of mung bean residue incorporation on sorghum and succeeding lentil productivity along with different doses of phosphorus (0, 30 and 60 kg/ha) applied to these crops. When 60 kg P_2O_5 /ha was applied and mung bean residue incorporated then the result showed that sorghum grain yield increased significantly and available soil nitrogen, phosphorus, and organic carbon content were increased with incorporation of mung bean residue. Naeem et al. (2009) evaluated the efficacy of organic and inorganic fertilizers and mung bean residue under wheat-mung bean-wheat cropping system where mung bean was grown with basal dose of 25-60 kg N- P_2O_5 /ha was applied. After mung bean harvest, three residues management practice, i.e. R + (mung bean residue incorporated into soil), R- (mung bean residue removed) and F (fallow) were performed in the main plots. The result obtained microbial biomass C, N, mineralizable C and N from R+ compared with fallow increased by 33.7, 47.4, 21.4 and 32.2% at surface and 36.8, 51, 21.9 and 35.4% at sub-surface soil, respectively.

Singh et al. (2011) carried out a research in a mung bean/urd bean-wheat cropping system to assess the effect of different methods of legume residues management along with N fertilizer on succeeding wheat yield and soil properties. They analyzed collected data and reported significant improvement in the number of panicles/unit area with different residue management (10.2-20.9%) over residue removal. Moreover, residues incorporation presented higher soil available major nutrients *viz.* N by 24.6%, P by 11.5% and K by 18.5% over

initial levels and bulk density, particle density, percent pore space and water holding capacity (WHC) like soil physical properties were also improved in residues incorporated plots over residues removal plots. Buarach et al. (2014) conducted a study to determine the effect of tillage systems and soil organic amendments on rice growth, yield and carbon sequestration on paddy soil including two factors: the first factor was included conventional tillage; t_0 and minimum tillage; t_1 (tillage systems) and the second factor was included mung bean; p_0 , *Sesbania*; p_1 , sunhemp; p_2 and rice straw; p_3 (soil organic amendments). The results demonstrated that tillage systems with a mung bean amendment tend to increase the rice yield as also soil organic matter was the highest in sunhemp amendment in soil as followed by Sesbania, mung bean and rice straw, respectively.

Ghassan et al. (2017) conducted a pot trial to analyze the influence of mung bean residue on the properties of soil involving three levels of mung bean residue at three incubation periods (5% + 1 week, 5% + 2 weeks, 5% + 4 weeks, 10% + 1 week, 10% + 2 weeks, 10% + 4weeks, 15% + 1 week, 15% + 2 weeks and 15% + 4 weeks) and control treatment. The results showed that the values of calcium, potassium, magnesium, CEC, phosphorous, organic carbon, organic matter and carbon: nitrogen ratio were significantly (p < 0.05) higher than that of the control treatment and they concluded that mung bean residue is effective in increasing the fertility of the soil.

Farooq et al. (2018) examined the integrated effect of allelopathic residues and NPK fertilizer treatments including T_0 (control), T_1 (200-150-100 kg NPK/ha), T_2 (100-75-50 kg NPK/ha + mung bean straw 4 t/ha), T_3 (100-75-50 kg NPK/ha + rice straw 4 t/ha), T_4 (mung bean straw 8 t/ha) and T_5 (rice straw 8 t/ha) in wheat production under different water regimes on soil fertility. The result revealed that among fertilizer treatments, mung bean residue caused a greater increase in soil organic carbon, available nitrogen and available phosphorus, while there was a maximum percent increase in available potassium with T_1 (200-150-100 kg NPK/ha). But maximum wheat grain yield (30% and 33%) was obtained from the treatment T_2 (100-75-50 kg NPK/ha + mung bean straw 4 t/ha) during 2014-2015 and 2015-2016, respectively.

2.2.3 Cow dung incorporation and soil properties

Rahman (2001) reported that the inclusion of green manure in the cropping pattern and addition of cow dung resulted in a slight decrease in pH. There was a little increase in available P, K, S and Zn contents in soil after three crop cycles. Matsi et al. (2003) investigated the influence of liquid cattle manure on winter wheat (*Triticum aestivum* L.) germination, growth, and nutrient utilization by applying four treatments in the same plots in a 4 years field experiment with winter wheat: (i) application of 40 Mg/ha/yr liquid dairy cattle manure (wet weight basis) before sowing; (ii) single application of 120 and 26 kg/ha/yr N and P, respectively, as inorganic fertilizers before sowing; (iii) as in ii, but with split application of N, half the amount before sowing and the rest at tillering; and (iv) no fertilization. The results showed that application of manure did not affect seed germination, but resulted in a significant increase in dry biomass at the two growth stages and in grain yield and nutrient uptake, similar to the inorganic N and P fertilization.

Adeniyan et al. (2011) conducted a pot experiment to compare different organic manures with NPK fertilizer and results showed that application of 5 t/ha of each of the evaluated organic manures and 100 kg/ha NPK 15:15:15 fertilizer improved chemical properties of both acid and nutrient depleted soils compared with unfertilized soil. Cow dung application resulted in the highest pH levels of 6.37 and 6.50 in acid soil and nutrient depleted soil, respectively, while NPK fertilizer gave lowest pH levels of 5.28 and 5.74 for both soils. Moreover, application of different types of organic manures enhanced soil organic C, total N, available P, exchangeable K and CEC better than NPK fertilizer in both soils. Santillan et al. (2014) studied the effect of cow manure application on soil chemical properties and concluded that the sites with manure application, organic matter content increased significantly compared to the control plot (1.24 and 1.43%), the addition of manure for several years contributed to lower soil pH value.

Ram (2017) stated that the most important significance of cow dung and cow manure is to maintain the organic microbial and mineral micronutrient richness of soil and as a medicine for plants. Report made on cattle manure contains an average of 1.04 percent nitrogen, 0.15 percent potassium, 0.78 percent phosphorus and 32 other micronutrients having the characteristics of low nitrogen release and spread over time.

They further stated that manure applications result in increases in pH, water holding capacity, hydraulic conductivity, infiltration rates and soil amending properties of this great natural fertilizer are unbelievable.

2.2.4 Compost incorporation and soil properties

Prasad and Kerketta (1991) conducted an experiment to assess the soil fertility, crop production and nutrient removal under different cropping sequences in the presence of recommended doses of fertilizer and cultural practices along with 5 t/ha compost applied to the crops. There was an overall increase in organic C, an increase in total N (83.9%), available P (117.3%) and CEC (37.7%). Compost application also decreases soil bulk density (Park et al. 1995). Ouedraogo et al. (2001) assessed the impact of compost on improvement of crop production and soil properties where compost were applied at the rate of 0 and 10 Mg/ha and 5 and 0 Mg/ha. They found no significant difference in soil organic matter content between treatments receiving compost and no-compost. However, compost application increased soil cation exchange capacity (CEC) range 4 to 6 cmol (+)/kg and soil pH was also increased by the compost application. Sorghum yield tripled on the 10 Mg/ha compost plots and increased by 45% on the 5 Mg/ha compost plots, compared to no-compost plots.

Eghball et al. (2004) conducted a study to determine the residual effects of annual or biennial application of N and P-based compost and manure on corn. Results showed that residual effects of N and P-based compost and manure application on corn grain yield and N uptake remain at least one growing season, whereas the effects on soil properties were lasting longer. The residual effects of compost and manure increased significantly soil pH, electrical conductivity and plant available P and NO₃-N concentration. Gil et al. (2007) described an alternative approach for cattle manure management on intensive livestock farms in industrial-scale composting plants. The resulting compost was applied to a field to study the viability of applying this compost combined with a nitrogen mineral fertilizer as a replacement for the mineral fertilization conventionally used for maize (*Zea mays* L.). One year later, soil pH, OM content and CEC were higher with the compost treatment. Total P, K, Ca and Na concentrations in compost-amended plots were higher than in mineral-fertilized ones.

Tejada et al. (2009) carried out an experiment and mentioned that among the advantages of compost as soil amendment is its potential to maintain soil organic matter, foster nutrient availability and increase soil microbial population and activity, thus enhancing soil quality and fertility. Vengadaramana and Jashothan (2012) obtained soil from 10 different areas in Jaffna peninsula and evaluated the effect of organic fertilizers such as compost fertilizer and cow dung on the water holding capacity (WHC) of those soils. A significant difference ($p \le 0.05$) was observed on mean WHC of each soil sample with compost fertilizer and cow dung treated separately when compared to the control. Addition of compost fertilizer and cow dung separately increased the mean WHC of each soil sample. Cow dung doubly increased the WHC of each soil sample.

Desta (2015) evaluated the effects of organic and inorganic fertilizers on soil fertility for two consecutive years using maize under rain fed conditions. Experimental treatments included factorial combinations of three rates of N (0, 60 and 120 kg/ha), compost (0, 5 and 10 t/ha) and S (0, 15 and 30 kg/ha) fertilizers which were laid out in RCBD with three replications. In comparison to the initial soil, results showed that integrated application of organic and inorganic fertilizers improved soil total porosity, pH, OC, total N, CEC, available P and S by 31.8, 0.9, 58.1, 20.0, 3.1, 29.8 and 38.9%, respectively but decreased bulk density by 26.1% in 0-30 cm soil depth. The plots were treated with compost applied 10 t/ha and S 30 kg/ha had revealed the lowest bulk density and the highest total porosity, while combined application of N 120 kg/ha, compost 10 t/ha and S 30 kg/ha showed the highest total N, available P and S. The highest OC and CEC were recorded in plots treated with N 60 kg/ha along with compost 10 t/ha.

Zaki (2016) conducted an experiment and reported that after addition of compost to soil at two rates (5 and 10 t/fed) in combination with three rates of N fertilization (35, 50 and 70 kg N/fed). The obtained data indicated that increasing N fertilization from 35 to 70 kg N/ fed increased significantly straw, grain and 1000-grain weight of rice as well as the high yield values of straw 2.88 t/fed, grain 2.11 t/fed and 1000-grain weight 35.50 g was obtained under a high level of the N fertilization and compost rate (70 kg N/fed and 10 t/ fed, respectively).

2.2.5 Vermicompost incorporation and soil properties

Gaind and Nain (2007) investigated to evaluate the relative contribution of organic fertilizers (paddy straw, microbial inoculants and vermicompost) and inorganic fertilizers (urea and superphosphate) in improving pH, C, N, humus, microbial biomass, dehydrogenase, phosphatase, cellulase, b-glucosidase and xylanase activities of soil under wheat crop. In wheat, soil vermicompost fertilization resulted in the highest microbial biomass, available phosphorus, and nitrogen content. Its effectiveness in minimizing the alkalinity of soil was also recorded compared to other treatments as indicated by pH change.

Ramesh et al. (2009) studied the combination effect of different organic manures on the productivity of crops and soil quality in India. Four cropping systems involving soybean (*Glycine max* L.), durum wheat (*Triticum durum* Desf.), mustard (*Brassica juncea* L.), chickpea (*Cicer arietinum* L.) and isabgol (*Plantago ovata* Forsk) were tested with the cattle dung manure (CDM), poultry manure (PM), and vermicompost (VC) vis-à-vis mineral fertilizers. The organic manures were applied based on the N-equivalent basis and nutrient requirement of individual crops. At the end of the 3-year cropping cycle, application of organic manures improved the soil-quality parameters *viz.* soil organic carbon (SOC), soil available nutrients (N, P and K) and soil bulk density. The highest SOC and accumulated higher soil available N, P and K recorded from the cropping systems, soybean-durum wheat. Where the manures applied in different combinations improved the soil quality and produced the grain yields which are at par with mineral fertilizers.

Tharmaraj et al. (2011) carried out an experiment to study the impact of various worm products such as vemicompost, vermiwash and mixture of vermicompost and vermiwash on soil physico-chemical properties during the pot culture studies of samba rice. They collected soil sampling and plant growth measurements for two months, i.e. during initial and final stages. Distinct enhancement were found in vermicompost treated soil involving physical properties such as the pH, electrical conductivity (EC), porosity, moisture content, water holding capacity and chemical properties like nitrogen, phosphorous, potassium, calcium and magnesium, whereas the corresponding physical and chemical values in

control were minimum. The physical properties were improved in vermicompost treated soil such as water holding capacity, moisture content and porosity. The addition of vermicompost resulted in a decrease of soil pH.

Choudhary and Kumar (2013) stated that replenishing nutrients through organic sources is essential to maintain the soil health and sustainability in Eastern Himalayan Region, India which is organic by default. Keeping this in mind an experiment was laid out on randomized block design with six treatments *viz*. T₁: Vermicompost (VC; 2.5 Mg/ha), T₂: Poultry manure (PM; 1.25 Mg/ha), T₃: Swine manure (SM; 3.0 Mg/ha), T₄: Cow dung manure (CDM; 10.0 Mg/ha), T₅: Farmyard manure (FYM; 10.0 Mg/ha) and T₆: Control, and replicated three times to study the effect of applied organic nutrients on growth and yield of maize. When the crop was supplied with FYM followed by CDM physical indication like porosity, maximum water holding capacity (MWHC), field capacity (FC), bulk density (BD) and moisture releasing pattern was measured better. Chemical parameters like pH, Soil organic carbon (SOC), available nitrogen (N), phosphorus (P) and potassium (K) were recorded better on VC followed by PM over control.

2.2.6 Poultry manure incorporation and soil properties

Das et al. (1992) observed that application of poultry manure at the rate of 5 t/ha increased the uptake of Ca, Mg, K, Fe and also soil organic carbon. Obi and Ebo (1995) also reported that application of organic sources, i.e. poultry manure significantly increased soil organic matter content, total porosity, infiltration rate and hydraulic conductivity. Khanam et al. (2001) observed that the continuous use of chemical fertilizers accelerate the depletion of soil organic matter and impairs physical and chemical properties of soil in addition to micronutrient deficiencies. Poultry manure contains a high amount of secondary and micronutrients in addition to N, P and K. When poultry manures are applied to the field, it may supply sufficient amounts of S, Zn and B to meet up the demand for the growth of rice plants.

Agbede et al. (2008) conducted an experiment using the same plots in three years to determine the effect of poultry manure to sorghum at the rate of 7.5 t/ha. The poultry manure significantly (p > 0.05) reduced soil bulk density and temperature but increased

porosity and moisture content. The poultry manure increased significantly soil organic matter, soil and leaf N, P, K, Ca and Mg concentrations. Furthermore, plant height, leaf area and weight of roots, shoot and grain yield were significantly increased. The mean grain yield was increased by 39%. The poultry manure had a cumulative effect on soil properties, growth and yield parameters over the three years of study.

Ru et al. (2012) worked on the effect of different chicken manure rates on crop yields and the soil nutrients accumulation were carried out by a field plot experiment aimed at resolving the unreasonable disposition and disadvantage to the environment from poultry manures. The results showed that appropriate application rate of chicken manure significantly increased the yield of wheat. The soil organic matter, total nitrogen (N), total phosphorus (P), total Zinc (Zn) and nitrate (NO_3^-), available P and available Zn contents significantly increased with the increasing rates of the chicken manure.

2.2.7 Green manure incorporation and soil properties

Mandal et al. (2003) conducted a field experiment on rice (*Oryza sativa* L.) and wheat (*Triticum aestivum* L.) with objectives to study the influence of different green manure (*Sesbania rostrata, Sesbania aculeata, Vigna radiata*) residues and in combination with different levels of nitrogen (0, 60, and 120 kg N/ha) on physical properties, organic matter and total nitrogen contents of soil. They found that organic matter and total soil nitrogen concentrations was higher under green manure treated plots than summer fallow and magnitude of reduction in bulk density due to green manure over fallow was 0.03-0.07 t/m³ in 0-15 cm soil layer and 0.05-0.09 t/m³ in 15-30 cm soil layer during the growth of rice and wheat. Green manure also improved the soil physical environments and saturated hydraulic conductivity than fallow.

Sultani et al. (2007) studied physical properties of soil influenced by various green manure legumes (*Sesbania*, cluster bean and rice bean) and different P levels (0, 30, 60, 90 kg P_2O_5/ha). They noticed that highest fresh biomass was observed in *Sesbania* (23 t/ha) followed by cluster bean (19 t/ha) and lowest in rice bean (17 t/ha) as also green manure crops, on average reduced soil bulk density (5%) enhanced total porosity (8%), and

macro pores and large meso pores (28%). An increase (11%) in total soil porosity and available water (17%) was observed in plots and decrease (7%) in soil bulk density where *Sesbania* was incorporated as a green manure crop. The application of chemical phosphorus showed meager positive impact on various soil physical properties but did not significantly increase porosity or reduced bulk density.

Deshpande and Devasenapathy (2010) studied the effect of green manure and different sources of organic manures on yield and soil chemical properties of rice. They reported green manure incorporation along with poultry manure application resulted in higher soil available N, P and increased K uptake and higher N and P uptake and increased soil available K was recorded with green manure and poultry manure application. In another context, incorporation of green manure in situ, vermicompost and poultry manure decreased the soil pH and increased the organic carbon content of soil compared to all other combinations of treatments. Jat et al. (2012) conducted a field experiment at New Delhi during the Kharif and Rabi season of 2006-2007 and 2007-2008 with maize (Zea mays L.)wheat (Triticum aestivum L.) cropping sequence as control, Sesbania green manure, wheat straw and their combination to evaluate the influence of organic sources. The result showed that Sesbania green manure in combination with wheat straw followed by Sesbania green manure alone recorded significant improvement in system productivity (12.5 t/ha), protein content of maize (8.51%) and wheat (10.24%) and N, P and K uptake of both the crops and the system. Furthermore, addition of Sesbania green manure along with wheat straw improved the available N, P, K and organic carbon in soil to the tune of 25.3, 49.3, 5.9 and 11.9% over control, respectively.

Phullan et al. (2017) applied manures (control, farmyard manure, *Sesbania* and cluster bean) as main split and mineral fertilizer rates (control, 40-30, 60-45, 80-60, 90-70 and 120-90 kg $N-P_2O_5/ha$) as sub-split. The manures significantly influenced soil properties along with shoot dry weight, N, P and K uptake. Conversely, the rates of mineral fertilizers did not have any effect on soil properties, however, significantly enhanced the shoot dry weight and N, P and K uptake. The combined use of manures and mineral fertilizers had a significant effect on shoot P uptake.

Pawar et al. (2018) carried out an experiment to study the effect of various crop residues and green manure on soil properties and yield of cotton in salt affected soils of purna valley. The results of the experiment indicated that various green manure treatments significantly decreased the pH value over initial status, whereas electrical conductivity was increased under almost all the treatments over initial status. The bulk density and mean weight diameter was slightly improved with *Dhaincha* in situ green manure as compared with control. The application of in situ *Dhaincha* green manure significantly increased available N (249.90 kg/ha), P₂O₅ (30.46 kg/ha) and K₂O (459.20 kg/ha).

2.3 Probiotic inoculation and soil amendments

2.3.1 *Trichoderma* spp. inoculation for soil

Singh et al. (2010) applied *Trichoderma* multiplied culture (TMC) of *T. harzianum* strain Th 37 at the rate of 20 kg/ha on the stubbles at the ratoon initiation stage and observed increased the availability of nitrogen (N), phosphorus (P) and potassium (K) by 27, 65 and 44%, respectively. Some micronutrients *viz*. Cu, Fe, Mn and Zn were enhanced, respectively by 6, 100, 79 and 66% and a considerable increase in organic carbon (55%) with associated decrease in soil pH (6%). Guong et al. (2010) Evaluated the effect of organic amendment on improving soil properties with application of 10 t/ha sugarcane filter cake compost plus *Trichoderma* spp. and 20 t/ha of fresh *Tithonia diversifolia* in orange orchards. They used the above organic treatments with recommended inorganic fertilizer (250g N -200g P₂O₅ -120g K₂O/plant) to compare with usual farmer practice (628g N - 327g P₂O₅ -64g K₂O/plant). Soil analyses indicated that amendment with sugarcane filter cake compost (+ *Trichoderma* spp.) and fresh *Tithonia diversifolia* led to an increase soil organic matter content, available nitrogen and phosphorus, CEC, soil aggregate stability and reduced soil compaction (p ≤ 0.05) compared with the farmers usual practice.

Suparno et al. (2016) aimed to evaluate the effectiveness of *Trichoderma* spp. isolates from tidal swamp fields against *Rhizoctonia solani*. The result of the study showed that *Trichoderma* spp. isolates from tidal swamp fields were suspected as *Trichoderma harzianum* and the isolates were able to decrease plant disease intensity caused by *Rhizoctonia solani* about 7.4%. As well, *Trichoderma* isolates contribute to the plant growth

which are represented by the number of growing tiller per clump and plant height (12.2 cm) and increases the content of nitrogen, phosphor, potassium in soils and soil pH value. Tapke (2017) opined that the chemical form of pesticides and use of fertilizers is not only financially burdening the farmers but also reducing fertility of the soil. Natural farming which involves zero-budget agriculture practices is a better option than even organic farming production from a natural *Trichoderma* culture using solid state fermentation as well as in biofertilizer use in cotton, groundnuts, maize, caster chilies all are uses in natural *Trichoderma* bio-fertilizers. Several farmers are doing well by natural farming in contrast to those depending on fertilizers.

2.4 Chemical fertilizer and soil properties

Baque et al. (2006) conducted a pot trial under greenhouse condition in Bangladesh having three levels of potassium (low: 39.0 medium: 156 and high 312 kg/ha) and three levels of soil moisture, namely control (less than 25% depletion from the field capacity, FC), mild stress (more than37.5% depletion from FC) and severe stress (more than 50% depletion from FC). The result revealed that higher levels of K improved the dry matter production, yield and yield contributing characteristics of wheat with the irrespective levels of soil moisture. Besides this, uptake of N, P and K were also enhanced with the increasing levels of K especially under water stress condition.

Dubey et al. (2012) carried out complete soil analysis to establish the status of nutrients in district Sidhi during cropping season and after harvesting (for summer and *Kharif* 2010). The obtained result clearly showed that the application of recommended doses of N, P and K increased the crop yield and application of 100% NPK substantially improved the available NPK over its initial value thereby indicating significant contribution towards sustaining the soil health. However, the fertility of soil appears to be adversely affected due to the imbalanced use of nutrients, i.e. N, P or N alone indicates necessity of the use of balanced fertilizers for sustaining soil fertility and productivity of crops.

Zhong et al. (2014) conducted an experiment to determine the effects of N fertilizer applications on soil physical-chemical characters and maize grain yield. The results indicated that the soil bulk densities were increased, whereas the soil porosity, field capacity and values pH were decreased by N application at different stages. Furthermore, N application could increase the N contents of leaf and stem, whereas less or excess N application should not significantly improve maize yield.

Azizi et al. (2016) studied the effect of different levels of chemical fertilizers on soil health and yield of maize (*Zea mays* L.), where fertilizers were used in different levels as nitrogen (120, 90,50,0 kg/ha), phosphorus (60, 45,30,0 kg/ha), potassium (40, 30, 20, 0 kg/ha) are applied under sandy loam soil. The result showed that application inorganic fertilizers increased soil physical properties like bulk density and particle density, but some parameters of soil physical properties as pores space and EC decreased. It was also recorded that the application of chemical fertilizers in treatment T_1 (120 kg N/ha, 60 kg P/ha, 20 kg K/ha) improved nitrogen and available phosphorus was found more in T_5 (60 kg N/ha, 60 kg P/ha, 20 kg K/ha).

Singh et al. (2017) opined that continuous application of balanced chemical fertilizers either alone or in combination with FYM or lime for forty two years significantly improved available N, P and K, total carbon (CT) and labile carbon (CL) as compared to control and unbalanced use of fertilizers. They also stated that application of 100 percent NPK + FYM was more effective for increasing available N, P and K as compared to application of chemical fertilizers alone and continuous use of urea alone had the most deleterious effect on soil properties and productivity of both the crops.

Li et al. (2018) studied to explore the influence on soil physico-chemical properties under a 38-years long-term fertilization in a brown soil where soil samples (0-20 cm) were taken from the six treatments of the long-term fertilization trial in October 2016: no fertilizer (CK), N_1 (mineral nitrogen fertilizer), N_1P (mineral nitrogen and phosphate fertilizer), N_1PK (mineral nitrogen, phosphate and potassic fertilizer), pig manure (M_2), M_2N_1P (pig manure,

mineral nitrogen and phosphate fertilizer). The results showed that the long-term application of chemical fertilizers reduced soil pH value, while the application of organic fertilizers increased pH value.

2.5 Organic amendments for wheat growth and yield

2.5.1 Non-leguminous crop residue incorporation

Rautaray et al. (2003) mentioned that application of rice straw 30 days before transplanting resulted in a higher number of panicles and grains per panicle. However, application of rice straw alone reduced the number of tillers but increased when applied in combination with *Sesbania rostrata*. Gurpreet et al. (2007) evaluated the effect of various combinations of green manure (GM), wheat straw (WS), rice straw (RS), farmyard manure (FYM) and urea alone (control) on physical and hydraulic properties of soil in a rice-wheat experiment (1988-2001) in India. With addition of GM to all these treatments; FYM + GM (0.59%), WS + GM (0.60%), WS + RS + GM (0.64%) and GM (0.47%), organic carbon content further increased significantly. Increased OC content of the soil in turn improved its aggregation status, infiltration rate and decreased the bulk density, dispersion ratio and soil strength correspondingly. The differences in yield of rice were not significant among urea, GM and WS applied alone or in combination. However, the highest yield produced FYM + GM + urea combination.

Singh et al. (2010) noted that incorporation of crop residues decreased bulk density (BD) and increased infiltration rate, WHC (water holding capacity), microbial population, soil fertility as compared to no residue treatment. They also concluded that the residue incorporation with NPK fertilizer resulted in the highest yield, nutrient uptake, improved residual soil fertility and soil microorganism's status. Polthanee et al. (2011) were undertaken a research with treatments consisted of (1) rice straw incorporation into the soil, (2) rice straw combined with cattle manure, (3) rice straw combined with bio-extracted fertilizer. The significant effect on plant height was found from the application of different types of organic fertilizer combined with rice straw. At harvest, total above ground dry weight and

panicle number were significantly affected by the application of different types of organic fertilizer. The rice straw combined with cattle manure and bio-extract fertilizer gave the maximum panicle number. Maximum grain yields were recorded from rice straw combined with cattle manure but did not show any significant difference from the treatment of rice straw combined with cattle manure and bio-extracted fertilizer.

Dahri et al. (2018) assessed the influence of wheat straw incorporation and timing prior to seeding at 6 Mg/ha (S+), relative to no straw (S-), on maize (*Zea mays* L.) growth and yield parameters, as well as on soil characteristics. The soil organic matter showed a weakly significant ($0.05 \le p \le 0.06$) increase under straw amendment compared with S-. Some parameters like seedling emergence, plant height, cob length, the number of grain rows/plant and the number of grains/cob, as well as 1000-grain weight and yield were the highest under S+, and the lowest under S-.

Meena et al. (2018) conducted a field experiment at the 'Research farm' of Division of Agronomy, Indian Agricultural Research Institute, New Delhi, during 2014-2015 and 2015-2016 to study the growth parameter, yield and economies in maize-wheat-mung bean cropping system as influenced by crop residue and potassium management. During both the years, wheat was positively influenced by variable crop residue and potassium management at growth stages. The highest value of plant height, DMA, LAI, mean crop growth rate, net assimilation rate of wheat were noticed with treatment receiving fungal consortium incorporation which was statistically at par with crop residue incorporation. Similar results were also revealed for grain yield, straw yield and significant interaction grain yield in the mentioned years.

2.5.2 Leguminous residue incorporation

Bakht et al. (2009) evaluated the effects of residue retention, fertilizer N application and mung bean (*Vigna radiata*) on wheat in a mung bean-wheat sequence where associated treatments were- (a) crop residue retained (+ residue), (b) removed (- residue), (c) 120 kg N/ha applied to wheat, (d) 160 kg N/ha to maize and (e) no nitrogen applied under a

crop rotation wheat with maize or wheat with mung bean. The experimental outcome was crop residues incorporation increased the wheat grain yield on average by 1.31 times and straw yield by 1.39 times and application of N fertilizer produced on average 1.59 times more grain and 1.77 times more straw yield of wheat over the 0 N kg/ha treatment.

Ali et al. (2009) evaluated the suitability of different sources of organic materials for integrated use with chemical fertilizers considering the treatment T_1 : control, T_2 : 70% NPKS, T_3 : 100% NPKS, T_4 : 70% NPKS + rice straw (RS) 5 t/ha, T_5 : 70% NPKS + *Dhaincha* (DH) 15 t/ha, T_6 : 70% NPKS + mung bean residue (MBR) 10 t/ha, T_7 : 70% NPKS + cow dung (CD) 5 t/ha and T_8 : 70% NPKS + poultry manure (PM) 3 t/ha. In this study they applied organic manure or crop residue to transplant *Aman* rice and their residual effects were observed in the following *Boro* rice and reported that total grain yield in the cropping pattern ranged from 5.14 t/ha in T_1 (control) treatment to 12.29 t/hain T_3 (100% NPKS) treatment and application 3 t/ha PM with 70% NPKS (T_8) produced the total yield of 12.09 t/ha followed by 11.59 t/ha in the treatment containing 10 t/ha MBR plus 70% NPKS (T_6).

Jan et al. (2011) studied the response of mung bean residue (0, 10, 20 and 30 Mg/ha), nitrogen levels (0, 25, 50 and 75 kg/ha) and their interaction on barley. They observed that 30 Mg/ha mung bean residue presented values as emergence/m² (50), plant height (109 cm), leaf area/tiller (106 cm²), lodging score (5.55), termites attack (3.4%), grains/spike (67), biological yield (12.80 Mg/ha) and grain yield (2.32 Mg/ha) were significantly ($p \le 0.05$) higher compared to other levels. Similarly plant height (110 cm), lodging score (5.29) and biological yield (13.75 Mg/ha) were higher at 75 kg/ha N compared to other levels of N and interaction between residue and nitrogen indicated that 10 Mg residue and 50 kg N/ha is recommended to achieve maximum net return under comparable conditions.

Polthanee et al. (2012) conducted a research to investigate the effect of different rates of organic fertilizers with or without mung bean crop residues incorporated into the soil after harvesting, on growth and yield of transplanted rice. The result indicated that incorporation of mung bean residue into the soil provided 3.2 t dry matter/ha containing

50.2 kg N, 9.8 kg P and 166.2 kg K/ha along with significant increased plant height and tiller number/hill. They also found that incorporation of mung bean residue produced an increase in rice grain yield over fallow treatment of 416 kg/ha (or 17%) and significantly increased panicle number/hill.

2.5.3 Cow dung incorporation

Matsi et al. (2003) stated that liquid cattle manure should be applied to soil in such a manner that would improve soil fertility and crop production without causing salinity problems or increasing NO_2 levels. A four years experiment with winter wheat showed that application of manure did not affect seed germination but resulted in a significant increase in dry biomass at two growth stages and grain yield relative to control and was similar to that of fertilizer treatments.

Ramesh et al. (2009) reported combined effect of cattle dung manure (CDM), poultry manure (PM), and vermicompost (VC) *vis-à-vis* mineral fertilizers tested in four cropping systems involving soybean, durum wheat, mustard, chickpea and isabgol. They were applied organic manures based on the N-equivalent basis and nutrient requirement of individual crop and found the grain yields of durum wheat and isabgol were higher in the treatment that received from a combination of CDM + VC + PM, whereas in mustard, CDM + PM and in chickpea, CDM + VC recorded the higher yields. Hossain (2011) reported that cow dung significantly influenced on the characters related to the growth, yield and yield contributing characters except plant height, panicle length and 1000-grain weight. The highest grain yield (5.49 t/ha) was obtained from 8 t/ha cow dung application which was statistically identical with 6 t/ha cow dung application but significantly differ from cow dung rates. The lowest grain yield (4.59 t/ha) was obtained from control treatment.

Hossaen et al. (2011) studied the efficacy of different organic manure and inorganic fertilizer on the yield and yield attributes of *Boro* rice (*Oryza sativa* L.). The experiment consisted of 8 treatments, T₀: Control, T₁: 100% $N_{100}P_{15}K_{45}S_{20}$ (recommended dose), T₂: 50% NPKS + 5 t/ha cow dung, T₃: 70% NPKS + 3 t/ha cow dung, T₄: 50% NPKS + 4 t/ha poultry manure, T₅: 70% NPKS + 2.4 t/ha poultry manure T₆: 50% NPKS + 5 t/ha

vermicompost, T₇: 70% NPKS + 3 t/ha vermicompost. The highest number of effective tillers/hill (13.52), the longest panicle (24.59 cm), maximum number of total grain/plant (97.45), the highest weight of 1000-seeds (21.80 g), the maximum grain yield (7.30 t/ha) and straw yield (7.64 t/ha) was recorded from T₅, whereas the lowest number of effective tillers/hill (6.07), the shortest panicle (16.45 cm), the minimum total grain/plant (69.13), the lowest weight of 1000-seeds (16.73 g), the lowest grain yield (2.06 t/ha) and straw yield (4.63 t/ha) was observed from T₀.

Raj et al. (2014) mentioned that cow dung is very effective alternatives to chemical fertilizers by enhancing productivity in long term with maintaining the soil health and enhancing the microbial population. Cow dung manure and vermicompost increase soil organic matter content, and this leads to improved water infiltration and water holding capacity as well as an increased cation exchange capacity. It also enhances productivity of yield and minimizes the chances of bacterial and fungal pathogenic disease.

Taheri et al. (2017) conducted a field experiment to assess the impact of cow manure and vermicompost on the improvement of rice grain yield and quality in Rasht, Iran in 2015 and 2016. The experimental factors were as two cow manure (0, 10 and 20 t/ha) and vermicompost (0, 5 and 10 t/ha). The results revealed that the application of cow manure and vermicompost increased grain yield components such as the number of fertile tillers and the number of grain. The highest grain yield was obtained from the application of 30 t/ha cow manure + 10 t/ha vermicompost in the first year (3537 kg/ha) and in the second year (3958 kg/ha). Although the combined application of various rates of cow manure and vermicompost improved plant growth and nutrient uptake but the influence of vermicompost on the grain yield and quality was much stronger.

Khatun et al. (2018) carried out a field research to evaluate the growth, yield and yield attributes of aromatic rice (cv. Tulshimala) under the fertilization of cow dung (organic manure) and zinc (micronutrient). The increased the number of total tillers/hill, number of productive tillers/hill, panicle length (cm), grain yield/hill (g), straw yield/hill (g),

grain yield (t/ha), straw yield (t/ha) and biological yields were produced over control by the application of different levels of cow dung and zinc fertilizers. Here the treatment combination 10 t/ha cow dung and 12 kg/ha $ZnSO_4$ produced the highest grain yield (2.79 t/ha) and straw yield (5.80 t/ha) over other treatments.

2.5.4 Compost incorporation

Garcia et al. (2008) stated that application of compost is one of the most effective amendments and it improves not only the physico-chemical properties like aeration, cation exchange capacity, buffer capacity or porosity, but biotic factors too. They also mentioned from a biological point of view, compost can positively affect microbial population and their enzymatic activities and stimulates the development of plants by means of the presence of growth factors. Ibrahim et al. (2008) investigated the impact of organic manure and compost on productivity of wheat (*Triticum aestivum* L. cv. Inqlab-91) in sandy clay loam soil. They stated that organic amendments had positive but variable effects. They also observed that the organic manures/pot application increased the wheat yield by 11.13 (105%) to 13.53 (128%) g relative to the control. The wheat plant height, number of tillers, spike length, straw yield, grain yield and 1000-grain weight all were statistically different from that of control. They concluded with the suggestion that crop productivity may be improved significantly by the application of various organic manures for a longer time.

Diacono and Montemurro (2009) stated that adding up of organic residues recurrently, particularly the composted ones, increased soil physical fertility, mainly by improving aggregate stability and decreasing soil bulk density and also crop yield increased by up to 250% by long-term applications of high rates of municipal solid waste compost. Stabilized organic amendments improve crop yield quality but not reduce it. Saini and Kumar (2014) conducted a field experiment during 2006-2007 to 2012-2013 in India with combined application of composts in maintaining the soil health and ensuring sustainability in crop productivity. During conversion to organic farming, for the first three years of study (2006-2008) the yields of maize were low, however, after the third year of study i.e. from 2009-2012 the yields were improved. Similarly, in the wheat-gram cropping system the

wheat equivalent yields were low during 2006-2007 and 2007-2008 but thereafter, an impressive increase was observed due to the improvement of soil health in the form of nutrients and microbial status.

Imran et al. (2017) used organic sources of fertilization to attain sustainability in agricultural production and noticed that soil physico-chemical characteristics were greatly influenced by charcoal and compost combine application. They also obtained results from the study revealed that maximum number of tillers (331), maximum productive tillers (312), promising number of grain/spike (56) and 1000 grain weight (44.1 g) of wheat was recorded with combine application of compost and charcoal 5+5 Mg/ha, whereas highest biological yield (8710.2 kg/ha) and grain yield (4023.2 kg/ha) was noted with compost application at the rate of 10 Mg/ha as compared to control plot and other treatments. With charcoal application at the rate of 10 Mg/ha along with combined application of compost and charcoal and inorganic fertilizers presented the highest plant height (93.7 cm).

2.5.5 Vermicompost incorporation

Prasad et al. (2003) reported that application of organic manure mainly vermicompost at the rate of 5 t/ha was found beneficial and significantly increased the dry matter yield of maize. Dry matter production per plant, LAI and silking were significantly higher with the application of vermicompost at the rate of 5 t/ha along with poultry manure at the rate of 14 t and 10 t FYM/ha. Gowda et al. (2010) reported that application of vermicompost 3.8 t/ha + poultry manure 2.45 t/ha in wheat recorded significantly higher plant height (86.30 cm), number of leaves (60.10), 1000-grain weight (42.73 g) and seed yield (3043 kg/ha), vigor index (3223) and seedling dry weight (311.27 mg) compared to other treatments.

Gopinath et al. (2008) studied to evaluate the effects of three organic amendments on the yield and quality of wheat (*Triticum aestivum* L.) and on soil properties during transition to organic production. The organic amendments with composted farmyard manure (FYMC), vermicompost and lantana (*Lantana* spp.) compost-1 applied to soil at four application rates (60 kg N/ha, 90 kg N/ha, 120 kg N/ha and 150 kg N/ha). The grain yield of wheat from all organic amendments was markedly lower (36-65% and 23-54% less in the first

and second year of transition, respectively) than with the mineral fertilizer treatment. Protein content of wheat grain was higher (85.9 kg/g) for mineral fertilizer treatment in the first year of transition but in the second year, there were no significant differences among the mineral fertilizer treatment and the highest application rate (150 kg N/ha) of three organic amendments.

Barlas et al. (2018) aimed to investigate the effects of increasing doses of vermicomposts combined to soil and peat on wheat (*Triticum vulgaris* L.) growth and nutrition. For this purpose, air dried soil and peat were mixed with three rates of vermicompost equivalent to 0% (control), 25% and 50% (v/v) combinations which was replicated three times. The data revealed that nutritional concentration of aerial parts was influenced significantly by the application of vermicompost in the growth media. This study suggests that the vermicompost use in plant production has a role as a source of nutrients for plant growth.

2.5.6 Poultry manure incorporation

Umanah et al. (2003) conducted a field trial to study the effect of different rates of poultry manure on the growth, yield components and yield of upland rice cv. Faro 43 in Nigeria, during the 1997 and 1998 early crop production seasons. The treatments comprised 0, 10, 20 and 30 t poultry manure/ha. There were significant differences in plant height, internodes length, number of tillers/hill, panicle number/stand, number of grains/panicle and dry grain yield. There was no significant variation among treatments for 1000-grain weight.

Silva et al. (2005) determined the possibilities of increasing crop yields and soil nutrients by combined application of organic manure (straw, cattle manure, poultry manure and compost) and chemical fertilizer under rice crop rotation in 2004 yala and 2004-2005 maha seasons. Results of the study revealed that higher crop growth and yield can be obtained by combining organic manures and chemical fertilizers. Among the organic manure and chemical fertilizer combinations tested, poultry manure + NPK showed the highest (493% in yaha and 256% in maha) and rice straw + NPK combination showed the lowest (361% in yaha and 145% in maha) grain yields and increase of soil nutrient status, respectively.

They concluded that the combined application of poultry manure and chemical fertilizer is better compared to the organic manure + NPK combinations in sustainable crop yield and soil nutrient status.

Parvez et al. (2008) carried out an experiment at the soil science field laboratory, Bangladesh Agricultural University, Mymensingh during *Aman* season of 2006 to evaluate the combined effects of poultry manure, cow dung, *Dhaincha* and fertilization the yield and nutrient uptake by BRRI *Dhan* 30. The treatment receiving 80% RFD (recommended fertilizer dose) with 3 t/ha poultry manure produced the higher grain yield 4.80 t/ha and straw yield 5.70 t/ha. The performance of manure may be rated in order of PM>DH>CD.

Enujeke (2013) evaluated the different rates of poultry manure as 0 t/ha, 10 t/ha, 20 t/ha and 30 t/ha for growing improved maize in Nigeria. The results obtained indicated that plants that received 30 t/ha of poultry manure were superior at 8 weeks after sowing in 2008 and 2009, with mean height of 209.3 cm, mean number of leaves of 13.1, mean leaf area of 682.6 cm², mean grain weight at 16 weeks after sowing of 2.14 t/ha and mean number of grains/cob is 518.4. Islam et al. (2014) applied different treatments such as T_0 (Control), T_1 [STB-CF (HYG)], T_2 [CD + STB-CF (HYG)], T_3 [PM + STB-CF (HYG)], T_4 [COM + STB-CF (COM)] and T_5 [FP (Farmers' practice)] to observe integrated effect of manures and fertilizers on the yield attributes as well as grain and straw yields of wheat. The treatment T_3 [PM + STB-CF (HYG)] produced the highest grain yield 4362 kg/ha (90.40% increase over control) and straw yield 5492 kg/ha (84.79% increase over control). The treatment T_3 comprising poultry manure in combination with chemical fertilizers on IPNS (Integrated Plant Nutrient System) basis was found to be the best combination of manures and fertilizers for obtaining the maximum yield and quality of wheat at BAU farm.

Uwah et al. (2014) studied the effect of four rates of poultry manure (PM) (0, 5, 10 and 15 t/ha) on soil chemical properties and agronomic performance of sweet maize (*Zea mays* L. Saccharata strut). PM significantly ($p \le 0.05$) raised soil pH, organic matter (OM) content, total N, available P, exchangeable K, Ca, Mg and the effective cation exchange capacity (ECEC) status of soil. The 15 t/ha PM rate maximized sweet maize growth attributes,

total dry matter (TDM) and grain yields and also hastened days to 50% tasseling and silking. On average over the two seasons, the application of 5, 10 and 15 t/ha PM rates increased TDM and grain yield by 9.11, 57.29 and 77.67%; and 52.32, 117.18 and 144.28%, respectively compared with the 0 t/ha PM rate.

Rasul et al. (2015) investigated the influence of different manures on some vegetative growth of wheat variety *Semito* during the winter growing season of 2013-2014. The treatments including $T_1 = \text{control}$, $T_2 = 20$ t/ha sheep manure, $T_3 = 20$ t/ha cow manure and $T_4 = 20$ t/ha poultry manure; using randomized complete block design (RCBD) with four replications. The results showed that the poultry manure is the most efficient one compared to sheep and cow manures. The values of grain yield, biological yield and grain protein content were 6.75 t/ha, 15.67 t/ha and 14.96%, respectively for poultry manure treatment.

Shah et al. (2017) studied the residual effect of organic N (poultry manure) and mineral N on maize crop in Pakistan during 2014-2015. Combined doses of N from both sources were 120 kg/ha applied to wheat crops alone and in different combinations making six treatments. From the treatment where 25% N applied from poultry manure + 75% from mineral N source applied to the previous wheat crop produced maximum grain/ear, 1000-grain weight, biomass and grain yield was obtained.

Mukhtiar et al. (2018) worked on the use of different organic manures for improving crop productivity and to select potential organic manure that improve crop productivity. The results revealed that plots receiving poultry manures have high spikes/m² (274), grains/spike (60), more 1000-grain weight (42.29 g), high biological yield (11435 kg/ha) and high grain yield (3996 kg/ha). Similarly, sheep manure had also great effect on wheat parameters such as higher emergence/m² (103), tillers/m² (308), plant height (104.50 cm), grains/spike (60) and similar 1000-grain weight compared to poultry manures (42.20 g). They also compared cattle and farmyard manure with legume residue and found them better but were found less important in improving wheat quality compared to poultry and sheep manure.

2.5.7 Green manure incorporation

Rajinder et al. (2000) studied the combined use of organic manure and fertilizer on yield and nutrient uptake by rice in rice-wheat cropping system. They observed that yield of rice increased 100% with recommended NPK, but replacement of 25 or 50% N with green manure, FYM or wheat straw generally gave similar yields. Mammad et al. (2011) used a spring wheat cultivar in an experiment and included five organic manures; green manure (GM), farmyard manure (FYM), poultry litter (PL), press mud (PM) and sewage sludge (SS) at the rate of 10 t/ha. Six different treatments were made with different combinations of these manures along with one treatment having recommended dose of NPK (150, 115 and 60 kg/ha NPK, respectively) and one control treatment with no fertilizer at all. The results indicated that the combination of GM + PL+ SS each 10 t/ha gave maximum economic yield (3.65 t/ha), which was 137% more from control. PL and SS each 10 t/ha followed by green manure should be used as organic manure in wheat crops.

Paul et al. (2014) studied the effect of plant nutrient recycling through crop residue management, green manure and fertility levels on yield attributes, a rice-wheat cropping system in India. The crop residue incorporation (CRI) with or without *Sesbania* green manure (SGM) significantly influenced the plant height, number of tillers/m², number of grains/panicle or /ear and 1000-grain weight. Rice and wheat mean yield revealed that CRI or crop residue burning (CRB) resulted in slightly greater yield over crop residue removal (CRR) treatment. The CRI + SGM treatment again showed significantly greater grain yields of 7.54 and 5.84 t/ha and straw yields of 8.42 and 6.36 t/ha in rice and wheat, respectively, over other crop residue management treatments. This messaged that crop residue management with *Sesbania* green manure practice could be a better option for nutrient recycling to sustain the crop productivity in India, China, Pakistan and Bangladesh.

Seufert et al. (2012) reported that organic farming is a system aimed at producing food with minimal harm to ecosystems, animals or humans. Analysis of data shows that overall organic yields are typically lower than conventional yields. Context, systems and site characteristics creates yield variation for instant organic yields is 5% lower (rain-fed legumes and perennials on weak acidic to weak-alkaline soils), 13% lower (when best organic practices are used) and 34% lower (when the conventional and organic systems are most comparable).

2.6 Probiotic inoculation for wheat growth and yield

2.6.1 Trichoderma spp. inoculation for wheat

Sliesaravicius and Altintas (2006) accomplished a field trial with biological agent *Trichoderma* spp. and concluded that the treated plots produced the highest healthy grains compared to untreated plots. The highest 1000-grain weight was measured under the plots where seeds were treated either alone or in combination with *T. harzianum* followed by foliar spray. Up to 52 g (Shatabdi variety) 1000-grain weight was obtained from the plot which received seed treatment in combination with *T. harzianum* + *Penicillium* spp. and foliar spray with *T. harzianum*. They also investigated that *Trichoderma* spp. increased the spike length, number of grains/spike, 1000-grain weight and ultimately the grain yield.

Pratibah et al. (2012) conducted an experiment on the growth promoting ability of *Trichoderma viride* in the popular wheat (*Triticum aestivum* L.) variety Raj 3765 at farmer's field through TIFA-DST project and was successfully demonstrated in two districts of Rajasthan *viz*. Jaipur and Kota belong to different agro ecological zones (AEZ). Rhizosphere competence along with its growth promotion effect on rootlets, tillers, weight of grains and grain yield were evaluated by using *Trichoderma* spp. at three of the crops *viz*. seed, flowering and pre-harvesting at the rate of 4 g/kg and 4 ml/L along with soil treatment with farmyard manure and formulation at the rate of 50:1 before sowing and has been observed that grain yield significantly increased after application of *Trichoderma* spp. as compared to control.

Doni et al. (2014) evaluated the effectiveness of seven isolates of *Trichoderma* spp. in rice to promote growth and increase physiological performance. This study indicated that all the *Trichoderma* spp. tested isolates increased several physiological processes of rice including net photosynthetic rate, stomatal conductance, transpiration, internal CO_2 concentration and water use efficiency. They concluded that beneficial fungi isolate

Trichoderma spp. SL2 has potential growth promoting ability of rice regarding plant height (70.47 cm), tiller number (12), root length (22.50 cm) and root fresh weight (15.21 g) compared to the plants treated with other *Trichoderma* spp.

Bokhtiar et al. (2015) conducted a research at Bangladesh to examine the effects of enriched compost with chemical fertilizers on soil fertility and productivity of sugarcane where four levels of inorganic fertilizers (100%, 75%, 50% and 0%) and four levels of pressmud (enriched by *Trichoderma harzianum* 7.5 t/ha, *T. viride* 7.5 t/ha, untreated raw pressmud 10 t/ha and control) were applied. It was observed that germination (90.34%), total chlorophyll content (2.58 mg/g), leaf area index (5.00), dry matter (3.41 kg/m²), tiller (137.94 × 000 /ha), millable cane stalk (99.15 × 000/ ha) and yield (111.32 t/ha) were found maximum in 100% recommended fertilizer (N₁₅₀ P₅₀ K₉₀ S₃₅ and Zn₄ kg/ha) with enriched pressmud which was statistically identical in comparison with 75% and 50% of recommended chemical fertilizers with enriched pressmud. Finally, they concluded that enriched pressmud is more effective than raw pressmud in increasing sugarcane yield and maintaining soil fertility in High Ganges River Floodplain soils.

Hossain et al. (2016) worked with five different plant extracts including neem (*Azadirachta indica*), mehedi (*Lawsonia alba*), garlic clove (*Allium sativum*), rhizome of ginger (*Zingiber officinales*), seeds of black cumin (*Nigella sativa*), and BAU-Bio-fungicide (a *Trichoderma* based preparation) to evaluate the performance or effectiveness of those biological control agent on *Bipolaris* leaf blight of wheat and related pathogen (*Bipolaris sorokiniana*). Moreover, effect of seed treatment on wheat plant was evaluated and higher normal seedlings were found with BAU-Bio-fungicide, extracts of garlic clove and neem leaf at the value of 13%, 12% and 10%, respectively and BAU-Bio-fungicide also resulted 26.6% higher vigor index over control. They further reported that multiplication effect of seed treatment plus foliar spray showed superior effect by BAU-Bio-fungicide including higher 1000-grain weight (43.92 g) and grain yield (2.75 t/ha) which revealed that BAU-Bio-fungicide increased grain yield (29.87%) over control.

Hajieghrari and Mohammadi (2016) worked to observe the effect of five indigenous isolates (*T. harzianum* T 969, *T. harzianum* T 447, *T. hamatum* T 614, *Trichoderma* sp. isolate T and *Gliocladium virens* G525) on wheat variety (Moghan 3) seed germination, seedling vigor and plant growth through seeds treatment by the conidia and culture filtrate of the isolates. The result in overall, seed germination rate was increased and the highest seed germination rate (95.8%) was observed for *T. harzianum* T 969 non-sterilized cultural filtrate treatment. Furthermore, the field experiments showed that the isolates had significant effect on parameters such as 1000-grain weight, ear fresh and dry weights, ear length and stem and root dry weight.

Talukder et al. (2017) evaluated the efficacy of *Trichoderma* fortified compost with different substrates to reduce the pre-emergence and post-emergence seedling mortality, diseases of stem and root of chickpea. After the experiment, they stated that *Trichoderma* fortified compost with poultry manure was found significantly effective in reducing pre- emergence and post-emergence seedling mortality, disease incidence and disease severity of chickpeas in the field. Besides this, all the treatments significantly increased but *Trichoderma* fortified compost with poultry manure was the best to boost seed yield and quality.

Mahato et al. (2018) conducted a study to find out the effects of *Trichoderma viride* on growth and yield of wheat including of seven treatments (T) - T₁: Control, T₂: Soil + NPK, T₃: Soil inoculated *Trichoderma*, T₄: *Trichoderma* + FYM, T₅: *Trichoderma* + ¹/₂ NPK, T₆: *Trichoderma* + NPK and T₇ = *Trichoderma* + NPK + FYM. The results showed that *Trichoderma viride* increased the plant height (4.6%), root weight (1.5%), leaf length (0.3%), panicle weight (9.1%), number of grains (3.8%), grain yield (36.5%), biological yield (13.7%), and biomass yield (2.7%) over control. They also opined that when *T. viride* and NPK were accompanied with farmyard manure, most of the growth and yield parameters showed the highest value as more introducing farmyard manure to *T. viride* gives better yield than *T. viride* alone.

2.7 Chemical fertilizer for wheat growth and yield

Tarique and Paul (2005) carried out an experiment in order to evaluate the effects of different doses of nitrogen fertilizer on the yield of two wheat cultivars (Kanchan and Sourav) in the botanical field Rajshahi University, Bangladesh. In this study seven treatments were applied as $T_0 = \text{control}$, $T_1 = 20 \text{ kg N/ha}$, $T_2 = 40 \text{ kg N/ha}$, $T_3 = 60 \text{ kg N/ha}$, $T_4 = 80 \text{ kg N/ha}$ $T_5 = 100 \text{ kg N/ha}$ and $T_6 = 120 \text{ kg N/ha}$ which presented that plant height, dry matter/plant, tiller/plant, panicle length, spikelet/panicle, 1000-grain weight, harvest index and grain yield of wheat increased significantly with the increase of nitrogen level and the rate 120 kg N/ha considered as most suitable for the above mentioned characters.

Alam et al. (2009) conducted a research to ascertain the effect of K application on wheat at district Pabna in Bangladesh during *Rabi* season. In this study five different levels of K were applied and the 36 kg K/ha showed the remarkable increase in grain, straw and total biomass yield of wheat. Here the highest protein content was recorded from 36 kg K/ha, which was 6.86% and 4.98% higher over omission of K and recommended dose (100% estimated K). Iqbal et al. (2012) investigated that the effect of seeding rates and different levels of nitrogen on yield and yield component of wheat by using cultivar Fareed-2006 at seeding rates of 125, 150 and 175 kg/ha with five nitrogen levels as 0, 75, 100, 125 and 150 kg N/ha. They noticed that the plant height, number of tillers/m², spike length, number of spikelet/spike, 1000-grain weight, grain yield, biological yield and harvest index were highest at nitrogen at 125 kg N/ha and lowest at 0 kg N/ha level of nitrogen.

Debnath et al. (2014) conducted a field trial in Bangladesh Agricultural University, Mymensingh from November 2011 to March 2012 to determine the effect of nitrogen (N) and boron (B) fertilization on the performance of wheat. The trial comprised of four levels of N *viz*. 0, 80, 120, 160 kg/ha and three levels of B *viz*. 0, 1 and 2 kg/ha where grain was found to be significantly and positively correlated with number of effective tillers/plant, number of fertile spikelet/spike, number of grains/spike and straw yield. The result also presented that wheat grain yield increased with increasing levels of N and B up to 120 kg/ha and 2 kg/ha, respectively.

Akhtar et al. (2018) conducted a field experiment entitled "Yield maximization through nutrient management in irrigated wheat (*Triticum aestivum* L.)" comprising ten treatments of nutrient management. The experimental soil was clayey in texture, medium in available N, P, K and low in available zinc where wheat variety (GW-336) was sown at 22.5 cm row spacing. The experimental results revealed that significantly higher values of growth parameters *viz.* plant height, dry matter/plant, number of total tillers and effective tillers, and yield attributes *viz.* length of spike, number of grains/spike, grain weight/spike and 1000-grain weight, higher grain yield (4227 kg/ha) and straw yield (5792 kg/ha), quality parameters *viz.* protein content and protein yield were recorded significantly higher under the treatment of RDF (120-60-60 kg N -P₂O₅- K₂O/ha) + ZnSO₄ 25 kg/ha.

Chapter III

Materials and Methods

3. MATERIALS AND METHODS

The present piece of research work was conducted at Plant Biotechnology and Genetic Engineering Laboratory, Institute of Biological Sciences and its experimental field and at Plant Pathology Laboratory, Department of Agronomy and Agricultural Extension, University of Rajshahi during the period of July 2016 to June 2019 to study the sustainable improvement of wheat (*Triticum aestivum* L.) yield through soil amendments with probiotic and organic manures. In this chapter the details of different materials used and methodologies followed during the study are presented.

3.1 Description of the experimental site

3.1.1 Location

The experimental field is located at the western side of Rajshahi University, Bangladesh. Geographically situated at $24^{0}17^{\text{N}}$ N latitude and $88^{0}28^{\text{N}}$ E longitude at an elevation of 20 m above the sea level belonging to the Agro-Ecological Zone-11(AEZ-11).

3.1.2 Soil properties

The soil of the investigation plot has characteristics like poorly drained with moderately permeability, textural class loam and slightly alkaline in nature.

Moisture	PD	BD	Porosity	Sand	Silt	Clay	Textural class	
content (%)	(g/cc)	(g/cc)	(%)	(%)	(%)	(%)	Textural class	
19.40	2.65	1.27	51.34	47	40	13	Loam	

3.1.2.1 Physical properties

3.1.2.2 Chemical properties

pН	Organic matter (%)	K cmol (+)/kg	Total N (%)	P (µg/g)	S (µg/g)	Zn (µg/g)	EC (µs/cm)	C:N
8.10	1.20	0.150	0.07	26.30	12.50	0.75	145	10:1

3.1.3 Climatic condition

The field experimental site was under subtropical monsoon having heavy precipitation during the month of April to October and insufficient rainfall during the month of November to March. During the experiment period monthly average highest temperature $(28^{\circ}C)$ and precipitation (133 mm) was recorded in the month of April to September, whereas lowest temperature $(15^{\circ}C)$ and no precipitation found in December to January. Generally, winter is cool and dry in Bangladesh. Some meteorological records regarding temperature, rainfall, relative humidity and sunshine hours during the research period are presented in Table 4 (Appendix).

3.2 Cropping season

Rabi, *Kharif-I* and *Kharif-II* are the major cropping seasons in Bangladesh. *Rabi* season covers the month of middle October to mid March, *Kharif-I* season covers from mid March to the end of June and *Kharif-II* season from early July to mid October. Green manure (*Sesbania rostrata*) was grown in the *Kharif-II* season, whereas wheat in the *Rabi* season in this study.

3.3 Cropping history of the experimental plot

The plots of the research field are commonly used for agronomic crop production. Before 2016, the field was engaged with barley cultivation for two years.

3.4 Experiment

To achieve the objectives of the study the designed experiment was conducted for the three uninterrupted years.

3.4.1 Factors and treatments

The following three wheat varieties and nine soil amendments were included in the experiment:

Factor A. Wheat varieties (V): 3

- $V_1 = BARI$ wheat-28
- $V_2 = BARI$ wheat-29
- $V_3 = BARI$ wheat-30

Factor B. Soil amendments (T): 9

$T_0 =$	Control (without treatment)
$T_1 =$	Rice straw + vermicompost + green manure
T_2 =	Cow dung + vermicompost + green manure
T ₃ =	Compost + vermicompost + green manure
$T_4 \hspace{0.1in}=\hspace{0.1in}$	Poultry manure + vermicompost + green manure
T ₅ =	<i>Trichoderma harzianum</i> + vermicompost + green manure
T ₆ =	Mung bean residue + vermicompost + green manure
$T_7 \hspace{0.1 cm} = \hspace{0.1 cm}$	Trichoderma viride + vermicompost + green manure
T ₈ =	Chemical fertilizer (recommended dose)

3.4.2 Experimental design

The experiment was laid out in randomized complete block design (RCBD) with three replications. Each block was compacted with a 27 unit plot. Thus the total numbers of unit plots were 81. The unit plot was 2.5 m \times 2.0 m = 5.0 m² having plot to plot 0.5 m and bed to bed distance 0.25 cm and 1m from surrounding the boundary. The unit plots were separated with earthen bunds to avoid nutrient transfer to besides plot by lateral seepage.

Chapter III



BARI wheat-28



BARI wheat-29



BARI wheat-30



Green manure



Rice straw



Mung bean residue



Cow dung




Compost



Vermicompost



Poultry manure



Trichoderma spp.



Chemical fertilizer

Figs. 1 (a-l): Different treatment materials used in the experiment.

3.4.3 Description of wheat varieties

Three modern wheat varieties namely BARI wheat-28, BARI wheat-29 and BARI wheat-30 were used as planting materials. All the varieties were developed by Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur. Some features of these wheat varieties are given below:

Wheat variety ⇔ Wheat features ↓	BARI wheat-28	BARI wheat-29	BARI wheat-30
Sowing time	November	November	November
Plant height	95-100 cm	98-100 cm	100-103 cm
No. of tillers/plant	4-5	4-5	4-5
Days of heading	55-60	62-65	60-65
Grains/spike	45-50	42-43	45-47
Grain color	White	White amber	White amber
1000 seed weight	43-48 g	42-45 g	43-45 g
Life length	102-108 days	105-110 days	100-105 days
Grain yield	4.0-4.5 t/ha	4.0-5.0 t/ha	4.5-5.0 t/ha
Released on	2012	2014	2014

3.4.4 Green manuring

3.4.4.1 Land preparation

For experimentation during the year 2016-2017, the land was opened on 01 August, 2016 by power tiller followed by two ladders. Then weeds and stubbles were removed from the field and according to the experimental design 81 unit plots were prepared.

3.4.4.2 Seed sowing

As per treatment specification on 05 August 2016, seeds of *Dhaincha (Sesbania rostrata)* were sown on the experimental plots using at the rate of 50 kg/ha (BARI 2014) following broadcast method.

3.4.4.3 Harvesting

After 50 days of sowing the young succulent green manure plants were cut into pieces and incorporated into the soil to the respective plots (Dubey et al. 2015). After mixing, light irrigation was applied and waited about 6 weeks for decomposition of the plant materials. These similar practices were continued for the next two consecutive cropping seasons 2017-2018 and 2018-2019.

3.4.5 Establishment and management of wheat crop

3.4.5.1 Land preparation

After complete decomposition of green manure, the experimental plots were ploughed for three times followed by laddering to have a good tilt on 09 November 2016. Later on, weeds and stubbles were removed from the plots. For the experimentation the above work was repeated for next two years (2017 and 2018).

3.4.5.2 Treatment application

Crop residues (rice straw, mung bean residue), cow dung, compost and poultry litter were applied at the rate of 10 t/ha before 15 days of sowing. Other manures like vermicompost and *Trichoderma* spp. were applied as dose of 5 t/ha and suspension (10⁶ cfu/g at the rate of 5 kg/ha), respectively. Urea was applied at the rate of 200 kg/ha as three installments, first at final land preparation, second after first irrigation and third after second irrigation. All other fertilizers were applied as basal dose TSP 160 kg/ha, MP 45 kg/ha and Gypsum 115 kg/ha (BARI 2014) during final land preparation. Crop residues were collected from IBSc field laboratory. Cow dung and compost were collected from a trained farmers pit under DAE Bagatipara, Natore. Poultry litter was collected from a poultry farm. Vermicompost and chemical fertilizers were collected from Sumi Seed Vander, Rajshahi and *Trichoderma* spp. were cultured in Plant Pathology Laboratory, Agronomy and Agriculture Extension Department, University of Rajshahi, Bangladesh.

3.4.5.3 Collection of seed

Seeds of selected wheat varieties were collected from Regional Wheat Research Centre, Shyampur, Rajshahi, Bangladesh.

3.4.5.4 Seed sowing

In the 2016-2017 cropping year, wheat seeds were sown on 25 November 2016, likewise in the 2017-2018 and in 2018-2019 seeds were sown on 27 November 2017 and 25 November 2018. Seeds were sown in line following seed to seed and line to line distance 4 cm and 20 cm with a depth of about 4-5 cm opened by specially made an iron hand tine. The sown seeds were covered by soil manually. Seeds were applied as BARI recommended a dose 120 kg/ha. Bird boys were appointed for prevention from birds and other enemies during germination seeds and plant establishment.

3.4.5.5 Weeding

The major infesting weed species for wheat were *Chenopodium album*, *Cyperus rotundus* L., *Amaranthus spinosus* L. and *Cynodon dactylon* and weeds were controlled by hand weeding with niri for two times at 30 and 50 DAS.


3.4.5.6 Irrigation

Wheat crops were irrigated considering the presence of soil moisture. The crop provided 3 irrigations at crown root initiation (21 DAS), flowering (55 DAS) and grain filling (75 DAS) stages (BARI 2014). In every application flood method of irrigation was followed.

3.4.5.7 Harvesting and processing

After examining the proper maturity harvesting of crops was accomplished on 20 March 2017, 23 March 2018 and 25 March 2019, respective year of study. The harvested crop of each plot covering 1 m^2 was bundled individually, tagged properly and taken to the clean threshing floor. Then the crop was separately threshed, cleaned and winnowed for necessary data collection mentioned in the yield and yield contributing characters. Then grain yield and straw yield of each plot were recorded and the both yields were converted to hectare basis.

Chapter III




Prepared land



Seedling infection



Seedling blight



Sundry for oven dry




Oven dry



Booting stage



Panicle initiation



Crop at maturity



Bundle for data collection



Germinated seed



Soil collection



Data recording

Figs. 2 (a-l): Scenario of crop production and data collection.

3.4.6 Soil sample collection and analysis

3.4.6.1 Soil sample collection

Soil samples were collected randomly from nine places of the experimental field at the beginning of experimentation with the help of auger (SRDI 2017). Collected samples were packed in a polythene bag and were sent to the Department of Soil Science, Bangladesh Agricultural University, Mymensingh, for determination of moisture content, particle density and textural class.

3.4.6.2 Physical properties analysis

3.4.6.2.1 Moisture content

Moisture content of the soil is conveniently determined by gravimetric method. Soil samples (10 g each) were taken from a uniform depth. The fresh weight of the samples was recorded. Dry weight was determined after drying the soil in an oven for 24 hrs at 105°C to a constant weight, and the moisture percentage was calculated (Black 1965).

Percent of moisture =
$$\frac{W_1 - W_2}{W_1} \times 100$$

Where, W_1 = weight of sample before drying and W_2 = weight of sample after oven drying.

3.4.6.2.2 Particle density

Twenty five gram oven dry soil was poured in 100 ml volumetric flask and added 50 ml water in the flask. Then boiled the flask for 20 minutes in a water bath and cooled. After that distilled water was flask to have a desired volume. Then was reweighed the flask and measured the temperature of water to determine the density of water. The differences of two weights gave the weight of water. The volume of water was determined by dividing the weight of water by the density of water. Then volume of soil was recorded from subtracting the volume of water from 100. The particle density was then expressed by the formula (Balck 1965).

Particle density (Dp) =
$$\frac{W_s}{V_s}$$
 g/cc

Where, Ws = Weight of soil (g), Vs = Volume of solid soil (cc)

3.4.6.2.3 Bulk density and porosity

This method required a solid ring that pressed into the soil (Walter et al. 2016) to take an undisturbed core sample from 5-15 cm soil depth. Collected samples were dried at 105°C in an oven to get constant weights for 2-3 days depending on moisture content and then the mass of the dry soil sample was measured. The total volume of the soil sample was estimated as the internal volume of the core. The soil bulk density then was estimated as shown by the below formula (Yang et al. 2016).

Bulk density = $\frac{\text{Weight of oven dry soil}}{\text{Total volume of soil}}$ g/cc

Here, internal volume of core sampler (V) was calculated using the equation given below:

$$V = \pi r^2 h$$

Where, r is the radius of the core sampler and h is the height of the sampler.

The calculated bulk density to particle density and multiply by 100 gave the solid percentage of soil. Deduction of this result from 100 was presented the percentage of soil volume i.e. pore space or porosity as expressed by below formula:

Porosity (%) =
$$100 - \frac{\text{Bulk Density}}{\text{Particle Density}} \times 100$$

3.4.6.2.4 Textural class

Hydrometer method was used to determine soil texture by estimating the soil particles i.e. sand, silt and clay percent as per Bouyoucos (1936). Fifty grams of oven dry soil sample was separately taken in a dispersion cup and added 10 ml 5% calgon solution as also added 90 ml distilled water to the cup then kept overnight. The suspension was then stirred with an electrical stirrer for 10 minutes. The content of the dispersion cup was then transferred to 1 liter sedimentation cylinder and distilled water was added to make the volume up to the mark. A cork was placed on the mouth of the cylinder and the cylinder was inverted several times until the whole soil mass appeared in the suspension. The cylinder was set upright and the hydrometer reading was taken at 40 seconds and 2 hrs of sedimentation.

The temperature of the suspension was also recorded with a thermometer at 40 seconds and 2 hrs of sedimentation. The correction of hydrometer reading was made as the hydrometer was calibrated at 68°F. The percentage of sand, silt and clay were calculated as follows:

% (Silt + Clay) =
$$\frac{\text{Corrected hydrometer reading after 40 seconds}}{\text{Weight of soil}} \times 100$$

$$\% \text{ Clay} = \frac{\text{Corrected hydrometer reading after 2 hours}}{\text{Weight of soil}} \times 100$$

% Sand = 100 - % (Silt + Clay)

% Silt = % (Silt + Clay) - % Clay

The textural classes were determined by plotting the values of percentages of sand, silt and clay content on to the Marshall's Triangular Coordinate (Marshall 1947) following the USDA system.

3.4.6.2.5 Chemical analysis

For chemical analysis of soil samples were collected as mentioned in physical analysis. Here collected samples were air dried and powdered with a wooden roller. To remove coarse concretions, stones and organic debris each sample was passed through a 2 mm sieve. Then chemical properties mentioned pH, organic matter, N, P, K, Zn, EC and C:N was determined and analyzed at Soil Resource Development Institute (SRDI) Laboratory, Shyampur, Rajshahi, Bangladesh. Organic manures were also analyzed to determine nutrient elements content in the above mentioned institute shown in Table 3 (Appendix).

3.4.6.2.6 Soil pH

Soil pH was measured in 1:25 suspensions of soil and water on a glass electrode pH meter as outlined by Jackson (1962). Prior to making pH measurement, the electrode was calibrated using standard buffer solutions at pH 4.0 and 7.0

3.4.6.2.7 Organic matter

Organic carbon content of the soil was volumetrically determined by wet digestion method (Walkley and Black 1935). Organic matter was determined by multiplying organic carbon with Van Vemelon factor (1.724).

3.4.6.2.8 Total nitrogen

Total N in the soil was determined by the semi-micro Kjeldahl method. The sample was digested with concentrated sulphuric acid (H_2SO_4) and potassium sulphate (K_2SO_4) catalyst mixture (K_2SO_4 : CuSO₄.5H₂O: se = 100:10:1). Nitrogen in the digest was determined by distilling the digest with 10 N NaOH solution followed by titration of the distilled trapped in H_3BO_3 indicator solution with 0.01 N H_2SO_4 (Bremner and Mulvaney 1982).

3.4.6.2.9 Available phosphorus

Phosphorus was extracted by shaking the soil with 0.5 M sodium bicarbonate solution having pH 8.5. The extractable P in solution was then determined colorimetrically at 882 nm wave length after developing blue color using molybdate-ascorbic acid (Olsen and Sommers 1982).

3.4.6.2.10 Exchangeable potassium

Exchangeable potassium was extracted with neutral 1 N NH₄OAc as described by Jackson (1973) and was measured by Atomic Absorption Spectrophotometer.

3.4.6.2.11 Available sulphur

The available sulphur content in the soil was extracted by $0.01 \text{ M Ca}(\text{H}_2\text{PO}_4)_2$. The extracted S was estimated turbid metrically and the turbidity was measured by spectrophotometer at 420 nm (Page et al. 1982).

3.4.6.2.12 Available zinc

The available zinc content in the soil was extracted with 0.05 M HCl and the concentration of Zn in the extract were measured directly by Atomic Absorption Spectrophotometer at 214 nm (Page et al. 1982).

3.4.6.2.13 Electrical conductivity

The EC of collected soil samples was determined electrometrically (1:5: Soil: Water ratio) by a conductivity meter using 0.01 M KCl solution to calibrate the meter following the procedure described by Ghosh et al. (1983). After harvesting wheat crop soil samples were immediately collected on 20 March 2017 regarding nine treatments each sample were analyzed to obtain relevant physical and chemical properties which were defined before cultivation of green manure. The mentioned three years experimental plot soils physical and chemical properties analysis was done in the above institutions.

3.4.7 Crop characters studied

3.4.7.1 Seedling infection

The seedlings which were found yellow and rotted at the base are considered as infected. From germination up to 14 days regular observation was made and infected seedlings were counted in each plot. Infection was estimated by the following formula:

% Seedling infection = $\frac{\text{Number of infected seedlings}}{\text{Total number of seedlings}} \times 100$

3.4.7.2 Seedling blight

Seedlings which were dead and became straw in color defined as seedling blight. Seedling infection is treated as the first step of seedling blight. Thus, blighted seedlings were counted at 21 DAS in every plot. It was estimated as by the following formula:

% Seedling blight =
$$\frac{\text{Number of blighted seedlings}}{\text{Total number of seedlings}} \times 100$$

3.4.7.3 Leaf chlorophyll content

The atLeaf chlorophyll meter was used to assess leaf greenness. The meter determines light transmittance through the leaf at 660 and 940 nm wavelengths. The readings were obtained by inserting the middle portion of the topmost fully expanded leaf in the slit of the meter. At least three readings from leaves of randomly selected plants in each plot were recorded and mean value was determined. Abnormally looking or insect attacked plants were not selected for measurement (Ali et al. 2020).

3.4.7.4 Growth parameters of wheat

To explain the physiological basis of yield variation due to treatments, growth behavior of wheat was studied. Study of growth was started from 25 days after sowing (DAS) and ended at 85 DAS having a 15 days interval.

3.4.7.4.1 Total dry matter

In measurement of total dry matter (TDM) content, five plants were randomly collected from each plot at 25, 40, 55, 70 and 85 DAS. After collection of samples, the roots were cleaned with water to remove excess soil particles and kept in sun light to reduce moisture. The collected samples were then packed individually in labeled brown paper bags and were placed in an oven at 70-80°C for 72 hrs to get constant weight. After having constant weight each of them were measured with electrical balance and expressed as g/m^2 .

3.4.7.4.2 Leaf area index and crop growth rate

For determination of leaf area index (LAI) five plants were also collected randomly from each plot at 25, 40, 55, 70 and 85 DAS as stated above. At each time, collected plants were separated into leaf, stem and spike (when appeared) and leaf area of the collected leaves was measured by disc method. To determine leaf area every leaf of each plant was segmented into three portions and the length and breadth of the middle portion of each leaf were recorded. Among the total leaves of a plant, each leaf was packed including three segments individually in a labeled brown paper bag and was placed in the oven to get constant weight for 72 hrs at 70-80°C. Then leaf area was calculated by applying the following formula:

Area of leaf = $\frac{\text{Weight of leaves} \times \text{Area of segments}}{\text{Weight of segments}} \text{cm}^2$

Leaf area index (LAI) and crop growth rate (CGR) were calculated by following the standard formulae as shown below (Radford 1967).

Leaf area index (LAI) =
$$\frac{\text{Leaf area}}{\text{Ground area}}$$

Crop growth rate (CGR) = $\frac{(W_2 - W_1)}{(t_2 - t_1)} \text{g/m}^2/\text{d}$

Here, W_2 and W_1 are the total dry weight plant at different DAS and t_2 and t_1 are the harvesting time at the latter and former respectively.

3.4.7.5 Yield and yield contributing characters of wheat

3.4.7.5.1 Plant height

Randomly five plants were selected from the harvested crop regarding each plot and treatments to measure the plant height. The height of the plant (cm) was measured from the base of the plant up to the tip of the upper most tip of the spike.

3.4.7.5.2 Total plant/m²

Before harvesting 1 m^2 area was selected with quadrate and the number of total plants was counted. This operation was done in each plot preferably choosing the five rows in the middle of the plot.

3.4.7.5.3 Total tiller/plant

Tillers which had at least one leaf visible were counted from each plot that was collected with quadrate for representative area. It included both effective and non-effective tillers.

3.4.7.5.4 Effective tiller/plant

The numbers of tillers bearing panicles were counted at harvest with the help of quadrate from $1 \text{ m} \times 1 \text{ m}$ area, which expressed as effective tillers/plant.

3.4.7.5.5 Awn length

Awn length (cm) was measured with scale from the representative samples at the end portion of top spikelet to the apex of glum.

3.4.7.5.6 Spike length

Five spikes were selected randomly from each plot and their length was measured from the neck node of the rachis to the apex of last grain of each spike without awn length and average length of spikes was expressed in cm.

3.4.7.5.7 Spikelet/spike

Five spikes were selected randomly to count the number of spikelet/spike at harvest from each plot and data was presented as number of spikelet/spike by calculating the average.

3.4.7.5.8 Fertile spikelet/spike

Presence of any food material in the spikelet was considered as a fertile spikelet present on each spike were counted and average value five of spike were recorded as collected from each plot.

3.4.7.5.9 Grains/spike

Presence of any food material in the spikelet was considered as grain and the total number of grains present on each spike was counted. Later on mean was calculated.

3.4.7.5.10 Deformed grains/spike

Misshaped grains were regarded as deformed grains/spike. Deform grains/spike was separated from total grains/spike and then average value was recorded on plot basis.

3.4.7.5.11 Grain weight/spike

Grain weight/spike is cumulative estimation of grains/spike and deforms grains/spike. This value was also average, estimated from sampled spike and expressed in gram (g).

3.4.7.5.12 1000-grain weight

One thousand grains were counted from grains of harvesting crops grown in one square meter area at the center of each treatment plot, dried properly and weighed (g) by using an electric balance.

3.4.7.5.13 Grain yield

Grain yields were determined by harvesting crops grown in one square meter area of each plot. The harvested samples were then threshed, dried and weighed by using balance and finally the values were expressed in t/ha.

3.4.7.5.14 Straw yield

After separation of grains from plants, the straw obtained from one square meter area of each unit plot was sun dried and weighed by using balance and finally these values were converted into t/ha.

3.4.7.5.15 Harvest index

It denotes the ratio of grain yield (economic yield) to biological yield and was calculated with the following formula:

Harvest index (%) =
$$\frac{\text{Grain yield}}{\text{Biological yield}} \times 100$$

$$= \frac{\text{Grain yield}}{(\text{Grain yield} + \text{Straw yield})} \times 100$$

Here, Biological yield = grain yield and straw yield.

3.4.8 Seed quality parameters of wheat

3.4.8.1 Germination

3.4.8.1.1 Germination media

90 mm glass petri dishes were taken for germination tests. Two thirds of each petri dish was filled up with sands and made saturated by water and then leveled. Four hundred seeds were taken for each treatment for germination as recommended by ISTA (1985). Twenty five seeds were set in each petri dish for germination test. The media were kept saturated for 10 days by adding water at one day interval.

3.4.8.1.2 Determination of germination

Data on germination of wheat seeds were collected regularly up to 10 days and after 10 days total number of germinated seeds was counted. Percentage of germination was determined as follows:

Germination (%) =
$$\frac{\text{No. of germinated seedling}}{\text{No. of seed set for germination}} \times 100$$

3.4.8.2 Vigor index

After setting seeds in petri dishes data were collected regularly from the beginning to the end of the experiment. Germination percentage was calculated from the final count. Vigor index was found out by using following formula (Maguire 1962):

Vigor index =
$$\frac{\text{No. of normal seedlings (first count)}}{\text{Days to first count}} + - - - - - +$$


No. of normal seedlings (final count) Days to final count

3.4.8.3 Total soluble protein content of wheat seed

A technique formerly defined by Guy et al. (1992) was performed with some modification to estimate the entire soluble protein using spectrophotometer. Initially, wheat seeds were taken from the respected samples. Then the samples were weighed and washed with cold distilled water. Afterwards the samples were crushed with mortar and pestle using ice cold buffer containing 0.04% (v/v) 2-mercaptoethanol, mM EDTA (ethylene di-amine tetraacetic corrosive) and 50 mM Tris-HCl and buffer pH was fixed to 7.5. Then the sample solution was centrifuged for 12 minute at 10000 rpm at 25°C to remove the cell garbage. Afterwards the clear supernatants were separated in a glass cuvette and added 1 ml Coomassie Brilliant Blue (CBB) and measured the absorbance in a spectrophotometer (GENESYS 10S UV-Vis) at 595 nm. Finally, bovine serum albumin (BSA) calibration curve was applied to measure the total concentration of the sample's soluble protein.

3.4.9 Statistical analysis

The recorded data were compiled and tabulated for statistical analysis. The trial data for all morphological and yield parameters were analyzed statistically for analysis of variance (ANOVA) and Duncan's multiple range test (DMRT) (Gomez and Gomez 1984) was performed through r-studio (http://www.rstudio.com/) of "Agricolae" package (https://CRAN.R-project.org/package=agricolae). However, in case of mean comparison of soil parameters least significant difference (LSD) tests were used because of pairwise comparison such as 'Initial value versus 1st year value', '1st year value versus 2nd year value' and '2nd year value to 3rd year value'.



4. RESULTS

To achieve the objectives of the study various observations were completed. Some physical and chemical properties of soil and seedling viability, chlorophyll content, growth parameters, yield and yield contributing characters and seed quality of wheat were studied in this research. The collected data were analyzed statistically and presented in the form of tables, figures and appendices. Based on the above results detailed discussions are presented in this chapter.

4.1 Soil properties

4.1.1 Physical properties

4.1.1.1 Soil moisture

There was a significant influence of various soil amendments on soil moisture content in the year 2016-2017 (Table 1). Here the soil moisture content was found to be decreased significantly under all amendments except T₃. The highest rate of decrease (-3.33%) was found in the treatment T₈ (Chemical fertilizer). On the other hand, the least decrease in moisture content (-0.07%) was recorded in T₃ (Compost + vermicompost + green manure). As the consecutive year 2017-2018, soil moisture was varied by the application of different amendments. The treatment T₃ was demonstrated the higher increase of soil moisture (+0.60%) followed by T₂ (+0.40%), T₆ (+0.35%) and T₁ (+0.32%), whereas decline in soil moisture was found from T₀ (-1.27%) and T₈ (-1.80%). At the ending year of the study (2018-2019), the moisture content of soil was slightly increased for all the applied treatments except in T₀ (Control) and T₈. The most significant gain of soil moisture was recorded in T₃ (+0.87%) but major decline was found in T₈ (-2.13%) followed by T₀ (-2.00%).

Table 1: Changes of soil moisture content (%) as influenced by different soil amendments for three consecutive years of wheat production.

		2016-2017			2017-201	8	2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T ₀	19.40	16.73	-2.67*	16.73	15.47	-1.27*	15.47	13.47	-2.00*
T_1	19.40	17.85	-1.55*	17.85	18.17	+0.32 ^{ns}	18.17	18.65	+0.48 ^{ns}
T_2	19.40	18.65	-0.75*	18.65	19.05	$+0.40^{ns}$	19.05	19.77	+0.72*
T ₃	19.40	19.33	-0.07 ^{ns}	19.33	19.93	+0.60 ^{ns}	19.93	20.80	+0.87*
T_4	19.40	17.33	-2.07*	17.33	17.57	+0.23 ^{ns}	17.57	17.93	+0.37 ^{ns}
T ₅	19.40	17.10	-2.30*	17.10	17.27	+0.17 ^{ns}	17.27	17.53	+0.27 ^{ns}
T_6	19.40	18.05	-1.35*	18.05	18.40	+0.35 ^{ns}	18.40	18.95	+0.55 ^{ns}
T ₇	19.40	16.90	-2.50*	16.90	16.98	+0.08 ^{ns}	16.98	17.20	+0.22 ^{ns}
T_8	19.40	16.07	-3.33*	16.07	14.27	-1.80*	14.27	12.13	-2.13*
LSD (0.05)		0.41			0.71			0.69	
CV (%)		1.30			2.40			2.40	

4.1.1.2 Bulk density

In general, variation in bulk density of soil was recorded among the treatments in the whole study period (Table 2). There was a remarkable increase of soil bulk density under different soil amendments, the increase was pronounced (+0.05 g/cc) with T₈ treatment followed by T₀ (+0.03 g/cc). The decline trend was found in T₃ (-0.05 g/cc) followed by T₂, T₄ and T₆ in 2016-2017. Considering the differences between initial and final value of bulk density, the higher significant decrease was noted from T₃ with the value (-0.07 g/cc) oppositely higher significant increase was recorded from T₈ (+0.07 g/cc) during 2017-2018. Similar results were observed in the final year (2018-2019), where the greater significant deviation on bulk density was found in T₃ (-0.08 g/cc) but significant increase was found in T₈ (+0.10 g/cc) and in T₀ (+0.06 g/cc).

Table 2: Changes of soil bulk density (g/cc) as influenced by different soil amendments for three consecutive years of wheat production.

	2	2016-201	7	2	2017-201	8	2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T_0	1.27	1.30	$+0.03^{ns}$	1.30	1.34	$+0.04^{ns}$	1.34	1.41	+0.06*
T_1	1.27	1.24	-0.03 ^{ns}	1.24	1.21	-0.03 ^{ns}	1.21	1.17	-0.04 ^{ns}
T_2	1.27	1.23	-0.04*	1.23	1.20	-0.03 ^{ns}	1.20	1.15	-0.05 ^{ns}
T_3	1.27	1.22	-0.05*	1.22	1.16	-0.07*	1.16	1.08	-0.08*
T_4	1.27	1.24	-0.04*	1.24	1.18	-0.05 ^{ns}	1.18	1.14	-0.04 ^{ns}
T_5	1.27	1.24	-0.03 ^{ns}	1.24	1.21	-0.04 ^{ns}	1.21	1.17	-0.03 ^{ns}
T_6	1.27	1.24	-0.04*	1.24	1.20	-0.04 ^{ns}	1.20	1.15	-0.05 ^{ns}
T_7	1.27	1.25	-0.02 ^{ns}	1.25	1.24	-0.01 ^{ns}	1.24	1.20	-0.03 ^{ns}
T_8	1.27	1.32	+0.05*	1.32	1.39	+0.07*	1.39	1.49	+0.10*
LSD (0.05)		0.03			0.05			0.05	
CV (%)		1.30			2.30			2.50	

4.1.1.3 Particle density

The particle density of soil did not differ significantly by the activities of various soil amendments all over the research period (Table 3). From the records of soil analysis in 2016-2017, the particle density remained unchanged at the T₀ (+0.00 g/cc) but the lowest decrease (-0.01 g/cc) was reported by T₁, T₅ and T₇. The + 0.01 g/cc was the only positive value of particle density noticed in T₈. There were no significant changes of particle densities observed from the applied treatments. But the treatment T₀ (+0.01 g/cc) and T₈ (+0.02 g/cc) demonstrated slight enhancement over initial value in 2017-2018. Without major variation the particle density remained unchanged in 2018-2019, where T₃ had the minimum decrease of particle density (-0.02 g/cc) which had the equality with T₁, T₂ and T₅.

Table 3: Changes of soil particle density (g/cc) as influenced by different soil amendments for three consecutive years of wheat production.

	2	016-201	7		2017-202	18	2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T_0	2.65	2.65	+0.00 ^{ns}	2.65	2.66	+0.01 ^{ns}	2.66	2.67	+0.01 ^{ns}
T_1	2.65	2.64	-0.01 ^{ns}	2.64	2.63	-0.01 ^{ns}	2.63	2.61	-0.02 ^{ns}
T_2	2.65	2.63	-0.02^{ns}	2.63	2.63	-0.01 ^{ns}	2.63	2.61	-0.02^{ns}
T_3	2.65	2.63	-0.02^{ns}	2.63	2.61	-0.02^{ns}	2.61	2.61	-0.02^{ns}
T_4	2.65	2.63	-0.02 ^{ns}	2.63	2.61	-0.02 ^{ns}	2.61	2.60	-0.01 ^{ns}
T_5	2.65	2.64	-0.01 ^{ns}	2.64	2.63	-0.01 ^{ns}	2.63	2.61	-0.02^{ns}
T_6	2.65	2.63	-0.02 ^{ns}	2.63	2.61	-0.02 ^{ns}	2.61	2.60	-0.01 ^{ns}
T_7	2.65	2.64	-0.01 ^{ns}	2.64	2.62	-0.02 ^{ns}	2.62	2.61	-0.01 ^{ns}
T_8	2.65	2.66	+0.01 ns	2.66	2.68	$+0.02^{ns}$	2.68	2.70	$+0.02^{ns}$
LSD (0.05)		0.02			0.03			0.03	
CV (%)		0.40			0.60			0.70	

4.1.1.4 Soil porosity

All the applied soil amendments showed significant differences in respect of soil porosity in the experimental period 2016-2017 (Table 4). Here among the treatments, the maximum decrease of soil porosity (-0.96%) was obtained from T₈, though the maximum increase was reported by T₃ (+2.16%) and then T₂ (+1.92%). As the consecutive year 2017-2018, the treatments demonstrated increase in porosity except T₇ (-0.05%), T₀ (-1.36%) and T₈ (-2.24%). There the maximum porosity was obtained from T₃ (+2.31%) oppositely the minimum porosity was observed in T₁ (+0.99%). After three years of study, the significant increase of soil porosity (+2.83%) was recorded in T₃ but the most significant decrease (-3.40%) was noted in T₈.

Table 4: Changes of soil porosity (%) as influenced by different soil amendments for three consecutive years of wheat production.

	2	016-2017	7	,	2017-201	8	, ,	2018-2019	9
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T_0	51.34	50.94	-0.40*	50.94	49.58	-1.36 ^{ns}	49.58	47.37	-2.21*
T_1	51.34	52.91	+1.57*	52.91	53.90	$+0.99^{ns}$	53.90	55.00	$+1.09^{ns}$
T_2	51.34	53.26	+1.92*	53.26	54.29	$+1.03^{ns}$	54.29	55.93	$+1.64^{ns}$
T_3	51.34	53.50	+2.16*	53.50	55.80	+2.31*	55.80	58.74	+2.83*
T_4	51.34	52.98	+1.64*	52.98	54.73	+1.76*	54.73	56.33	$+1.63^{ns}$
T_5	51.34	52.82	+1.48*	52.82	54.15	$+1.33^{ns}$	54.15	54.98	+0.83 ^{ns}
T_6	51.34	53.04	+1.70*	53.04	54.29	$+1.25^{ns}$	54.29	55.81	$+1.52^{ns}$
T_7	51.34	52.84	+1.50*	52.84	52.79	-0.05 ^{ns}	52.79	53.81	$+1.02^{ns}$
T_8	51.34	50.38	-0.96*	50.38	48.14	-2.24*	48.14	44.74	-3.40*
LSD (0.05)		0.02			1.74			1.92	
CV (%)		1.20			2.00			2.20	

4.1.1.5 Soil sand particle

The changes in soil sand particles showed a decreasing trend due to practicing different organic amendments during 2016-2017 (Table 5). Although most of the treatments showed insignificant variation but the significant variation of sand particles occurred in T_8 (+1.33%). Furthermore, the decrease of sand particles was recorded by T_2 and T_3 . There was a similar trend of change of sand particles in the consecutive year 2017-2018. Here the maximum increase was noted from T_8 (+1.67%) and oppositely minimum from T_3 (-1.67%). Most of the values regarding sand particles of soil were affected by various soil amendments in 2018-2019. Likewise the previous year, T_8 was also increased (+2.00%) over the initial one followed by T_0 (+1.33%). Whereas the maximum decrease of sand particles were recorded from T_3 (-2.33%) followed by T_2 (-2.00%).

Table 5: Changes of soil sand particle (%) as influenced by different soil amendments for three consecutive years of wheat production.

		2016-201	7		2017-201	18		201	8-2019
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T ₀	47.00	48.00	$+1.00^{ns}$	48.00	49.33	+1.33 ^{ns}	49.33	50.67	+1.33 ^{ns}
T_1	47.00	46.33	-0.67 ^{ns}	46.33	46.00	-0.33 ^{ns}	46.00	45.00	-1.00 ^{ns}
T_2	47.00	46.00	-1.00 ^{ns}	46.00	44.67	-1.33 ^{ns}	44.67	42.67	-2.00*
T_3	47.00	46.00	-1.00 ^{ns}	46.00	44.33	-1.67*	44.33	42.00	-2.33*
T_4	47.00	46.67	-0.33 ^{ns}	46.67	46.00	-0.67 ^{ns}	46.00	44.67	-1.33 ^{ns}
T_5	47.00	46.67	-0.33 ^{ns}	46.67	46.00	-0.67 ^{ns}	46.00	45.00	-1.00 ^{ns}
T_6	47.00	46.33	-0.67 ^{ns}	46.33	46.00	-0.33 ^{ns}	46.00	44.67	-1.33 ^{ns}
T_7	47.00	46.67	-0.33 ^{ns}	46.67	46.33	-0.33 ^{ns}	46.33	45.33	-1.00 ^{ns}
T_8	47.00	48.33	+1.33*	48.33	50.00	+1.67*	50.00	52.00	+2.00*
LSD (0.05)		1.01			1.51			1.60	
CV (%)		1.30			2.00			2.10	

4.1.1.6 Soil silt particle

The influence of various soil amendments on soil silt particles was found to be increased in the year 2016-2017 (Table 6). The highest significant rate of increase was found in the treatment T_3 (+1.67%). On the contrary, the significantly highest decrease in silt particle (-2.00%) was recorded from T_8 . As the repeated year 2017-2018, the treatments T_3 (+2.00%), T_2 (+1.67%) and T_4 (+1.33%) were demonstrated the higher increase of soil silt particles. Where, the greater deviations in silt particles over initials were found from T_8 (-2.67%) and T_0 (-2.33%). At the final year of the research, the result presented that the silt particle of soil was slightly increased for all the applied treatments except T_0 and T_8 . The most significant achievement of the silt particle was recorded with T_3 having value +2.67%. But compared to initial value, significant decline of silt particles was found in T_8 (-3.00%) and T_0 (-2.67%).

Table 6: Changes of soil silt particle (%) as influenced by different soil amendments for three consecutive years of wheat production.

	2016-2017				2017-20)18		20	18-2019
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T ₀	40.00	38.67	-1.33*	38.67	36.33	-2.33*	36.33	33.67	-2.67*
T_1	40.00	41.00	+1.00 ^{ns}	41.00	42.00	+1.00 ns	42.00	43.67	$+1.67^{ns}$
T_2	40.00	41.33	+1.33*	41.33	43.00	+1.67 ^{ns}	43.00	45.33	+2.33*
T_3	40.00	41.67	+1.67*	41.67	43.67	+2.00*	43.67	46.33	+2.67*
T_4	40.00	40.67	$+0.67^{ns}$	40.67	42.00	+1.33 ^{ns}	42.00	44.00	+2.00*
T_5	40.00	40.67	$+0.67^{ns}$	40.67	41.67	+1.00 ns	41.67	43.00	$+1.33^{ns}$
T_6	40.00	41.00	$+1.00^{ns}$	41.00	42.00	+1.00 ns	42.00	44.00	+2.00*
T_7	40.00	40.33	$+0.33^{ns}$	40.33	41.00	+0.67 ns	41.00	42.00	$+1.00^{ns}$
T_8	40.00	38.00	-2.00*	38.00	35.33	-2.67*	35.33	32.33	-3.00*
LSD (0.05)		1.20			1.85			1.91	
CV (%)		1.80			2.80			2.80	

4.1.1.7 Soil clay particle

In general, insignificant variation of clay particles of soil was recorded among the treatments in 2016-2017 (Table 7). There was a decrease of clay particles under different soil amendments but the increase was pronounced with T_8 (+0.67%) and in T_0 (+0.33%). The treatment T_3 showed the maximum decrease (-0.67%) on this parameter. Considering the differences between initial and final value of clay particles, the higher increase was noted from T_8 and T_0 with the equal value +1.00% during 2017-2018. Whereas the maximum decrease of clay particles was recorded in T_1 , T_4 and T_6 with the value -0.67%. Similar results with insignificant change rate were observed in final year compare with the second year, where the greater difference on clay particle was found in T_1 , T_4 and T_6 having the similar value -0.67% though parallel increase was noted from T_8 (+1.00%).

Table 7: Changes of soil clay particle (%) as influenced by different soil amendments for three consecutive years of wheat production.

Treatment		2016-201	7		2017-20	18	2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T ₀	13.00	13.33	+0.33 ^{ns}	13.33	14.33	$+1.00^{ns}$	14.33	15.67	+1.33 ^{ns}
T_1	13.00	12.67	-0.33 ^{ns}	12.67	12.00	-0.67 ^{ns}	12.00	11.33	-0.67 ^{ns}
T_2	13.00	12.67	-0.33 ^{ns}	12.67	12.33	-0.33 ^{ns}	12.33	12.00	-0.33 ^{ns}
T_3	13.00	12.33	-0.67 ^{ns}	12.33	12.00	-0.33 ^{ns}	12.00	11.67	-0.33 ^{ns}
T_4	13.00	12.67	-0.33 ^{ns}	12.67	12.00	-0.67 ^{ns}	12.00	11.33	-0.67 ^{ns}
T_5	13.00	12.67	-0.33 ^{ns}	12.67	12.33	-0.33 ^{ns}	12.33	12.00	-0.33 ^{ns}
T_6	13.00	12.67	-0.33 ^{ns}	12.67	12.00	-0.67 ^{ns}	12.00	11.33	-0.67 ^{ns}
T_7	13.00	13.00	$+0.00^{ns}$	13.00	12.67	-0.33 ^{ns}	12.67	12.67	$+0.00^{ns}$
T_8	13.00	13.67	$+0.67^{ns}$	13.67	14.67	$+1.00^{ns}$	14.67	15.67	$+1.00^{ns}$
LSD (0.05)		1.54			2.33			2.24	
CV (%)		7.20			11.00			10.60	

4.1.1.8 Soil textural class

Textural class is output of the combination of soil particles in a particular soil. This parameter is a vital remark of soil type which affected the selection of soil for its utilization. The initial composite soil texture was reported as loam by analysis at the beginning of the study. The application of different soil amendments did not change the textural classes in the consecutive three years study (Table 8). Though it had no differences in the textural class all over the study period, there were variations in the percentage of different aforesaid soil particles.

		201	6-2017			201	7-2018			201	8-2019	
Treat.	Sand (%)	Silt (%)	Clay (%)	Textural class	Sand (%)	Silt (%)	Clay (%)	Textural class	Sand (%)	Silt (%)	Clay (%)	Textural class
T ₀	48.00	38.67	13.33	Loam	49.33	36.33	14.33	Loam	50.67	33.67	15.67	Loam
T_1	46.33	41.00	12.67	Loam	46.00	42.00	12.00	Loam	45.00	43.67	11.33	Loam
T_2	46.00	41.33	12.67	Loam	44.67	43.00	12.33	Loam	42.67	45.33	12.00	Loam
T ₃	46.00	41.67	12.33	Loam	44.33	43.67	12.00	Loam	42.00	46.33	11.67	Loam
T_4	46.67	40.67	12.67	Loam	46.00	42.00	12.00	Loam	44.67	44.00	11.33	Loam
T_5	46.67	40.67	12.67	Loam	46.00	41.67	12.33	Loam	45.00	43.00	12.00	Loam
T_6	46.33	41.00	12.67	Loam	46.00	42.00	12.00	Loam	44.67	44.00	11.33	Loam
T_7	46.67	40.33	13.00	Loam	46.33	41.00	12.67	Loam	45.33	42.00	12.67	Loam
T ₈	48.33	38.00	13.67	Loam	50.00	35.33	14.67	Loam	52.00	32.33	15.67	Loam

Table 8: Changes of soil textural class as influenced by different soil amendments for three consecutive years of wheat production.

 T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer.

4.1.2 Chemical properties

4.1.2.1 Soil pH

Soil pH was not significantly influenced by various soil amendments during the research period (Table 9). The increment of soil pH was noted higher (+0.13 unit) in T_8 and the (+0.10 unit) in T_4 in 2016-2017. While the least decrease (-0.07 unit) of pH was observed in T_2 and T_6 and remains unchanged in T_1 , T_5 and T_7 . In the year 2017-2018, the pH was decreased by most of the treatments except T_0 , T_4 and T_8 . There the bigger increase (+0.13 unit) of pH was found in T_8 but the minimum equal losses (-0.03 unit) were noticed in T_5 and T_7 . Although there was insignificant variation among the treatments in 2018-2019, however, the treatment T_8 was maintained the highest increment (+0.17 units). On the other hand, the maximum decrease (-0.10 unit) was recorded from T_2 and T_3 .

Table 9: Changes of soil pH as influenced by different soil amendments for three consecutive years of wheat production.

Treatment		2016-201	7		2017-20)18	2018-2019		
Treatment -	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change
T ₀	8.10	8.17	+0.07 ns	8.17	8.20	+0.03 ^{ns}	8.20	8.27	+0.07 ^{ns}
T_1	8.10	8.10	$+0.00^{ns}$	8.10	8.03	-0.07 ^{ns}	8.03	7.97	-0.07 ^{ns}
T_2	8.10	8.03	-0.07 ^{ns}	8.03	7.93	-0.10 ^{ns}	7.93	7.83	-0.10 ^{ns}
T_3	8.10	8.00	-0.10 ^{ns}	8.00	7.87	-0.13 ^{ns}	7.87	7.77	-0.10 ^{ns}
T_4	8.10	8.20	$+0.10^{ns}$	8.20	8.27	$+0.07^{ns}$	8.27	8.30	+0.03 ^{ns}
T_5	8.10	8.10	$+0.00^{ns}$	8.10	8.07	-0.03 ^{ns}	8.07	8.00	-0.07 ^{ns}
T_6	8.10	8.03	-0.07 ^{ns}	8.03	7.93	-0.10 ^{ns}	7.93	7.87	-0.07 ^{ns}
T_7	8.10	8.10	$+0.00^{ns}$	8.10	8.07	-0.03 ^{ns}	8.07	8.03	-0.03 ^{ns}
T_8	8.10	8.23	$+0.13^{ns}$	8.23	8.37	+0.13 ^{ns}	8.37	8.53	+0.17 ^{ns}
LSD (0.05)		0.15			0.21			0.17	
CV (%)		1.10			1.60			1.30	

4.1.2.2 Soil organic matter

Most of the cases soil organic matter content was improved by different soil amendments in the season 2016-2017. Significantly the major deviation (-0.12%) was noticed in T_8 then in T_0 (-0.10%), while major accumulation was recorded in T_3 (+0.04%) followed by T_2 (+0.03%). There was a trend of enhancement of soil organic matter by most of the amendments in 2017-2018. Where the greater significant increase (+0.06%) of organic matter content was observed in T_3 and considering the same parameter, the maximum significant decline (-0.15%) was noted from T_8 . Organic matter content of soil after completion of three years trial under different soil amendments followed a similar movement as in 2017-2018. The treatment T_3 greatly improved organic matter status of soil (+0.14%) but T_8 showed significant decline (-0.17%) of organic matter content of soil (Table 10).

Table 10: Changes	of soil organic	matter content (%) as influenced	by different soil
amendme	ents for three cons	secutive years of w	heat production	

Treatment -	2016-2017				2017-2018			2018-2019		
	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T ₀	1.20	1.10	-0.10*	1.10	0.98	-0.12*	0.98	0.85	-0.13*	
T_1	1.20	1.21	$+0.01^{ns}$	1.21	1.25	$+0.04^{ns}$	1.25	1.31	+0.06*	
T_2	1.20	1.23	$+0.03^{ns}$	1.23	1.28	$+0.05^{ns}$	1.28	1.39	+0.11*	
T_3	1.20	1.24	$+0.04^{ns}$	1.24	1.29	+0.06*	1.29	1.43	+0.14*	
T_4	1.20	1.22	$+0.02^{ns}$	1.22	1.27	$+0.05^{ns}$	1.27	1.36	+0.09*	
T_5	1.20	1.21	+0.01 ^{ns}	1.21	1.24	$+0.03^{ns}$	1.24	1.28	$+0.04^{ns}$	
T_6	1.20	1.22	$+0.02^{ns}$	1.22	1.26	$+0.04^{ns}$	1.26	1.33	+0.08*	
T_7	1.20	1.20	$+0.00^{ns}$	1.20	1.21	$+0.01^{ns}$	1.21	1.25	$+0.03^{ns}$	
T_8	1.20	1.08	-0.12*	1.08	0.93	-0.15*	0.93	0.76	-0.17*	
LSD (0.05)		0.05			0.05			0.04		
CV (%)		2.40			2.70			1.80		

4.1.2.3 Total nitrogen

At the beginning of the study, i.e. in 2016-2017, due to application of different soil amendments resulted decrease of total nitrogen content of the experimental plots. The maximum significant loss of nitrogen (-0.025%) was observed in T_0 but a minor non-significant decrease was noticed in T_5 (-0.001%). Negligible differences of total nitrogen over initial were recorded by most of the treatments in 2017-2018. The treatment T_4 exhibited the greater significant increase of nitrogen (+0.016%), while T_0 (-0.015%) and T_8 (-0.011%) had the decreasing trend. In the last year, treatments T_4 and T_3 were showed the maximum significant gain in total nitrogen with the value +0.013% and +0.010%. Furthermore, soil total nitrogen was reduced (-0.015%) under the treatment T_8 (Table 11).

Table 11: Changes of soil total nitrogen (%) as influenced by different soil amendments for three consecutive years of wheat production.

Treatment		2016-20	17		2017-2018			2018-2019		
Heaunent	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T_0	0.070	0.045	-0.025*	0.045	0.031	-0.015*	0.031	0.022	-0.010*	
T_1	0.070	0.051	-0.019*	0.051	0.054	+0.001 ^{ns}	0.054	0.056	+0.003 ^{ns}	
T_2	0.070	0.057	-0.014*	0.057	0.061	+0.005*	0.061	0.068	+0.007*	
T_3	0.070	0.063	-0.008*	0.063	0.069	+0.007*	0.069	0.079	+0.010*	
T_4	0.070	0.068	-0.002 ^{ns}	0.068	0.084	+0.016*	0.084	0.097	+0.013*	
T_5	0.070	0.070	-0.001 ^{ns}	0.070	0.076	+0.006*	0.076	0.084	+0.008*	
T_6	0.070	0.056	-0.015*	0.056	0.058	+0.003 ^{ns}	0.058	0.062	+0.005*	
T_7	0.070	0.066	-0.004*	0.066	0.070	$+0.004^{ns}$	0.070	0.075	+0.005*	
T_8	0.070	0.050	-0.020*	0.050	0.039	-0.011*	0.039	0.024	-0.015*	
LSD (0.05)		0.003			0.004			0.003		
CV (%)		3.100			3.800			2.700		

4.1.2.4 Exchangeable potassium

In 2016-2017, exchangeable potassium (K) was significantly influenced under various soil amendments (Table 12). Data revealed that K was distinctly depleted under all the treatments with the maximum depletion in T_0 (-0.063 cmol (+)/kg) and then T_8 (-0.045 cmol (+)/kg). In the second year (2017-2018), there was an increasing trend of K being noticed from all the treatments but not in T_0 . The higher increased value of K (+0.013 cmol (+)/kg) was obtained from T_4 , whereas the lower one was recorded from T_0 (-0.002 cmol (+)/kg). At the end of the study (2018-2019), exchangeable potassium ranged from -0.008 to +0.015 (cmol (+)/kg), where the upper value (+0.015 cmol (+)/kg) was demonstrated by the treatment T_4 and the lower one was observed in T_0 (-0.008 cmol (+)/kg).

 Table 12: Changes of soil exchangeable potassium (cmol (+)/kg) as influenced by different soil amendments for three consecutive years of wheat production.

Traatmant		2016-20	17		2017-2018			2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T ₀	0.150	0.088	-0.063*	0.088	0.085	-0.002 ^{ns}	0.085	0.078	-0.008 ^{ns}	
T_1	0.150	0.110	-0.040*	0.110	0.118	+0.007 ^{ns}	0.118	0.122	$+0.004^{ns}$	
T_2	0.150	0.118	-0.033*	0.118	0.123	+0.005 ^{ns}	0.123	0.130	$+0.008^{ns}$	
T ₃	0.150	0.123	-0.028*	0.123	0.133	$+0.010^{ns}$	0.133	0.145	+0.012*	
T_4	0.150	0.133	-0.018*	0.133	0.145	+0.013*	0.145	0.160	+0.015*	
T_5	0.150	0.128	-0.023*	0.128	0.135	$+0.008^{ns}$	0.135	0.143	$+0.008^{ns}$	
T_6	0.150	0.114	-0.036*	0.114	0.122	+0.008 ^{ns}	0.122	0.130	$+0.009^{ns}$	
T_7	0.150	0.125	-0.025*	0.125	0.130	+0.005 ^{ns}	0.130	0.136	$+0.006^{ns}$	
T_8	0.150	0.105	-0.045*	0.105	0.110	$+0.005^{ns}$	0.110	0.120	+0.010*	
LSD (0.05)		0.007			0.010			0.009		
CV (%)		3.300			5.000			4.200		

4.1.2.5 Available phosphorus

The statistical analysis indicated mostly significant differences among the various soil amendments on the available phosphorus in 2016-2017 (Table 13). Available phosphorus was found decreasing in the first year of the experiment where the treatment T_0 led to the maximum deviation (-13.35 µg/g) followed by T_8 (-11.18 µg/g). Compared with the final readings between 2016-2017 and 2017-2018, a little enhancement was recorded by all the treatments except in T_0 . The treatment T_3 led to the highest enhancement of available phosphorus (+5.42 µg/g), while the opposite result with lessen value (-0.70 µg/g) was observed from T_0 . The year 2018-2019, the advancement trend of phosphorus remains unchanged and the value +4.05 µg/g was led to a higher increase but -1.07 µg/g was indicated the reduction by T_3 and T_0 respectively.

Table 13: Changes of soil available phosphorus ($\mu g/g$) as influenced by different soilamendments for three consecutive years of wheat production.

Treatment		2016-20	017	2	2017-2018			2018-2019		
Treatment .	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T_0	26.30	12.96	-13.35*	12.96	12.26	-0.70 ^{ns}	12.26	11.19	-1.07 ^{ns}	
T_1	26.30	23.49	-2.82*	23.49	24.42	$+0.93^{ns}$	24.42	25.85	$+1.44^{ns}$	
T_2	26.30	25.65	-0.66 ^{ns}	25.65	29.36	+3.72*	29.36	31.54	+2.18*	
T_3	26.30	26.09	-0.22 ^{ns}	26.09	31.50	+5.42*	31.50	35.55	+4.05*	
T_4	26.30	25.22	-1.09*	25.22	27.33	+2.11*	27.33	29.34	+2.02*	
T_5	26.30	19.23	-7.07*	19.23	20.52	$+1.29^{ns}$	20.52	21.57	$+1.05^{ns}$	
T_6	26.30	21.41	-4.89*	21.41	22.96	+1.55*	22.96	24.14	+1.18 ^{ns}	
T_7	26.30	16.79	-9.51*	16.79	17.70	+0.91 ^{ns}	17.70	18.60	$+0.90^{ns}$	
T_8	26.30	15.13	-11.18*	15.13	16.53	$+1.40^{ns}$	16.53	17.50	$+0.98^{ns}$	
LSD (0.05)		1.00			1.43			1.87		
CV (%)		2.60			4.00			4.80		

4.1.2.6 Available sulphur

Available sulphur (S) content of soil mostly showed an increasing trend due to addition of different soil amendments after the completion of the first year experiment (Table 14). The large significant gain (+4.51 µg/g) of available sulphur occurred in T₃. On the other hand, a small decrease (-0.05 µg/g) on this parameter was recorded under T₈ over the initial value. A little gain was demonstrated by all the treatments except control (T₀) at the second year of the study. However, the utmost enhance (+1.32 µg/g) and decline (-0.92 µg/g) were reported by the treatments T₄ and T₀ correspondingly. The treatments had the similar tendency in the final year that was shown in the previous year on this element. But there the higher and the lower value was denoted as +1.45 and -0.97 µg/g under the treatment T₃ and T₀, respectively.

Table 14: Changes of soil available sulphur ($\mu g/g$) as influenced by different soil amendmentsfor three consecutive years of wheat production.

Treatment		2016-201	17		2017-2018			2018-2019		
	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T_0	12.50	11.56	-0.94 ^{ns}	11.56	10.65	-0.92 ^{ns}	10.65	9.68	-0.97 ^{ns}	
T_1	12.50	14.96	+2.46*	14.96	15.73	$+0.78^{ns}$	15.73	16.32	$+0.59^{ns}$	
T_2	12.50	16.17	+3.67*	16.17	17.42	$+1.25^{ns}$	17.42	18.53	$+1.11^{ns}$	
T_3	12.50	17.01	+4.51*	17.01	18.20	$+1.20^{ns}$	18.20	19.65	$+1.45^{ns}$	
T_4	12.50	15.39	+2.89*	15.39	16.71	$+1.32^{ns}$	16.71	18.02	+1.31 ^{ns}	
T ₅	12.50	13.79	+1.29*	13.79	14.83	+1.05 ^{ns}	14.83	15.78	+0.95 ns	
T_6	12.50	14.21	+1.71*	14.21	15.36	$+1.15^{ns}$	15.36	16.21	$+0.86^{ns}$	
T_7	12.50	13.39	$+0.89^{ns}$	13.39	14.42	$+1.03^{ns}$	14.42	15.35	+0.93 ^{ns}	
T_8	12.50	12.45	-0.05 ^{ns}	12.45	13.05	$+0.60^{ns}$	13.05	13.83	$+0.78^{ns}$	
LSD (0.05)		1.06			1.46			1.58		
CV (%)		4.80			6.00			6.10		

4.1.2.7 Available zinc

Data presented in the table (Table 15) revealed that the available zinc was significantly influenced by different soil amendments except T₃ and T₄ in 2016-2017. The maximum significant depletion of zinc (-0.15 μ g/g) was observed in T₀ but a minor decline was noticed in T₄ (-0.02 μ g/g). Positive balances of available zinc over initial were recorded by the amendments but not in T₀ and T₈ in 2017-2018. The treatment T₄ (+0.05 μ g/g) and T₃ (+0.05 μ g/g) were presented with a greater increase of zinc, however, only T₀ (-0.03 μ g/g) had the decrease effect. Incase of 2018-2019, treatment T₄ followed by T₃ showed the maximum significant gain in total zinc with the values of +0.06 and +0.05 μ g/g, respectively (Table 15). Furthermore, total available zinc was slightly reduced under the treatment T₀ (-0.04 μ g/g) and T₈ (-0.02 μ g/g).

Table 15: Changes of soil available zinc ($\mu g/g$) as influenced by different soil amendmentsfor three consecutive years of wheat production.

Treatment	2016-2017				2017-2018			2018-2019		
	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T_0	0.75	0.60	-0.15*	0.60	0.56	-0.03 ^{ns}	0.56	0.54	-0.04 ^{ns}	
T_1	0.75	0.63	-0.13*	0.63	0.65	$+0.03^{ns}$	0.65	0.69	$+0.04^{ns}$	
T_2	0.75	0.70	-0.05 ^{ns}	0.70	0.74	$+0.04^{ns}$	0.74	0.78	$+0.04^{ns}$	
T_3	0.75	0.73	-0.03 ^{ns}	0.73	0.77	$+0.05^{ns}$	0.77	0.82	$+0.05^{ns}$	
T_4	0.75	0.74	-0.02*	0.74	0.79	$+0.05^{ns}$	0.79	0.84	$+0.06^{ns}$	
T_5	0.75	0.68	-0.07*	0.68	0.72	$+0.04^{ns}$	0.72	0.75	$+0.03^{ns}$	
T_6	0.75	0.64	-0.12*	0.64	0.66	$+0.02^{ns}$	0.66	0.68	$+0.02^{ns}$	
T_7	0.75	0.66	-0.10*	0.66	0.67	$+0.02^{ns}$	0.67	0.70	$+0.03^{ns}$	
T_8	0.75	0.62	-0.13*	0.62	0.61	-0.01 ^{ns}	0.61	0.59	-0.02 ^{ns}	
LSD (0.05)		0.05			0.08			0.12		
CV (%)		4.00			7.20			10.00		

4.1.2.8 Electrical conductivity

The statistical analysis indicated differences among the various soil amendments on electrical conductivity in 2016-2017 (Table 16). Electrical conductivity (EC) was decreased under the control treatment (T₀) and all other amendments were increased the EC. Here T₄ had the maximum (+10.50 μ s/cm) EC than the other treatments. In the year 2017-2018, positive EC was recorded by most of the treatments, but not in T₀. Here the treatment T₄ led to the highest enhancement of EC (+12.50 μ s/cm), while the decreased EC (-6.50 μ s/cm) was observed in T₀. Except T₀, there was also found positive deviation of EC over initial by addition of other treatments in 2018-2019. Nevertheless, the advancement of EC +13.00 μ s/cm led to the higher increase which was shown by T₄ but -7.00 μ s/cm was indicate by T₀, respectively.

Table 16: Changes of soil electrical conductivity (μ s/cm) as influenced by different soilamendments for three consecutive years of wheat production.

Treatment		2016-2	017		2017-2018			2018-2019		
	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T_0	145.00	140.00	-5.00*	140.00	133.50	-6.50*	133.50	126.50	-7.00*	
T_1	145.00	147.00	$+2.00^{ns}$	147.00	150.00	+3.00 ^{ns}	150.00	153.50	+3.50 ^{ns}	
T_2	145.00	150.50	+5.50*	150.50	160.00	+9.50*	160.00	167.50	+7.50*	
T_3	145.00	152.50	+7.50*	152.50	164.00	+11.50*	164.00	175.00	+11.00*	
T_4	145.00	155.50	+10.50*	155.50	168.00	+12.50*	168.00	181.00	+13.00*	
T_5	145.00	147.00	$+2.00^{ns}$	147.00	149.50	$+2.50^{ns}$	149.50	152.50	$+3.00^{ns}$	
T_6	145.00	149.00	+4.00*	149.00	154.00	$+5.00^{ns}$	154.00	160.50	+6.50*	
T_7	145.00	146.50	$+1.50^{ns}$	146.50	147.50	$+1.00^{ns}$	147.50	149.50	$+2.00^{ns}$	
T_8	145.00	147.50	$+2.50^{ns}$	147.50	151.50	$+4.00^{ns}$	151.50	156.50	$+5.00^{ns}$	
LSD (0.05)		3.58			5.58			5.85		
CV (%)		1.50			2.20			2.30		

4.1.2.9 Carbon nitrogen ratio

There was a significant influence of various soil amendments on carbon nitrogen ratio (C:N) in the year 2016-2017 (Table 17). Here the C:N was found to be greater (+4.15) under the treatment T_0 , while the plots under treatment T_5 provided the lower C:N (+0.12) of soil. As the consecutive year 2017-2018, C:N was declined insignificantly by the application of different soil amendments without T_0 and T_8 . The treatment T_4 was demonstrated the higher decrease of C:N (-0.69), whereas the increase of C:N was found from T_0 (+4.42) and T_8 (+1.31). At the end year of the study, the result revealed that the C:N of soil was also decreased for all the applied treatments except T_0 and T_8 . The most decrease of C:N was recorded in T_4 (-0.60) but a major positive value was found in T_8 (+4.88).

 Table 17: Changes of soil carbon nitrogen ratio (C:N) as influenced by different soil amendments for three consecutive years of wheat production.

Treatment		2016-20)17	,	2017-2018			2018-2019		
Treatment	Initial	Final	Change	Initial	Final	Change	Initial	Final	Change	
T ₀	10.00	14.15	+4.15*	14.15	18.64	+4.42*	18.64	22.91	+4.22*	
T_1	10.00	13.80	+3.80*	13.80	13.52	-0.28 ^{ns}	13.52	13.70	-0.14 ^{ns}	
T_2	10.00	12.64	+2.64*	12.64	12.19	-0.44 ^{ns}	12.19	11.87	-0.32 ^{ns}	
T_3	10.00	11.51	+1.51*	11.51	10.88	-0.63 ^{ns}	10.88	10.52	-0.35 ^{ns}	
T_4	10.00	10.47	+0.47*	10.47	8.75	-0.69 ^{ns}	8.75	8.16	-0.60 ^{ns}	
T_5	10.00	10.12	+0.12*	10.12	9.44	-0.68 ^{ns}	9.44	8.89	-0.55 ^{ns}	
T_6	10.00	12.72	+2.72*	12.72	12.57	-0.15 ^{ns}	12.57	12.48	-0.45 ^{ns}	
T_7	10.00	10.58	+0.58*	10.58	10.05	-0.52 ^{ns}	10.05	9.64	-0.41 ^{ns}	
T_8	10.00	12.49	+2.49*	12.49	13.80	$+1.31^{ns}$	13.80	18.69	+4.88*	
LSD (0.05)		0.91			2.22			2.20		
CV (%)		5.00			11.20			10.70		

4.2 Wheat

4.2.1 Seedling infection

After 14 days of sowing the seedling infection was recorded in the three consecutive cropping years (Table 18). The treatment used for the study showed significant results in reducing seedling infection in the field during the study periods 2016-2017, 2017-2018 and 2018-2019. Among the selected three varieties of wheat, the variety BARI wheat-28 (V_1) showed the higher infection (1.24%), while the variety BARI wheat- 30 (V_3) exhibited minimum seedling infection (0.92%).

Regarding treatment effect, the lower rate of seedling infection (0.32%) was found after application *Trichoderma harzianum* (T₅) in comparison with the maximum infection from T₀ (2.32%) in 2018-2019. All the treatments in the experimental period presented significant differences among each other. But T₁ (Rice straw + vermicompost + green manure) and T₈ (Chemical fertilizer) was statistically identical. Reduction of seedling infection at 14 DAS under different soil amendments showed more or less the following order as T₅>T₇>T₄>T₃>T₂>T₆>T₁>T₈>T₀ during the whole research period (Table 19).

Though all the amendments in combination with variety showed significant performance in reducing seedling infection at 14 DAS, but T_5 with each variety was presented the minimum infection in the three consecutive years. The lowest (0.25%) and the highest (2.69%) infection of seedling were found in 2018-2019 from V_3T_5 and V_1T_0 , respectively. Along with significant difference, parallel tendency were noticed regarding lower infection rate by V_2T_5 (0.33%) and V_1T_5 (0.38%) but higher infection values were recorded from V_2T_0 (2.40%) and V_3T_0 (1.88%) in same year of the study. Similar results were also noticed from the aforesaid combination during 2016-2017 and 2017-2018 (Table 20).

4.2.2 Seedling blight

The frequency of seedling blight at 21 DAS was recorded and found to be significantly different for all varieties. In the experiment, the higher seedling blight was found in V₁ and the lower was in V₃ throughout the study time (Table 18). Hence, the highest percentage survivability of seedling was obtained from V₃ (1.73%) in 2017-2018 and the lowest from V₁ (2.41%) in 2018-2019.

Under this study, recorded results showed that there was a significant variation among the treatments. The highest inhibition effect against seedling blight (0.87%) was observed by treatment T_5 and the lowest (4.74%) from control (T_0) during 2018-2019. Next to treatment T_5 , the treatment T_7 also showed remarkable effect in reducing seedling blight (1.07%) which was statistically identical (1.09%) with T_4 (Table 19).

The interaction effect of treatments and variety clearly demonstrated significant influence on seedling blight at 21 DAS. All the varieties presented the minimum blight incidence in combination with T_5 , while the maximum blight was found from combination with T_0 in the whole time of study (Table 20). The results indicated that among the treatment, V_3T_5 combination gave the comparatively lower seedling blight incidence (0.73%) in 2018-2019. On the other hand, V_1T_0 showed the higher seedling blight incidence (5.44%) in 2018-2019.

4.2.3 Leaf chlorophyll content

Leaf chlorophyll content of wheat was affected significantly due to varietal influences in the three years of this experiment. From the responses of individual variety, provided the following order $V_3 > V_2 > V_1$ in respect of chlorophyll content over the whole study period (Table 18). Among three varieties, V_3 produced the sharp increase of chlorophyll content and it was recorded maximum (45.87 SPAD) during 2018-2019 which exhibited a statistically similar result (45.42 SPAD) with V_1 at the same year.

A significant difference was observed from various soil amendments for the chlorophyll content of wheat leaf at 70 DAS. The biggest value of chlorophyll content (51.48 SPAD) was recorded from chemical fertilizer (T_8) and which was followed by T_4 (48.37 SPAD) and T_5 (48.08). In this regard, the smallest value (40.33 SPAD) was noticed from T_0 during the season 2018-2019. Due to soil amendment with chemical fertilizer (T_8) provided more chlorophyll content and the trend was maintained from beginning to the end of the study (Table 19).

Diverse soil amendments in combination with selected varieties attributed significant effect in leaf chlorophyll content of wheat. Among all combinations of treatments and variety, V_3T_8 presented higher value (52.52 SPAD), but V_2T_0 had the poorest value (38.40 SPAD) of chlorophyll content during 2018-2019. Indication of chlorophyll content as like above were noticed during the year 2016-2017 and 2017-2018, respectively (Table 20).

	See	dling infec (%)	etion	Se	edling blig (%)	ght	Leaf ch	nlorophyll (SPAD)	content
Variety	14 Da	ays after so	owing	21 Da	iys after so	owing	70 Da	ays after so	owing
	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019
V_1	1.26	1.22	1.28	2.24	2.09	2.41	42.29	43.61	45.42
v ₁	± 0.13a	± 0.12a	± 0.14a	± 0.17a	± 0.17a	± 0.26a	$\pm 0.65b$	$\pm 0.69b$	± 0.73a
V_2	1.15	1.12	1.10	2.04	1.92	1.98	41.31	42.87	44.13
v ₂	± 0.11b	± 0.11b	$\pm 0.12b$	$\pm 0.15b$	$\pm 0.15b$	± 0.21b	± 0.59c	$\pm 0.75c$	$\pm 0.81b$
X7	1.02	0.94	0.97	1.83	1.73	1.84	43.03	44.65	45.87
V ₃	± 0.09c	± 0.09c	± 0.10c	± 0.13c	± 0.13c	± 0.20c	± 0.75a	± 0.79a	± 0.73a
LS	**	**	**	**	**	**	**	**	*
CV (%)	5.42	6.89	4.38	5.59	6.16	4.30	0.80	0.79	5.09

 Table 18: Effect of variety on seedling infection, seedling blight and leaf chlorophyll content of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, DAS = Days after sowing, SPAD = The soil plant analysis development, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

	Seedlin	ng infection	n (%)	See	dling blight	(%)	Leaf chlor	ophyll conte	ent (SPAD)
	14 Da	ys after sov	wing	21 D	ays after so	wing	70 E	Days after so	wing
Treatment	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019
T ₀	2.14	1.99	2.32	3.52	3.30	4.74	38.26	38.82	40.33
10	± 0.12a	$\pm 0.09a$	$\pm 0.13a$	± 0.16a	± 0.16a	$\pm 0.20a$	$\pm 0.85 h$	$\pm 0.90g$	$\pm 0.79 d$
T_1	1.49	1.47	1.46	2.53	2.41	2.48	40.18	41.62	42.77
11	$\pm 0.03b$	$\pm 0.05 b$	$\pm 0.09b$	$\pm 0.08b$	$\pm 0.08 b$	$\pm 0.19c$	$\pm 0.90 g$	$\pm 0.86 f$	$\pm 0.87c$
T_2	1.34	1.24	1.20	2.17	2.04	2.11	41.39	43.56	42.96
12	$\pm 0.04 d$	$\pm 0.05c$	$\pm 0.04c$	$\pm 0.06c$	$\pm 0.04 d$	$\pm 0.08 d$	$\pm 0.87 f$	± 1.05 de	$\pm 0.76c$
т	0.93	0.87	0.92	1.77	1.71	1.61	42.45	43.85	44.09
T ₃	$\pm 0.04e$	$\pm 0.03 d$	$\pm 0.05 d$	$\pm 0.05 d$	$\pm 0.05 e$	$\pm 0.04e$	$\pm 0.85 d$	± 1.04 d	$\pm 0.85c$
т	0.59	0.58	0.55	1.24	1.19	1.09	44.59	46.63	48.37
T_4	$\pm 0.02 f$	$\pm 0.02e$	$\pm 0.01e$	$\pm 0.05e$	$\pm 0.06 f$	$\pm \ 0.05 f$	$\pm 0.81 b$	$\pm 0.89 b$	$\pm 0.80 b$
т	0.35	0.34	0.32	0.93	0.87	0.87	43.03	44.73	48.08
T ₅	$\pm 0.02g$	$\pm 0.02g$	$\pm 0.02g$	$\pm 0.06 f$	$\pm 0.04g$	$\pm 0.05 g$	$\pm 0.85c$	$\pm 0.91c$	$\pm 0.61 b$
т	1.43	1.44	1.42	2.44	2.33	2.16	41.08	41.80	43.06
T ₆	$\pm 0.02c$	$\pm 0.06b$	$\pm 0.08b$	$\pm 0.07b$	$\pm 0.08 bc$	$\pm 0.11 d$	$\pm 0.89 f$	$\pm 0.92 f$	$\pm 0.63c$
т	0.57	0.46	0.45	1.29	1.11	1.07	41.78	43.45	45.11
T_7	$\pm 0.02 f$	$\pm 0.02 f$	$\pm 0.03 f$	$\pm 0.04e$	$\pm 0.03 f$	$\pm 0.06 f$	$\pm 0.92e$	$\pm 0.95e$	$\pm 0.71c$
т	1.48	1.47	1.42	2.44	2.26	2.57	47.14	48.92	51.48
T ₈	$\pm 0.10c$	$\pm 0.08 b$	$\pm 0.04b$	$\pm 0.09b$	$\pm 0.04c$	$\pm 0.11b$	$\pm 0.97a$	± 1.08a	± 0.70a
LS	**	**	**	**	**	**	**	**	**
CV (%)	5.42	6.89	4.38	5.59	6.16	4.30	0.80	0.79	5.09

Table 19: Effect of organic and probiotic soil amendments on seedling infection, seedling blight and leaf chlorophyll content of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

Table 20: Interaction effect of variety and soil amendments on seedling infection, seedling blight and leaf chlorophyll content of wheat.

Vor	riotry Br	Seed	ling infectio	n (%)	See	dling blight	(%)	Leaf chlore	ophyll conte	nt (SPAD)
	riety &	14 D	Days after so	wing	21 D	ays after so	wing	70 D	ays after sov	wing
Tre	atment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	т	2.52	2.26	2.69	4.01	3.81	5.44	37.79	39.22	40.90
	T ₀	$\pm 0.08a$	$\pm 0.07a$	$\pm 0.090a$	$\pm 0.18a$	± 0.16a	± 0.18a	± 1.861	± 1.36n	± 1.39
	т	1.55	1.58	1.78	2.74	2.60	3.21	40.85	41.75	43.05
	T_1	$\pm 0.04e$	± 0.07 de	$\pm 0.05 d$	$\pm 0.14d$	± 0.18 cd	$\pm 0.15c$	± 1.77 hi	$\pm 1.47k$	± 1.74
	т	1.48	1.37	1.28	2.32	2.05	2.39	41.82	44.85	43.02
	T ₂	$\pm 0.04 ef$	$\pm 0.04 f$	± 0.06 hi	$\pm 0.10 gh$	$\pm 0.08 hi$	$\pm 0.07 f$	± 1.69g	$\pm 1.94 f$	± 1.38
	т	0.98	0.95	1.10	1.86	1.79	1.62	42.22	42.57	45.14
	T ₃	$\pm 0.02 k$	$\pm 0.03i$	$\pm 0.05 j$	$\pm 0.07 \mathrm{jk}$	± 0.10 jk	$\pm 0.08 j$	± 1.74 fg	± 1.76ij	± 1.74
V	т	0.61	0.63	0.55	1.38	1.39	1.25	44.68	46.71	47.33
V_1	T_4	$\pm 0.03 m$	$\pm 0.02 kl$	± 0.031	$\pm 0.06 \mathrm{m}$	± 0.061	$\pm 0.06k$	± 1.43d	$\pm 1.45d$	± 1.52
	т	0.40	0.39	0.38	1.15	1.01	1.03	43.31	43.12	48.90
	T ₅	$\pm 0.02 op$	$\pm 0.02 mn$	$\pm 0.02 m$	$\pm 0.03 no$	$\pm \ 0.04mn$	± 0.051	$\pm 1.42e$	± 1.54 hi	± 1.04
	т	1.38	1.55	1.72	2.57	2.60	2.57	41.12	42.76	43.70
	T ₆	$\pm \ 0.05 fgh$	$\pm 0.07e$	$\pm 0.07 d$	± 0.06 de	$\pm 0.08 cd$	$\pm 0.05e$	$\pm 1.54h$	$\pm 1.89 hi$	± 1.14
	т	0.62	0.523	0.53	1.36	1.14	1.26	42.09	42.96	45.37
	T ₇	$\pm 0.02 m$	$\pm 0.02 kl$	± 0.021	$\pm 0.07 m$	$\pm 0.02 m$	$\pm 0.05 k$	± 1.61 fg	± 1.86 hi	± 1.40
	T ₈	1.84	1.72	1.54	2.74	2.38	2.92	46.73	48.53	51.38
	18	$\pm 0.08c$	$\pm 0.07c$	$\pm 0.02e$	$\pm 0.09 d$	$\pm 0.05 \text{ef}$	$\pm 0.08 d$	$\pm 1.49b$	± 1.94b	± 1.45
	T ₀	2.16	2.00	2.40	3.54	3.31	4.43	38.11	37.22	38.40
	10	$\pm 0.09b$	$\pm 0.10b$	$\pm 0.06b$	$\pm 0.10b$	$\pm 0.09 b$	$\pm 0.12b$	± 1.271	± 1.500	± 0.92
	т	1.52	1.51	1.40	2.53	2.42	2.16	39.32	41.00	42.08
	T_1	$\pm 0.04e$	$\pm 0.05 e$	$\pm 0.04 fg$	$\pm 0.06 \text{ef}$	$\pm 0.12 de$	$\pm 0.10g$	$\pm 1.82k$	± 1.551	± 1.70
	т	1.30	1.26	1.22	2.14	2.14	1.98	40.55	42.03	41.55
	T ₂	± 0.05 hij	$\pm 0.06 \mathrm{fg}$	$\pm 0.03 i$	$\pm 0.02 \text{hi}$	$\pm 0.02 ghi$	$\pm 0.05 \text{hi}$	\pm 1.52hij	± 1.94jk	± 1.38
	T ₃	1.02	0.90	0.85	1.82	1.75	1.68	41.78	43.21	42.27
	13	$\pm 0.05 k$	$\pm 0.04i$	$\pm 0.03k$	$\pm 0.08 kl$	± 0.08 jk	± 0.06j	$\pm 1.59g$	$\pm 1.92h$	± 0.92
V_2	T_4	0.61	0.58	0.56	1.26	1.16	1.04	43.35	45.67	48.57
2	14	$\pm 0.01 \text{mn}$	$\pm 0.02 kl$	± 0.021	$\pm 0.06 mno$	$\pm 0.02 m$	± 0.051	± 1.22e	± 1.71e	± 1.62
	T ₅	0.34	0.34	0.33	0.84	0.84	0.83	42.50	44.87	47.62
	15	$\pm 0.01 pq$	$\pm 0.02 mn$	$\pm 0.01 mn$	$\pm 0.03 p$	$\pm 0.02 no$	$\pm 0.03 mn$	$\pm 1.84 f$	$\pm 1.63 f$	± 1.36
	T_6	1.46	1.54	1.20	2.51	2.29	2.05	40.07	41.00	41.97
	•0	$\pm 0.04 efg$	$\pm 0.04e$	$\pm 0.05 \mathrm{i}$	$\pm 0.10 efg$	$\pm 0.08 efg$	$\pm 0.08 gh$	± 1.51j	± 1.671	± 1.08
	T_7	0.60	0.46	0.47	1.33	1.15	1.03	40.95	43.07	44.22
	• /	$\pm 0.02 mn$	$\pm 0.01 lm$	± 0.021	$\pm 0.06 mn$	$\pm 0.04 m$	± 0.041	\pm 1.89hi	\pm 1.87hi	± 1.20
	T_8	1.35	1.56	1.45	2.35	2.25	2.61	45.17	47.79	50.52
	- 0	± 0.06 ghi	± 0.06e	$\pm 0.04 f$	$\pm 0.07 fgh$	± 0.05 e-h	± 0.07e	± 1.45d	± 1.91c	± 0.93

	Contd.
eaf chlorophyll con	tent (SPA

Vori	ety &	Seed	ling infectio	n (%)	See	dling blight	(%)	Leaf chlor	ophyll conte	nt (SPAD)
	tment	14 D	ays after so	wing	21 D	ays after so	wing	70 D	ays after so	wing
IIca		2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
		1.74	1.70	1.88	3.00	2.79	4.36	38.89	40.02	41.70
	T ₀	± 0.04d	± 0.04 cd	± 0.09c	$\pm 0.07c$	$\pm 0.12c$	$\pm 0.21 b$	± 1.80k	± 1.85m	± 1.35
		1.41	1.31	1.21	2.33	2.19	2.07	40.36	42.11	43.17
	T_1	$\pm 0.03 fgh$	$\pm 0.02 \mathrm{fg}$	$\pm 0.05i$	$\pm 0.10 gh$	± 0.06 fgh	$\pm 0.05 gh$	± 1.64ij	± 1.98jk	± 1.69
	T	1.24	1.09	1.11	2.03	1.93	1.95	41.80	43.79	44.32
	T ₂	$\pm 0.05 \mathrm{j}$	$\pm 0.04 h$	$\pm 0.05 j$	$\pm 0.07 ij$	$\pm 0.05 ij$	$\pm 0.09 hi$	$\pm 1.81g$	± 1.90g	± 1.07
	T	0.79	0.76	0.82	1.64	1.59	1.54	43.36	45.76	44.88
	T ₃	$\pm 0.02l$	$\pm 0.01 j$	$\pm 0.02 k$	± 0.091	$\pm 0.05 k$	$\pm 0.07 j$	± 1.55e	± 1.80e	± 1.47
V_3	т	0.54	0.54	0.53	1.08	1.02	0.97	45.73	47.51	49.22
5	T_4	$\pm 0.03 \text{mn}$	± 0.01 kl	± 0.021	$\pm 0.04o$	$\pm 0.04 mn$	$\pm 0.04 lm$	± 1.72c	± 1.92c	± 1.36
	T	0.30	0.28	0.25	0.78	0.76	0.73	43.28	46.21	47.73
	T ₅	$\pm 0.02q$	$\pm 0.01 n$	$\pm 0.02 n$	$\pm 0.02 p$	$\pm 0.02o$	$\pm 0.03 n$	± 1.75e	± 1.56de	± 1.02
	т	1.45	1.22	1.36	2.23	2.10	1.86	42.06	41.65	43.52
	T ₆	$\pm 0.04 efg$	$\pm 0.04 gh$	$\pm 0.06 gh$	$\pm 0.08 hi$	± 0.03 ghi	$\pm 0.06i$	$\pm 1.94 \mathrm{fg}$	± 1.71k	± 1.21
	т	0.50	0.38	0.35	1.19	1.03	0.92	42.30	44.31	45.75
	T ₇	$\pm 0.01 \text{no}$	$\pm 0.01 mn$	$\pm 0.02 m$	± 0.06 mno	$\pm 0.05 mn$	$\pm 0.05 \text{lm}$	$\pm 1.87 \mathrm{fg}$	$\pm 1.79 \mathrm{fg}$	± 1.40
	т	1.25	1.20	1.27	2.23	2.16	2.18	49.51	50.45	52.52
	T_8	$\pm 0.05 ij$	$\pm 0.03 gh$	$\pm 0.04i$	$\pm 0.08 hi$	$\pm 0.04 gh$	$\pm 0.05g$	± 1.56a	± 2.19a	± 1.38
	LS	**	**	**	**	**	**	**	**	NS
	CV (%)	5.42	6.89	4.38	5.59	6.16	4.30	0.80	0.79	5.09

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

4.3 Growth attributes

4.3.1 Total dry matter

Generally, with the advancement of time the total dry matter (TDM) accumulation of wheat plant was positively increased in all varieties. TDM varied significantly due to various varietal influences at 25, 40, 55, 70 and 85 DAS during the experimental period (Fig. 3, 4 and 5). Results indicated that TDM was higher in V₃ followed by V₁, but it was lower in V₂ at all dates of sampling. At 85 DAS, the maximum TDM accumulation (722.34 g/m²) was recorded from V₃ in 2018-2019 and the minimum from V₂ (625.74 g/m²) in 2016-2017.

From the evaluation of different soil amendments, it has been found significant variation among the treatments at almost all DAS. Most of the treatments showed approximately similar patterns of TDM accumulation. However, the highest TDM was produced in T_8 followed in descending order by T_4 , T_5 , T_7 , T_3 , T_2 , T_6 and T_1 , while the lowest was found in T_0 at all sampling dates round the three years study (Fig. 6, 7 and 8). At 25 DAS, the treatment T_8 presented the uppermost value (34.84 g/m²) for TDM accumulation, while T_0 gave the lowest (17.03 g/m²) one in the 2018-2019 cropping season. Thus, considering the TDM production at 40, 55, 70 and 85 DAS the results revealed that the maximum and the minimum accumulation exhibited by T_8 (118.01, 242.53, 561.61, 798.15 g/m²) and T_0 (45.92, 124.08, 329.92, 532.33 g/m²), respectively from the sequential year of the study. At the peak (85 DAS) of TDM accumulation, the most statistical identity was found between T_8 (767.14 g/m²) and T_4 (750.91 g/m²) and similar trend was also found between T_7 (712.32 g/m²) and T_3 (706.39 g/m²) in 2018-2019.

In 2016-2017, TDM accumulation significantly influenced at 25, 40 and 55 DAS by the interaction of variety and different organic soil amendment treatments. Here the great and less values of TDM were recorded on the aforesaid different DAS as V_3T_8 (30.34, 96.97, 222.63 g/m²) and V_2T_0 (15.91, 43.40, 117.18 g/m²). However, among the combination, V_2T_8 (207.95 g/m²) reported higher which was statistically at par with V_3T_4 (205.31 g/m²) at 55 DAS.

Variation of TDM in respect of combination of variety and different organic soil amendments were significantly differed only at 25 and 55 DAS in the year 2017-2018. But at 25 DAS, V_3T_8 provided the maximum value (31.11 g/m²) and V_2T_0 showed the minimum value (17.11 g/m²) of TDM content in the plant. In case of 55 DAS, the highest TDM (230.93 g/m²) was obtained from the treatment combination of V_3T_8 and the lowest (127.47 g/m²) was observed in V_2T_0 . Due to treatment application significant differences of TDM content was notified at 25, 40 and 55 DAS in 2018-2019 and similar trend was observed in 2016-2017. From the above mentioned sampling dates, it was noticed that the production of TDM ranked as followed by the combination V_3T_8 (252.05 g/m²) > V_1T_8 (242.89 g/m²) > V_2T_8 (232.65 g/m²) and V_3T_0 (157.92 g/m²) > V_1T_0 (145.62 g/m²) > V_2T_0 (139.22 g/m²) in 2018-2019, where statistical identity was noticed between V_3T_8 and V_1T_8 (Table 21).




Fig. 3: Effect of variety on total dry matter content of wheat in 2016-2017.




Fig. 4: Effect of variety on total dry matter content of wheat in 2017-2018.




Fig. 5: Effect of variety on total dry matter content of wheat in 2018-2019.




Fig. 6: Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2016-2017.




Fig. 7: Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2017-2018.




Fig. 8: Effect of organic and probiotic soil amendments on total dry matter content of wheat in 2018-2019.

Variet	v &			2016-2017					2017-2018					2018-2019		
Treatn	•							Days	after sowing	(DAS)						
ITeath	lont	25	40	55	70	85	25	40	55	70	85	25	40	55	70	85
	T_0	18.95	45.05	123.57	325.24	528.50	20.06	50.92	137.44	368.48	556.70	16.42	60.34	145.62	400.52	576.84
	10	$\pm 0.82n$	$\pm 1.95 n$	$\pm 5.35r$	± 14.08	± 22.89	$\pm \ 0.87m$	± 2.21	$\pm 5.95 o$	± 15.96	± 24.11	$\pm 0.71 m$	$\pm2.613o$	$\pm 6.31 p$	± 17.34	± 24.9
	T_1	20.671	61.86	156.99	386.93	588.54	21.80	67.93	167.09	397.28	600.46	22.32	75.42	178.56	431.28	618.6
	11	$\pm 1.01 m$	$\pm 3.04 k$	\pm 7.70no	± 18.99	± 28.88	± 1.071	± 3.33	\pm 8.20mn	± 19.50	± 29.47	$\pm 1.10k$	$\pm 3.70 m$	$\pm 8.76n$	± 19.43	± 30.3
	T_2	21.56	76.75	175.13	411.80	632.97	23.50	80.65	190.15	435.25	652.51	24.28	85.67	200.41	457.43	679.2
	12	± 0.67 ijk	$\pm 3.64 fg$	$\pm 8.38 m$	± 18.00	± 27.31	$\pm 0.78ij$	± 4.00	\pm 8.96kl	± 15.89	± 31.90	$\pm 0.83 ij$	$\pm 4.234j$	\pm 8.68jkl	± 20.64	± 29.3
	T ₃	22.72	78.72	188.57	428.25	647.41	25.07	84.11	201.83	454.77	674.34	26.83	96.57	213.86	484.89	702.6
	13	$\pm 0.92 gh$	$\pm 3.18 efg$	\pm 7.62f-j	± 17.31	± 26.17	$\pm 1.01 gh$	± 3.40	\pm 8.16ghi	± 18.38	± 27.25	$\pm 1.08 \mathrm{fg}$	$\pm 3.903 h$	\pm 8.64e-i	± 19.60	± 28.4
v	T_4	25.76	86.99	198.72	469.71	725.94	28.71	92.83	219.34	503.23	751.01	30.75	108.24	232.68	530.34	780.0
V_1	14	$\pm 1.19d$	$\pm 4.02c$	$\pm 9.18d$	± 21.70	± 33.53	$\pm 1.33 cd$	± 4.29	$\pm 10.13 bcd$	± 23.24	\pm 34.69	$\pm 1.42 bc$	$\pm 4.999 de$	$\pm 10.75 cd$	± 24.50	± 36.0
	T 5	23.36	81.42	191.18	447.14	691.15	25.87	87.49	202.08	463.43	689.32	27.32	101.57	214.43	492.03	730.2
	15	$\pm \ 0.67 defg$	$\pm 2.35 de$	$\pm 5.52 efgh$	± 12.91	± 19.95	$\pm 0.75g$	± 2.53	± 5.83 ghi	± 13.38	± 19.90	$\pm 0.79 \mathrm{fg}$	$\pm 2.932 f$	$\pm 6.19 efgh$	± 14.20	±21.0
	т	21.25	63.92	162.86	398.99	610.13	22.53	71.63	173.71	413.07	632.02	23.94	80.64	193.32	451.62	668.1
	T_6	± 0.80 jkl	$\pm 2.40 jk$	± 6.11n	± 14.97	± 22.90	± 0.85 jkl	± 2.69	$\pm 6.52m$	± 15.50	± 23.72	$\pm 0.90 ij$	± 3.0261	\pm 7.26lm	± 16.95	± 25.0
	T ₇	23.04	82.98	182.16	435.42	663.30	25.27	85.12	195.32	449.35	678.68	26.73	93.85	205.72	473.42	709.8
	17	$\pm 0.80 fg$	$\pm 2.88d$	$\pm 6.31 jk$	± 15.08	± 22.98	$\pm 0.88 gh$	± 2.95	\pm 6.77ijk	± 15.57	± 23.51	$\pm 0.93g$	$\pm 3.251 h$	\pm 7.13hijk	± 16.40	± 24.5
	т	27.41	92.25	216.35	499.29	745.75	29.35	95.55	227.34	536.63	763.57	36.77	117.81	242.89	570.00	797.2
	T_8	$\pm 1.35c$	$\pm 4.53b$	$\pm 10.62b$	± 24.50	± 36.60	$\pm 1.44 bc$	± 4.69	$\pm 11.16ab$	± 26.34	± 37.47	$\pm 1.81a$	$\pm 5.782b$	$\pm 11.92ab$	± 27.97	± 39.1
	T ₀	15.91	43.40	117.18	317.52	516.75	17.11	48.96	127.47	348.74	526.51	15.86	52.70	139.22	372.22	549.4
	1 ₀	$\pm 0.32o$	$\pm 0.88 n$	$\pm 2.37s$	± 6.42	± 10.44	$\pm 0.35n$	± 0.99	$\pm 2.58 p$	± 7.05	± 10.64	$\pm 0.32 m$	± 1.065p	$\pm 2.81 p$	± 7.52	±11.1
	T_1	19.96	55.87	146.08	368.60	563.78	22.42	65.18	158.46	387.37	565.83	23.10	70.13	171.44	413.99	595.8
	1	$\pm0.75m$	± 2.101	± 5.48p	± 13.83	± 21.16	± 0.84 jkl	± 2.45	$\pm 5.95 n$	± 14.54	± 21.23	$\pm 0.87 jk$	$\pm 2.632n$	$\pm 6.43n$	± 15.54	± 22.3
	T ₂	20.84	67.61	170.31	402.04	619.33	22.96	77.53	189.65	427.57	640.11	24.56	83.30	197.76	438.59	665.1
	12	$\pm 0.51 kl$	$\pm 2.94 \text{hi}$	$\pm 7.12 m$	± 18.58	± 26.41	$\pm 0.62 jkl$	± 3.45	\pm 8.12kl	± 18.75	± 27.49	± 0.70 hij	± 3.751 jkl	\pm 8.54klm	± 18.75	± 28.7
	T ₃	21.68	77.09	183.94	425.42	642.39	24.39	83.32	195.29	442.48	660.82	26.03	93.37	206.39	468.65	692.8
	13	$\pm 0.56 ij$	$\pm 2.00 fg$	$\pm 4.78 ijk$	± 11.05	± 16.69	$\pm 0.63 \text{hi}$	± 2.17	$\pm 5.08 ijk$	± 11.50	± 17.17	$\pm 0.68 gh$	$\pm 2.426 \text{hi}$	\pm 5.36g-k	± 12.18	± 18.0
\mathbf{V}_2	T_4	23.77	83.13	193.13	450.79	697.75	27.84	90.52	211.57	485.25	713.10	29.20	102.70	223.05	509.60	740.1
	14	$\pm 1.03 \text{ef}$	$\pm 3.60 d$	$\pm 8.36 defg$	± 19.52	± 30.21	$\pm 1.21 de$	± 3.92	$\pm 9.16 def$	± 21.01	± 30.88	± 1.27 de	$\pm 4.447 f$	\pm 9.66de	± 22.07	± 32.0
	T_5	23.07	80.18	189.15	428.28	652.61	25.99	87.70	205.11	469.76	692.07	26.97	97.09	216.13	485.46	709.0
	15	$\pm 0.47 fg$	$\pm 1.62 de$	± 3.82 fghi	\pm 8.65	± 13.19	$\pm 0.53g$	± 1.77	$\pm 4.15 fgh$	± 9.49	± 13.99	$\pm 0.55 fg$	± 1.96 gh	$\pm 4.37 efg$	± 9.81	± 14.3
	T 6	20.64	61.58	152.56	383.73	588.58	22.23	68.06	168.61	396.26	606.71	24.11	81.23	188.96	425.45	636.1
	16	$\pm 0.78 \text{lm}$	$\pm 2.31k$	± 5.730	± 14.40	± 22.09	± 0.83 kl	± 2.55	± 6.33m	± 14.87	± 22.77	± 0.91ij	± 3.05 kl	$\pm 7.09 \mathrm{m}$	± 15.97	± 23.8

Table 21: Interaction effects variety and of soil amendments on total dry matter content of wheat (g/m^2) .

Contd...

Varie	N 87			2016-2017					2017-2018					2018-2019		
Treatn								Days	after sowing ((DAS)						
mean	lent	25	40	55	70	85	25	40	55	70	85	25	40	55	70	85
	T ₇	22.58	78.58	185.33	431.03	643.13	25.62	84.52	192.24	446.37	669.46	26.55	94.62	209.25	470.87	687.
	17	$\pm \ 0.98 gh$	$\pm 3.40 efg$	\pm 8.03hijk	$\pm \ 18.66$	± 27.85	$\pm 1.11g$	± 3.66	\pm 8.32jkl	± 19.33	± 28.99	$\pm 1.15g$	$\pm 4.10 h$	\pm 9.06ghij	± 20.39	± 29.
	T_8	25.85	87.98	207.95	483.23	707.36	27.60	93.22	217.32	507.40	730.12	29.99	111.13	232.65	532.14	754.
	18	$\pm 0.77 d$	$\pm 3.99c$	$\pm 9.65c$	± 21.07	± 30.98	$\pm 0.86 de$	± 4.27	± 9.56 cde	± 20.59	\pm 32.17	$\pm 0.98 cd$	$\pm 4.62 cd$	\pm 12.09cd	± 21.88	± 33
	T ₀	19.18	49.31	131.50	347.01	551.74	20.01	52.21	145.22	378.70	572.48	18.82	63.50	157.92	409.58	597.
	10	$\pm 0.78n$	$\pm 1.99 m$	$\pm 5.32q$	± 14.02	± 22.30	$\pm 0.81 m$	± 2.11	$\pm 5.87 \mathrm{o}$	± 15.31	± 23.14	± 0.761	$\pm 2.57o$	± 6.380	± 16.55	± 24
	T_1	21.22	65.33	168.84	395.54	603.4	22.36	73.76	183.67	425.05	637.49	24.15	84.53	194.08	455.35	645
	11	± 0.59 jkl	$\pm 3.07 ij$	$\pm 7.53 m$	± 19.08	± 27.33	$\pm 0.65 jkl$	± 3.47	± 8.341	± 18.70	± 29.19	$\pm 0.75 ij$	± 4.06 jk	$\pm 8.91 \text{lm}$	± 21.51	± 29
	т	22.13	76.33	180.17	424.12	651.88	24.33	82.27	197.17	453.63	672.83	24.88	93.96	206.71	475.19	698
	T_2	$\pm 0.83 \text{hi}$	$\pm 2.86g$	± 6.76 kl	± 15.92	± 24.46	$\pm \ 0.91 hi$	± 3.09	\pm 7.40hijk	± 17.02	± 25.25	$\pm 0.93 \text{hi}$	$\pm 3.53h$	\pm 7.76g-k	± 17.83	± 26
	T ₃	24.10	79.67	194.90	446.46	674.47	26.07	86.17	203.98	467.37	696.77	27.05	100.83	212.70	506.74	723.
	13	$\pm 0.42e$	$\pm 1.38 ef$	$\pm 3.38 def$	± 7.73	± 11.68	$\pm 0.45g$	± 1.49	± 3.53 fghi	± 8.10	± 12.07	$\pm 0.47 fg$	$\pm 1.75 \mathrm{fg}$	\pm 3.68fghi	± 8.78	± 12
V	T_4	28.33	87.83	205.31	495.64	765.39	30.32	97.01	223.40	526.57	788.63	31.92	114.45	234.36	545.37	810
V ₃	14	$\pm 0.98 b$	$\pm 3.04c$	± 7.11c	\pm 17.17	± 26.51	$\pm 1.05 ab$	± 3.36	\pm 7.74abc	± 18.24	± 27.32	$\pm 1.11b$	$\pm 3.97 bc$	\pm 8.12bc	± 18.89	± 28
	T 5	25.42	82.82	197.21	458.30	716.77	27.32	91.82	209.20	487.38	727.71	29.95	106.70	219.65	518.13	757
	15	$\pm 1.25 d$	$\pm 4.07 d$	\pm 9.68de	± 22.49	± 35.18	$\pm 1.34 ef$	± 4.51	\pm 10.27efg	± 23.92	± 35.71	± 1.47 cd	\pm 5.24	$\pm 10.78 ef$	± 25.43	± 37
	T_6	21.97	69.63	174.24	412.87	630.54	23.23	75.50	190.61	441.94	656.13	24.88	89.68	204.24	469.88	687
	16	$\pm 0.70 hij$	$\pm 2.21 h$	$\pm 5.53 \text{lm}$	± 13.11	± 20.02	$\pm 0.74 ijk$	± 2.40	± 6.05 kl	± 14.03	± 20.84	$\pm 0.97 \text{hi}$	$\pm 2.85i$	± 6.49 ijk	± 14.92	± 21
	T ₇	23.81	80.89	187.68	449.44	694.76	26.26	88.00	200.31	470.60	705.01	28.28	100.39	210.69	496.52	739.
	17	$\pm 1.03 ef$	$\pm 3.50 de$	\pm 8.13ghij	± 19.46	± 30.08	$\pm 1.14 fg$	± 3.81	\pm 8.68ghij	± 20.38	± 30.53	$\pm 1.23 ef$	$\pm 4.35 fg$	± 9.12 fghi	± 21.50	± 32
	т	30.34	96.97	222.63	529.52	780.03	31.11	101.35	230.93	558.67	807.73	37.76	125.04	252.05	582.68	843
T_8		$\pm 1.00a$	$\pm 4.46a$	$\pm 9.84a$	± 21.80	± 34.76	± 1.04a	± 4.69	± 9.11a	± 22.10	\pm 36.20	$\pm 1.39a$	$\pm 5.21a$	$\pm 10.26a$	± 24.50	± 34
LS		**	**	*	NS	NS	**	NS	**	NS	NS	**	*	**	NS	Ν
CV (%)	1.91	2.22	2.02	1.86	1.83	2.23	2.41	2.27	2.21	2.15	2.59	2.50	2.26	2.17	2.1

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

4.3.2 Leaf area index

Leaf area index (LAI) was significantly affected by the selected varieties of wheat in the length of the research. The upper limit of LAI was recorded from V_3 and the lower limit from V_1 (Fig. 9, 10 and 11). In case of V_3 , LAI achieved the maximum value (3.48) at 55 DAS in 2018-2019. On the contrary, the minimum value (2.84) at same DAS was recorded in 2016-2017. Normally, the leaf area index enlarged gradually with age of the wheat plants and reached the peak at 55 DAS. After that, LAI decreased slowly and this happened for all varieties. Wheat plants at 55 DAS produced higher leaves/plant that resulted in the highest LAI.

All soil amendments provided significant results for leaf area index at every sampling date in each year of study. The treatment T_8 performed better, followed by T_4 than that of T_0 in respect of LAI at all DAS (Fig. 12, 13 and 14). The highest value (4.34) of LAI was recorded from T_8 and the lowest one (2.14) was from T_0 at the peak period (55 DAS) during 2018-2019 and 2016-2017, respectively. The leaf area had an ascending tendency up to 55 DAS, whereas descending nature exhibited in the next 70 DAS and 85 DAS in the three years of the experiment. Due to slow release of plant nutrient from organic manure, LAI at 85 DAS was not higher than chemical fertilizer but statistically identical with chemical fertilizer during first cropping season by T_8 (3.73) and T_4 (3.39) and also from T_8 (2.58) and T_4 (2.46) in second year.

The combination of variety and different soil amendments had significant effects on LAI at 25 DAS, 40 and 55 DAS in 2016-2017 (Table 22). Initially at 25 and 40 DAS, the comparative higher LAI was noticed in V_3T_8 (0.70 and 2.20) and at 55 DAS, V_3T_8 once more gave the comparative greater value (3.88). The variation of LAI for interaction of variety and different soil amendments was significant at 25 and 55 DAS in 2017-2018, which maintained the parallel pattern as in 2016-2017. In the year 2018-2019, among the organic amendments of soil, poultry manure (T₄) greatly influenced the LAI, however, in every case, all other treatments showed good effect than no soil amendment (T₀). In the last year 2018-2019, LAI influenced significantly as a result of selected treatments combinations. The treatment combination V_3T_8 showed the highest value (4.38) of LAI than the other treatment combinations, whereas V_2T_0 gave the lowest value (2.34) at peak period (55 DAS) of LAI. Furthermore, the above higher and lower order of LAI was noticed during rest of the sampling dates (Table 22).




Fig. 9: Effect of variety on leaf area index of wheat in 2016-2017.




Fig. 10: Effect of variety on leaf area index of wheat in 2017-2018.




Fig. 11: Effect of variety on leaf area index of wheat in 2018-2019.




Fig. 12: Effect of organic and probiotic soil amendments on leaf area index of wheat in 2016-2017.




Fig. 13: Effect of organic and probiotic soil amendments on leaf area index of wheat in 2017-2018.




Fig. 14: Effect of organic and probiotic soil amendments on leaf area index of wheat in 2018-2019.

Variety	N 87		2	016-2017					2017-2018					2018-201	9	
Treatm								Days	after sowing	(DAS)						
mean	icitt	25	40	55	70	85	25	40	55	70	85	25	40	55	70	85
	T_0	0.23	1.21	2.20	2.01	1.37	0.27	1.45	2.54	2.24	1.52	0.24	1.35	2.42	2.20	1.68
	10	$\pm 0.01 n$	$\pm 0.03 \text{lm}$	$\pm 0.10n$	± 0.10	± 0.02	$\pm 0.01 \text{op}$	± 0.06	$\pm 0.12m$	± 0.11	± 0.03	$\pm 0.01 p$	$\pm 0.07 m$	$\pm 0.12p$	$\pm 0.11q$	$\pm 0.07 n$
	т	0.24	1.30	2.29	2.08	1.56	0.28	1.54	2.65	2.41	1.50	0.31	1.60	2.70	2.66	1.84
	T_1	$\pm \ 0.01 mn$	± 0.06 klm	$\pm 0.10 mn$	± 0.09	± 0.05	$\pm 0.01 nop$	± 0.08	± 0.111	± 0.08	± 0.05	$\pm 0.01 n$	$\pm 0.06 jk$	$\pm 0.12o$	$\pm 0.12 \text{mno}$	± 0.071
	T_2	0.33	1.56	2.56	2.33	1.65	0.37	1.73	3.02	2.74	1.94	0.38	1.80	3.21	3.01	2.25
	12	$\pm 0.00 j$	± 0.05 g-j	$\pm 0.13ij$	± 0.08	± 0.02	$\pm 0.01 k$	± 0.06	$\pm 0.10i$	± 0.03	± 0.07	$\pm 0.02k$	$\pm 0.07 i$	$\pm 0.18 j$	$\pm 0.12k$	$\pm 0.11j$
	T ₃	0.42	1.64	2.78	2.60	1.89	0.47	1.99	3.13	3.05	2.17	0.53	2.24	3.38	3.12	2.48
	13	$\pm 0.01 h$	$\pm 0.03 e$ -h	$\pm 0.10 gh$	± 0.13	± 0.08	$\pm 0.01 i$	± 0.08	$\pm 0.12 h$	± 0.11	± 0.10	$\pm 0.02 h$	$\pm 0.10e$	$\pm 0.13i$	$\pm 0.13j$	$\pm 0.09 fg$
\mathbf{V}_1	T_4	0.54	1.94	3.38	2.88	2.03	0.62	2.24	3.98	3.56	2.51	0.63	2.41	4.17	3.84	2.77
\mathbf{v}_1	14	$\pm 0.01 d$	$\pm 0.07 bc$	$\pm 0.14 cd$	± 0.09	± 0.08	$\pm \ 0.01 cd$	± 0.07	$\pm 0.10c$	± 0.11	± 0.11	$\pm \ 0.03 cd$	$\pm 0.09 bc$	$\pm 0.10 \text{cd}$	$\pm 0.11d$	$\pm 0.09c$
	T_5	0.49	1.88	3.15	2.92	2.02	0.52	2.05	3.56	3.28	2.27	0.57	2.28	3.64	3.47	2.50
	15	$\pm 0.01e$	$\pm 0.07 bcd$	$\pm 0.15 ef$	± 0.08	± 0.07	$\pm 0.01 fgh$	± 0.10	$\pm 0.15 f$	± 0.09	± 0.08	$\pm 0.02 ef$	$\pm 0.06 de$	$\pm 0.17g$	$\pm 0.16g$	$\pm 0.11 f$
	T ₆	0.27	1.38	2.41	2.23	1.58	0.32	1.65	2.73	2.65	1.92	0.38	1.66	2.98	2.74	2.13
	16	$\pm 0.01 kl$	± 0.07 jkl	$\pm 0.08 kl$	± 0.07	± 0.07	$\pm 0.01 lm$	± 0.07	± 0.13 kl	± 0.05	± 0.08	$\pm 0.02k$	$\pm 0.07 j$	$\pm 0.10k$	$\pm 0.10 m$	$\pm 0.07 k$
	T_7	0.47	1.74	3.08	2.62	1.81	0.50	1.87	3.34	3.11	2.14	0.53	2.07	3.53	3.26	2.24
	17	$\pm 0.01 \text{ef}$	$\pm 0.06 d\text{-}g$	$\pm 0.12 f$	± 0.08	± 0.06	± 0.01 ghi	± 0.06	$\pm 0.12g$	± 0.06	± 0.07	$\pm 0.02 h$	$\pm 0.10 f$	$\pm 0.14 h$	$\pm 0.13i$	$\pm 0.10j$
1	T_8	0.65	2.22	3.67	3.14	2.20	0.65	2.33	4.14	3.79	2.66	0.68	2.56	4.39	4.12	2.87
	18	$\pm 0.01 b$	$\pm 0.07a$	$\pm 0.17b$	± 0.08	± 0.09	$\pm 0.01 bc$	± 0.10	$\pm 0.11 ab$	± 0.14	± 0.11	$\pm 0.02b$	$\pm 0.08a$	$\pm 0.20a$	$\pm 0.20b$	$\pm 0.11b$
	T_0	0.22	1.15	1.95	1.81	1.26	0.22	1.30	2.22	2.11	1.47	0.22	1.24	2.34	2.05	1.45
	10	$\pm 0.01 n$	$\pm 0.04 m$	$\pm 0.04o$	± 0.10	± 0.06	$\pm 0.01 q$	± 0.05	$\pm 0.10 n$	± 0.11	± 0.07	$\pm 0.01 q$	$\pm \ 0.05 n$	$\pm 0.12q$	$\pm 0.09r$	± 0.060
	T_1	0.24	1.25	2.27	2.0270	1.47	0.27	1.44	2.64	2.22	1.61	0.28	1.40	2.75	2.57	1.72
	11	$\pm \ 0.01 mn$	$\pm 0.051 \text{m}$	$\pm 0.10 mn$	± 0.08	± 0.08	$\pm 0.01 nop$	± 0.06	± 0.131	± 0.10	± 0.07	$\pm 0.01o$	$\pm 0.07 m$	$\pm 0.12o$	$\pm 0.08 op$	$\pm \ 0.08 mn$
	T_2	0.28	1.30	2.46	2.18	1.53	0.33	1.60	2.86	2.59	1.74	0.35	1.57	2.95	2.66	1.84
	12	$\pm 0.00 k$	$\pm 0.05 klm$	$\pm 0.07 jk$	± 0.09	± 0.06	± 0.011	± 0.07	$\pm 0.10j$	± 0.11	± 0.05	± 0.011	$\pm \ 0.07 k$	± 0.10 kl	$\pm 0.11 \text{mn}$	± 0.071
	T ₃	0.39	1.56	2.72	2.24	1.59	0.43	1.65	3.10	2.72	1.94	0.44	1.84	3.24	2.88	2.11
	13	$\pm 0.00i$	± 0.08 g-j	$\pm 0.10 h$	± 0.11	± 0.05	$\pm 0.01 j$	± 0.06	$\pm 0.13 hi$	± 0.10	± 0.06	$\pm 0.02i$	$\pm 0.04 \text{hi}$	$\pm 0.12 j$	± 0.131	$\pm 0.06k$
V_2	T_4	0.52	1.87	3.33	2.79	2.03	0.60	2.17	3.85	3.14	2.28	0.63	2.35	4.10	3.49	2.62
v ₂	14	$\pm 0.00 d$	$\pm \ 0.05 bcd$	$\pm 0.13 d$	± 0.11	± 0.10	$\pm 0.01 d$	± 0.06	$\pm 0.19 d$	± 0.13	± 0.06	$\pm 0.0361 d$	$\pm \ 0.11 cd$	$\pm 0.16d$	$\pm 0.11g$	$\pm 0.10e$
	T_5	0.49	1.94	3.13	2.85	1.94	0.53	2.04	3.52	3.09	2.11	0.55	2.11	3.75	3.13	2.31
	15	$\pm 0.01e$	$\pm 0.09 bc$	$\pm 0.14 ef$	± 0.12	± 0.10	$\pm 0.01 \text{ef}$	± 0.08	$\pm 0.15 f$	± 0.13	± 0.11	$\pm 0.02g$	$\pm \ 0.07 f$	$\pm 0.16 f$	$\pm 0.14j$	$\pm 0.07 ij$
	т	0.26	1.41	2.40	2.18	1.49	0.30	1.46	2.73	2.38	1.64	0.33	1.50	2.88	2.64	1.79
	T_6	$\pm 0.01 klm$	± 0.07 jkl	$\pm 0.07 kl$	± 0.09	± 0.08	$\pm 0.01 mn$	± 0.06	$\pm 0.08 kl$	± 0.09	± 0.04	$\pm \ 0.01 mn$	± 0.071	$\pm 0.14 lm$	± 0.10 no	$\pm 0.08 \text{lm}$
	T	0.46	1.67	3.05	2.60	1.73	0.49	1.76	3.28	2.85	1.97	0.52	1.87	3.56	2.99	2.14
	T ₇	± 0.01 fg	$\pm 0.08\text{e-h}$	$\pm 0.13 f$	± 0.12	± 0.07	± 0.02 hi	± 0.08	$\pm 0.14g$	± 0.12	± 0.09	$\pm 0.02h$	± 0.10 hi	± 0.14 gh	$\pm 0.11k$	$\pm 0.06k$
	m	0.57	2.04	3.64	2.95	2.07	0.63	2.23	4.05	3.41	2.35	0.65	2.43	4.24	3.73	2.72
	T_8	$\pm 0.01c$	± 0.10ab	± 0.18b	± 0.13	± 0.10	$\pm 0.01 \text{cd}$	± 0.02	$\pm 0.11 bc$	± 0.13	± 0.11	$\pm 0.02c$	$\pm 0.10b$	± 0.16bc	± 0.16e	± 0.13 cd

Table 22: Interaction effects of variety and soil amendments on leaf area index of wheat.

Contd...

Variety	8.		2	016-2017					2017-2018					2018-2019)	
Treatm								Days	after sowing	(DAS)						
Treatin	ient -	25	40	55	70	85	25	40	55	70	85	25	40	55	70	85
	To	0.22	1.30	2.26	2.03	1.49	0.25	1.46	2.64	2.31	1.61	0.24	1.41	2.78	2.52	1.71
	10	$\pm 0.01 n$	± 0.03 klm	$\pm 0.09 mn$	± 0.06	± 0.05	$\pm 0.01 p$	± 0.06	± 0.121	± 0.07	± 0.05	$\pm 0.01 p$	$\pm 0.07 m$	$\pm 0.12 no$	$\pm 0.10 p$	$\pm 0.07 mn$
	T_1	0.24	1.44	2.34	2.05	1.52	0.29	1.78	2.71	2.47	1.85	0.33	1.84	2.86	2.61	1.87
	11	$\pm 0.01 \text{mn}$	$\pm 0.05 ijk$	$\pm 0.08 ml$	± 0.09	± 0.04	$\pm \ 0.01 mno$	± 0.02	$\pm 0.11 kl$	± 0.11	± 0.05	$\pm 0.01 \text{lm}$	$\pm 0.04 hi$	$\pm 0.14 \text{mn}$	$\pm 0.12 no$	± 0.091
	T ₂	0.35	1.62	2.61	2.56	1.74	0.41	1.81	3.03	2.95	2.00	0.42	1.97	3.23	3.13	2.39
	12	$\pm 0.01 j$	$\pm 0.03 ext{f-i}$	± 0.11i	± 0.08	± 0.05	$\pm 0.01 j$	± 0.09	$\pm 0.15i$	± 0.13	± 0.08	$\pm 0.02j$	$\pm 0.05g$	$\pm 0.12j$	± 0.11j	± 0.10 hi
	T ₃	0.49	1.76	2.88	2.75	2.05	0.55	2.00	3.27	3.06	2.21	0.55	2.33	3.41	3.28	2.53
	13	$\pm 0.01e$	± 0.03 c-f	± 0.15g	± 0.07	± 0.09	$\pm 0.03 \text{ef}$	± 0.09	$\pm 0.18g$	± 0.08	± 0.11	$\pm 0.03g$	$\pm 0.11d$	± 0.13i	$\pm 0.13 hi$	$\pm 0.07 f$
V ₃ T ₄	0.68	1.92	3.47	3.07	2.16	0.68	2.36	4.02	3.65	2.57	0.69	2.43	4.28	3.97	2.85	
V ₃	14	± 0.01a	$\pm 0.08 bcd$	$\pm 0.14c$	± 0.10	± 0.05	$\pm 0.015 ab$	± 0.02	$\pm 0.17c$	± 0.12	± 0.06	$\pm 0.02b$	$\pm 0.11b$	$\pm 0.17b$	$\pm 0.17c$	$\pm 0.07 b$
	т	0.47	1.82	3.20	2.88	2.00	0.56	2.15	3.76	3.35	2.320	0.58	2.30	3.88	3.58	2.64
	T ₅	$\pm 0.01 \text{ef}$	± 0.08 cde	± 0.15e	± 0.05	± 0.04	$\pm 0.01e$	± 0.09	$\pm 0.18e$	± 0.06	± 0.04	$\pm 0.02e$	± 0.11 de	$\pm 0.178e$	$\pm 0.13 f$	± 0.06de
	T	0.28	1.57	2.44	2.24	1.60	0.32	1.83	2.78	2.66	1.95	0.38	1.91	2.89	2.85	2.27
	T_6	$\pm 0.00k$	± 0.05 g-j	± 0.11 kl	± 0.08	± 0.04	$\pm 0.01 \text{lm}$	± 0.03	± 0.10jk	± 0.12	± 0.09	$\pm 0.02k$	± 0.09 gh	$\pm 0.11 lm$	± 0.131	± 0.05j
	T	0.44	1.54	3.08	2.70	1.81	0.53	1.99	3.50	3.17	2.21	0.56	2.13	3.63	3.36	2.41
	T_7	$\pm 0.01 gh$	± 0.03 hij	$\pm 0.10 f$	± 0.10	± 0.03	$\pm 0.01 fg$	± 0.07	$\pm 0.14 f$	± 0.04	± 0.06	$\pm 0.03 gf$	$\pm 0.09 f$	$\pm 0.14g$	$\pm 0.17 h$	± 0.06 gh
	T	0.70	2.20	3.88	3.27	2.23	0.70	2.45	4.19	3.85	2.71	0.74	2.59	4.38	4.24	2.99
	T_8	$\pm 0.01a$	$\pm 0.07a$	$\pm 0.17a$	± 0.10	± 0.08	$\pm 0.01a$	± 0.03	$\pm 0.18a$	± 0.14	± 0.04	$\pm 0.03a$	± 0.13a	± 0.19a	± 0.18a	$\pm 0.09a$
LS		**	*	**	NS	NS	**	NS	**	NS	NS	**	**	**	**	**
CV (%	%)	3.70	6.39	2.18	6.29	6.59	4.08	6.38	1.63	5.81	6.43	2.35	2.40	1.41	1.64	2.13

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

4.3.3 Crop growth rate

In this present study, crop growth rate (CGR) was found as significant at all sampling dates by all varieties during 2016-2017 (Fig. 15). From 25-40 to 55-70 DAS, CGR increased sharply and then went down slowly. Where the maximum (16.97 g/m²/d) and the minimum (15.83 g/m²/d) CGR recorded from V₃ and V₂, respectively when the wheat plant reached the peak growth (55-70 DAS). The maximum and statistically equal crop growth rate derived from V₃ (3.50 g/m²/d) and V₁ (3.45 g/m2/d) at 25-40 DAS. Wheat plant also showed significant variation in CGR caused by varietal influence in 2017-2018 (Fig. 16), where descending order of CGR was found asV₃>V₁>V₂. Here V₃ variety had produced the greater CGR (17.97 g/m²/d) and less CGR (17.09 g/m²/d) recorded by V₂. A similar tendency was obtained from the selected varieties of wheat in 2018-2019, which refers to the year 2016-2017. But only at 40-55 DAS, V₃ (7.51 g/m²/d) and V₁ (7.47 g/m²/d) showed the similarity of significance (Fig. 17).

The data indicated that soil amendments had a significant effect on wheat CGR at all DAS in the whole experimental period. Most of the sampling dates revealed that the treatment T_8 had the highest activity in CGR followed in sequence by T_4 , T_5 , T_7 , T_3 , T_2 , T_6 and T_1 and the lowest was in T_0 (Fig. 18, 19 and 20). At the end (70-85 DAS) of CGR sampling, T_4 showed the greater influence. Through the three years of study maximum CGR value (21.24 g/m²/d) was reported at 55-70 DAS by T_8 and T_4 (19.89 g/m²/d) in 2018-2019. Whereas the minimum value (16.386 g/m²/d) was recorded at the same DAS from T_0 in 2018-2019. Variation in CGR among the soil amendments was exhibited to a higher extent at 55-70 DAS and afterwards decreased due to leaf senescence and tiller drying.

The crop growth rate showed significant variation at various combinations of treatments and varieties during 2016-2017 at all DAS. Increasing development of crop growth rate was noticed from initial to till 40-55 DAS, where the combination V_3T_8 (20.46 g/m²/d), V_1T_8 (18.86 g/m²/d) and V_2T_8 (18.03 g/m²/d) produced greater CGR, while the lowest value was observed in V_2T_0 (13.36 g/m²/d), V_1T_0 (13.72 g/m²/d) and V_3T_0 (14.37 g/m²/d).

With the statistical significance, maximum CGR were recorded from V_3T_8 (21.85 g/m²/d) followed by V_1T_8 (20.62 g/m²/d) and V_2T_8 (19.94 g/m²/d) at 55-70 DAS in 2017-2018. Comparable results were found in 2018-2019 with 2016-2017 regarding CGR from starting to till 55-70 DAS. But the maximum increased values of CGR was documented with statistical identity in case of V_3T_8 (22.04 g/m²/d) and V_1T_8 (21.72 g/m²/d) at 55-70 DAS, oppositely the minimum decreased of CGR was showed statistical identity in case of V_3T_4 (17.67 g/m²/d) and V_3T_8 (17.37 g/m²/d) at 70-85 DAS (Table 23).




Fig. 15: Effect of variety on crop growth rate of wheat in 2016-2017.




Fig. 16: Effect of variety on crop growth rate of wheat in 2017-2018.




Fig. 17: Effect of variety on crop growth rate of wheat in 2018-2019.




Fig. 18: Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2016-2017.




Fig. 19: Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2017-2018.




Fig. 20: Effect of organic and probiotic soil amendments on crop growth rate of wheat in 2018-2019.

$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Var	ioty &		201	6-2017			201	7-2018			201	8-2019	
$ \begin{array}{c c c c c c c c c c c c c c c c c c c $		-					•	Days after so	owing (DAS)					
$ \begin{array}{c} 1_{0} & \pm 0.091 & \pm 0.25m & \pm 0.67p & \pm 0.54m0 & \pm 0.07p & \pm 0.23k & \pm 0.58gr & \pm 0.59m & \pm 0.13m & \pm 0.25k & \pm 0.83j & \pm 0.51m \\ T_{1} & \pm 0.15i & \pm 0.32j & \pm 0.75k & \pm 0.67m0 & \pm 0.10n & \pm 0.31i & \pm 0.75gr & \pm 0.66l & \pm 0.171 & \pm 0.34i & \pm 0.74ij & \pm 0.61kl \\ T_{2} & \pm 0.13i & \pm 0.27i & \pm 0.61jm & \pm 0.54efg & \pm 0.07j & \pm 0.25k & \pm 0.75m & \pm 0.66l & \pm 0.171 & \pm 0.34i & \pm 0.74ij & \pm 0.61kl & \pm 0.72ir & \pm 0.66l & \pm 0.171 & \pm 0.34i & \pm 0.74ij & \pm 0.61kl & \pm 0.72ir & \pm 0.66l & \pm 0.171 & \pm 0.34i & \pm 0.74ij & \pm 0.61kl & \pm 0.72ir & \pm 0.66l & \pm 0.171n & \pm 0.32ir & \pm 0.67m & \pm 0.54efg & \pm 0.07ir & \pm 0.25k & \pm 0.77m & \pm 0.32ir & \pm 0.16kj & \pm 0.77m & \pm 0.32ir & \pm 0.16kj & \pm 0.72ir & \pm 0.66l & \pm 0.75r & \pm 0.66kl & \pm 0.75r & \pm 0.68kl & \pm 0.75r & \pm 0.58kl & \pm 0.95r & \pm 0.75r & \pm 0.58kl & \pm 0.75r & \pm 0.75$	1100	atment	25-40	40-55	55-70	70-85	25-40	40-55	55-70	70-85	25-40	40-55	55-70	70-85
$ \begin{array}{c} \begin{array}{c} 10091 & \pm 0.23m & \pm 0.07p & \pm 0.34m & \pm 0.07p & \pm 0.23k & \pm 0.03p & \pm 0.03m & \pm 0.0$		To												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		-0												
$ \begin{array}{c} 1.5 \\ V_1 \\ V_1 \\ V_1 \\ V_1 \\ V_2 \\ V_3 \\ V_3 \\ V_1 \\ V_1 \\ V_1 \\ V_1 \\ V_1 \\ V_1 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_2 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_3 \\ V_2 \\ V_3 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_3 \\ V_2 \\ V_3 \\ V_4 \\ V_3 \\ V_4 \\ V_3 \\ V_4 $		T ₁												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		11												
$ \begin{array}{c} 1 & 0.14g & \pm 0.21i & \pm 0.01j & \pm 0.03eq & \pm 0.01j & \pm 0.02eq & \pm 0.04eq & \pm 0.03eq & \pm 0.05eq & \pm 0.03eq & \pm 0.0$		Т												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		12				U	5			5				
$ \begin{array}{c} \mathbb{V}_{1} & \mathbb{T}_{4} & \frac{4.08}{10} & \frac{10.52e1}{10.57e1} & \frac{10.08}{10.57e1} & \frac{10.50e}{10.57e1} & \frac{10.50e}{10.57e1} & \frac{10.50e}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.52e}{10.52e} & \frac{10.7e1}{10.52e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.55e} & \frac{10.7e1}{10.55e} & \frac{10.7e1}{10.55e} & \frac{10.7e1}{10.55e} & \frac{10.7e1}{10.55e} & \frac{10.7e1}{10.53e} & \frac{10.7e1}{10.55e} $		Т												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		13												
$ V_2 = V_1 = \frac{10.20cd}{15} \pm 0.30cd \pm 0.87c} \pm 0.7bd \pm 0.12bd \pm 0.34bc \pm 0.83cd \pm 0.79b \pm 0.24c \pm 0.38bd \pm 0.92c \pm 0.75d \\ \pm 0.12d + 0.26d \pm 0.50de \pm 0.44d \pm 0.09fg \pm 0.21ef \pm 0.49hi \pm 0.47efg \pm 0.14d \pm 0.22fg \pm 0.53e \pm 0.46c \\ \pm 0.285 - 6.60 - 15.74 - 14.08 - 3.27 - 6.81 - 15.96 - 14.60 - 3.78 - 7.51 - 17.22 - 14.44 \\ \pm 0.212 \pm 0.26i \pm 0.60i + 0.55j + 0.10m \pm 0.25i \pm 0.59hop \pm 0.55hi j \pm 0.14k \pm 0.28fg \pm 0.65hi j \pm 0.54c \\ \pm 0.14d \pm 0.26i \pm 0.60i + 0.55j + 0.10m \pm 0.25i \pm 0.59hol \pm 0.55hi j \pm 0.14k \pm 0.28fg \pm 0.65hi j \pm 0.576 \\ \pm 0.14de \pm 0.25i \pm 0.59def \pm 0.33ef \pm 0.13gh \pm 0.23gh \pm 0.59ki \pm 0.53de \pm 0.16g + 0.26fg \pm 0.62fg \pm 0.55c \\ T_8 - 4.32 - 8.273 - 18.86 - 16.43 - 4.11 - 8.79 - 20.62 - 15.13 - 5.40 - 8.37 - 21.72 - 15.15 \\ \pm 0.22ab \pm 0.43a \pm 0.95b \pm 0.74cd \pm 0.17b \pm 0.41a \pm 0.93b \pm 0.81ef \pm 0.27b \pm 0.38a \pm 0.63a \pm 0.08a \\ -7.8 - 0.22ab \pm 0.43a \pm 0.95b \pm 0.74cd \pm 0.17b \pm 0.41a \pm 0.93b \pm 0.81ef \pm 0.27b \pm 0.38a \pm 0.38a \pm 0.69ef \\ T_1 - 2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -2.39 - 6.01 - 14.84 - 13.01 - 2.85 - 6.22 - 15.26 - 11.90 - 3.14 - 6.75 - 16.17 - 12.12 \\ -3.12 - 0.04k \pm 0.30h \pm 0.64hm \pm 0.376h \pm 0.028g \pm 0.028g \pm 0.376n \pm 0.44hk \pm 0.456h - 4.049 - 7.44 - 10.64 - 4.045 + 0.556h - 4.0370h \pm 0.65kh \pm 0.556h - 4.0370h \pm 0.65kh \pm 0.576h + 0.326h - 4.0370h \pm 0.44hk \pm 0.55hh \pm 0.316h - 4.0459h + 0.55hh + 0.306h + 0.056h + 0.326h + 0.44hh + 0.18hh + 0.71hh + 0.365h + 0.326h $	V.	Т.												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	• 1	14												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	3.87				4.11			15.06	4.95	7.52		15.88
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		15												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	2.85	6.60	15.74	14.08	3.27	6.81	15.96	14.60	3.78	7.51	17.22	14.44
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		16	± 0.12i	± 0.26i	$\pm 0.60i$ -l	± 0.55j-m	$\pm 0.10m$	± 0.25i	± 0.59nop	± 0.53 hij	$\pm 0.14k$	± 0.28 fg	± 0.65 hij	± 0.54ij
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	4.00	6.61	16.88	15.19	3.99	7.35	16.94	15.29	4.47		17.85	15.76
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		T_7	± 0.14 cde	± 0.25i	± 0.59def	± 0.53ef	± 0.13 ghi	± 0.23 gh	± 0.59jkl	± 0.53de	± 0.16g	± 0.26fg	± 0.62fg	± 0.55cd
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	4.32	8.273	18.86	16.43	4.41	8.79	20.62	15.13	5.40	8.37	21.72	15.15
$ V_{2} \begin{matrix} I_{0} \\ \pm 0.04kl \\ \pm 0.11n \\ \pm 0.30p \\ \pm 0.24no \\ \pm 0.08lp \\ \pm 0.10l \\ \pm 0.27s \\ \pm 0.27s \\ \pm 0.27n \\ \pm 0.27n \\ \pm 0.05o \\ \pm 0.12k \\ \pm 0.12k \\ \pm 0.31l \\ \pm 0.31l \\ \pm 0.31l \\ \pm 0.31l \\ \pm 0.24m \\ \pm 0.24m \\ \pm 0.27s \\ \pm 0.27s \\ \pm 0.27n \\ \pm 0.05o \\ \pm 0.12k \\ \pm 0.25n \\ \pm 0.49n \\ \pm 0.12m \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.49n \\ \pm 0.12m \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.48m \\ \pm 0.49n \\ \pm 0.12m \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.48m \\ \pm 0.12m \\ \pm 0.25i \\ \pm 0.61k \\ \pm 0.61k \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.46m \\ \pm 0.44k \\ \pm 0.15k \\ \pm 0.15k \\ \pm 0.31e \\ \pm 0.65k \\ \pm 0.55e \\ \pm 0.55e \\ \pm 0.55e \\ \pm 0.55e \\ \pm 0.44k \\ \pm 0.19k \\ \pm 0.38h \\ \pm 0.44k \\ \pm 0.17hi \\ \pm 0.19k \\ \pm 0.25k \\ \pm 0.38h \\ \pm 0.38h \\ \pm 0.17hi \\ \pm 0.19k \\ \pm 0.28k \\ \pm 0.74m \\ \pm 0.19k \\ \pm 0.28k \\ \pm 0.73op \\ \pm 0.44k \\ \pm 0.73op \\ \pm 0.44k \\ \pm 0.15k \\ \pm 0.38hi \\ \pm 0.18k \\ \pm 0.27k \\ \pm 0.38hi \\ \pm 0.38hi \\ \pm 0.20k \\ \pm 0.20k \\ \pm 0.38hi \\ \pm 0.20k \\ \pm 0.20k \\ \pm 0.20k \\ \pm 0.42m \\ \pm 0.38hi \\ \pm 0.17hi \\ \pm 0.18k \\ \pm 0.27k \\ \pm 0.38hi \\ \pm 0.20k \\ \pm 0.38hi \\ \pm 0.20k \\ \pm 0.38hi \\ \pm 0.20k \\ \pm 0.38hi \\ \pm 0.42m \\ \pm 0.38hi \\ \pm 0.17hi \\ \pm 0.21k \\ \pm 0.28k \\ \pm 0.30k \\ \pm 0.38hi \\ \pm 0.42m \\ \pm 0.38hi \\ \pm 0.27k \\ \pm 0.20k \\ \pm 0.20k \\ \pm 0.38hi \\ \pm 0.39k \\ \pm 0.30k \\ \pm 0.74k \\ \pm 0.27k \\ \pm 0.20k \\ \pm 0.36k \\ \pm 0$		18	$\pm 0.22ab$	± 0.43a	$\pm 0.95b$	± 0.74cd	$\pm 0.17 bc$	± 0.41a	$\pm 0.93b$	$\pm 0.81 \text{ef}$	$\pm 0.27b$	$\pm 0.38a$	± 0.38a	± 0.69ef
$ V_{2} \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	1.83	4.92	13.36	13.28	2.13	5.23	14.75	11.85	2.46	5.77	15.53	11.82
$ V_{2} \begin{array}{ccccccccccccccccccccccccccccccccccc$		10	± 0.04 kl	± 0.11n	± 0.30p	± 0.24no	± 0.081p	± 0.101	$\pm 0.27s$	± 0.27n	$\pm 0.05o$	± 0.12k	± 0.311	± 0.24m
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T	2.39	6.01	14.84	13.01		6.22	15.26	11.90	3.14	6.75	16.17	12.12
$ V_{2} \begin{array}{cccccccccccccccccccccccccccccccccccc$		\mathbf{I}_1	$\pm 0.11j$	± 0.23k	± 0.57no	$\pm 0.45o$	$\pm 0.07o$	± 0.23j	± 0.557rs	± 0.49n	± 0.12m	± 0.25i	± 0.61k	± 0.46lm
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		т	3.12	6.85	15.27	14.66	3.64	7.48	15.86	14.17	3.91	7.63	16.06	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		1 ₂	± 0.15h	± 0.30h	± 0.641mn	± 0.57f-i	± 0.12k	± 0.28 fg	± 0.73 opq	± 0.44jk	± 0.15 jk	± 0.31ef	± 0.65k	± 0.55ef
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T	3.69	7.12	16.10	14.46	3.93	7.46	16.48	14.56	4.49	7.54	17.48	14.95
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		13	$\pm 0.10g$	± 0.19fg	± 0.43hi	± 0.38 h-k	± 0.17 hij	± 0.19fg	± 0.421mn	± 0.38hij	± 0.18 g	± 0.20fg	± 0.45ghi	± 0.39efg
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T	3.96	7.33	17.18	16.46	4.18	8.07	18.26	15.19		8.02	19.10	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	V_2	\mathbf{I}_4	± 0.18 c-f	± 0.35ef	± 0.79de	± 0.66cd	± 0.15ef	± 0.32d	± 0.74ef	± 0.71ef	± 0.21d	± 0.35cd	± 0.83d	± 0.67de
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		-					4.11		17.64		4.67	7.94		
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T_5												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		T_6												
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		-												
T 4.14 8.00 18.03 15.26 4.37 8.27 19.34 14.85 5.39 8.10 19.97 14.79														
		T_8	± 0.15 bc	± 0.29b	$\pm 0.67c$	± 0.51e	± 0.15 bcd	± 0.28cd	$\pm 0.82c$	± 0.29e-h	± 0.17b	± 0.28bc	$\pm 0.69c$	± 0.70f-i

Table 23: Interaction effects of variety and soil amendments on crop growth rate of wheat $(g/m^2/d)$.

Contd...

			201	5-2017			201	7-2018			201	8-2019	
	ety &						Days after so	owing (DAS)					
Treat	tment	25-40	40-55	55-70	70-85	25-40	40-55	55-70	70-85	25-40	40-55	55-70	70-85
	Ŧ	2.01	5.48	14.37	13.65	2.15	6.20	15.57	12.92	2.98	6.29	16.78	12.54
	T_0	$\pm 0.09k$	± 0.251	$\pm 0.63o$	$\pm 0.52 klm$	$\pm 0.07 p$	$\pm 0.22j$	$\pm 0.58 pqr$	$\pm 0.55 m$	$\pm 0.12n$	$\pm 0.25 j$	$\pm 0.68j$	$\pm 0.51 k$
	т	2.94	6.90	14.96	14.02	3.43	7.33	16.09	14.16	4.02	7.30	17.42	12.68
	T_1	± 0.16 hi	$\pm 0.34 gh$	$\pm 0.74 mn$	$\pm 0.65 klm$	± 0.091	$\pm \ 0.32 gh$	± 0.70 nop	$\pm 0.53 jk$	$\pm 0.18ij$	$\pm 0.34 gh$	$\pm 0.81 ghi$	$\pm 0.38k$
	т	3.61	6.92	16.26	15.18	3.86	7.66	17.10	14.61	4.61	7.52	17.90	14.86
	T_2	$\pm 0.15g$	$\pm 0.29 gh$	± 0.64 ghi	$\pm 0.55 ef$	$\pm 0.13ij$	$\pm 0.26 ef$	± 0.61 ijk	± 0.57 g-j	$\pm 0.17 fg$	$\pm 0.28 fg$	$\pm 0.67 fg$	$\pm 0.56 \text{f-i}$
	т	3.70	7.68	16.77	15.20	4.01	7.85	17.56	15.29	4.92	7.46	19.60	14.46
	T ₃	$\pm 0.07 fg$	$\pm 0.14c$	$\pm 0.30 efg$	$\pm 0.27 ef$	$\pm 0.12 gh$	$\pm 0.13e$	± 0.29 ghi	$\pm 0.26 de$	$\pm 0.09 d$	$\pm 0.13 fg$	$\pm 0.34c$	± 0.25 hij
V.	V ₃ T ₄	3.97	7.83	19.36	17.98	4.45	8.43	20.21	17.47	5.50	7.99	20.73	17.67
v 3		$\pm 0.15c$ -f	$\pm 0.29 bc$	$\pm 0.70b$	$\pm 0.61a$	$\pm 0.15b$	$\pm 0.27 bc$	$\pm 0.67b$	$\pm 0.62a$	$\pm 0.19b$	$\pm 0.28cd$	$\pm 0.72b$	$\pm 0.61a$
	т	3.83	7.63	17.41	17.23	4.30	7.83	18.55	16.02	5.12	7.53	19.90	15.90
	T_5	$\pm 0.21 efg$	$\pm 0.38cd$	± 0.91d	$\pm 0.79b$	$\pm \ 0.12 cde$	$\pm 0.37e$	$\pm 0.85 de$	$\pm 0.85c$	$\pm 0.25c$	$\pm 0.37 fg$	$\pm 0.80c$	$\pm 0.69c$
	T	3.18	6.97	15.91	14.51	3.48	6.67	16.76	14.28	4.32	7.64	17.71	14.52
	T_6	$\pm 0.11h$	$\pm 0.24 gh$	± 0.53 h-k	$\pm 0.45 g$ -k	± 0.081	$\pm 0.22 ef$	$\pm 0.51 klm$	± 0.46 ijk	$\pm 0.14 h$	$\pm 0.24 ef$	$\pm 0.56 fgh$	± 0.46 ghi
	T_7	3.81	7.12	17.45	16.35	4.12	7.49	18.02	15.63	4.81	7.35	19.06	15.88
	17	$\pm 0.18 efg$	$\pm 0.32 fg$	± 0.78d	$\pm 0.68d$	$\pm 0.13 fg$	$\pm 0.31 fg$	$\pm 0.76 fg$	± 0.71 cd	$\pm 0.21 de$	$\pm \ 0.32 gh$	$\pm 0.83d$	$\pm 0.51c$
	т	4.44	8.38	20.46	16.94	4.68	8.64	21.85	16.60	5.82	8.46	22.04	17.37
	T ₈	$\pm 0.19a$	$\pm 0.35a$	$\pm 0.88a$	$\pm 0.67 bc$	$\pm 0.16a$	$\pm 0.34ab$	± 0.95a	$\pm 0.49b$	$\pm 0.21a$	± 0.34a	$\pm 0.83a$	$\pm 0.66a$
L	S	**	**	**	**	**	**	**	**	**	**	**	**
CV	/ %	4.03	1.79	1.95	2.02	2.05	1.77	1.73	1.68	2.01	1.64	1.67	1.66

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

4.4 Yield and yield contributing characters

4.4.1 Plant height

Except in 2017-2018, the height of wheat plants varied significantly due to varietal effect. In 2016-2017, the tallest plant (83.16 cm) was produced from V_3 and the lowest most (80.24 cm) recorded from V_1 . Here the V_2 had the intermediate value (81.62 cm) regarding plant height in between V_3 and V_1 . Like the first year (2016-2017), corresponding results were found in the last year (2018-2019) where the highest and lowest value represented as 86.88 and 84.42 cm from the variety V_3 and V_2 , respectively (Table 24).

With the significant variation, the treatment T_8 demonstrated the maximum and T_0 presented the minimum plant height among the treatments in the whole investigation period. Initially (2016-2017), treatment T_8 performed as the best (89.62 cm) plant height producer, while T_0 gave the least (75.76 cm). Rest of the treatments were influenced the plant height as descending order T_4 (87.19 cm) > T_5 (84.02 cm) > T_7 (82.08 cm) > T_3 (80.89 cm) > T_2 (79.35 cm) > T_6 (78.46 cm) > T_1 (77.61 cm) though T_3 and T_7 showed the statistical similarity. Result as like previous year, though T_8 and T_0 treatment activity showed greater (91.38 cm) and less (74.63 cm) plant height in 2017-2018, but had the same statistical interrelation among T_8 and T_4 , T_6 and T_2 . A little bit increase was found in plant height compared with the previous year from the different soil amendments in 2018-2019. At this time, the maximum (93.81 cm) and minimum (78.45 cm) plant height noticed with T_8 and T_0 , correspondingly (Table 25).

The Table 26 makes clear the varietal and treatment combination effect on the plant height under the investigation period from 2016-2017 to 2018-2019. The highest and the lowest plant height was obtained from V_3T_8 (95.20 cm) in 2018-2019 and V_1T_0 (74.21 cm) in 2017-2018 in that order.

4.4.2 Total plant/m²

Statistical difference was noticed by variety regarding total plant/m² at harvest (Table 24). Among the varieties, V_3 exhibited the higher frequency of total plant/m² which was followed by V_1 and V_2 from the first year to last year in this research. In 2016-2017, the variety V_3 (182.22) and V_1 (177.00) showed parallel activity in terms of total plant/m²

where V_2 had the second score (173.00). As cropping season 2016-2017, V_3 produced the maximum (187.67) total plant/m² having the statistical equality with V_1 (185.11), whereas V_2 (179.93) demonstrated the second highest during 2017-2018 (Table 24). The above trend of varietal performance was repeated in 2018-2019 with the greater plant/m² from V_3 (193.56) and lower plant/m² by V_1 (189.19).

Total plant/m² at harvest varied significantly due to different soil amendments all over the study period. The highest frequency of total plant/m² was found in T₈ (199.11) and the lowest result was recorded in T₀ (158.67) in 2016-2017 (Table 25). Where the next highest value was noticed from T₄ (191.56) and T₅ (185.00) had the statistical likeness. In the year 2017-2018, total plant/m² exhibited the matching trend due to various soil amendments as reported in 2016-2017. Where the highest (207.78) plant/m² produced by T₈ and had similar identity with the second most from T₄ (202.89). The lowest total plant/m² (163.89) was noticed in T₀ coupled with T₁ (170.22). Among the different soil amendments T₈ (216.11) performed superior results besides T₄ (209.89) to other treatments in 2018-2019. Rest of the treatments were showed also significant difference ranked as descending by T₅ (199.78), T₇ (190.22), T₃ (187.22), T₂ (179.00), T₆ (177.67) and T₁ (173.44), whereas the lowest one in T₀ (159.00).

Total plant/m² was not differed significantly due to interaction of variety and soil amendments during the whole period of the study. Although there was no significant variation among various aforesaid treatment combinations, numerical variations were observed from that combination. Here the maximum (225.00) and the minimum (153.00) values were recorded from the V_3T_8 and V_2T_0 in the 2018-2019 and 2016-2017 research period, respectively (Table 26).

4.4.3 Total tiller/plant

Considering the variety as a factor in this experiment, there was statistical significant deviation for total tiller/plant (Table 24). In 2016-2017, the highest total tiller/plant (3.11) was produced from V_3 and the loweest most (2.86) recorded from V_1 . As previous year 2016-2017, V_3 produced the maximum total tiller/plant (3.19) and V_1 demonstrated

minimum total tiller/plant (2.98) in the wheat growing period 2017-2018. The above trend of varietal performance was repeated in 2018-2019 with the greater and lower value of total tiller/plant from V_3 (3.51) and V_1 (3.15), correspondingly.

The rest of the treatments had also shown significant variations for total tiller/plant all over the study period. Though some treatments were acted as statistically similar, but the treatment T_8 and T_0 produced the maximum (3.67) and minimum (2.24) total tiller/plant, respectively in the period of 2016-2017. As a similar consequence on the number of total tiller/plant was reported in the year 2017-2018 compared to previous year. In this year, T_8 produced the highest total tiller/plant (3.80) and T_4 showed the second most total tiller/plant (3.62) but the least (2.32) in T_0 . During 2018-2019, treatment T_8 and T_0 furthermore demonstrated the greater (4.13) and the less (2.23) value of total tiller/plant (Table 25).

The number of total tiller/plant was insignificant at various combinations of varieties and treatments in all the three years of experiment. The above stated varieties and soil amendments showed great influence individually on total tiller/plant with mostly gradual increase. Although the interaction effect was insignificant but numerical values of this parameter greatly varied, where the maximum (4.40) and the minimum (2.27) number were obtained from V_3T_8 and V_2T_0 in 2018-2019 and in 2016-2017, correspondingly (Table 26).

4.4.4 Effective tiller/plant

The table 24 describes the significant variation among the selected varieties of wheat regarding effective tiller/plant from first year to third year on this study. A little bit difference was found among the varieties in the year of 2016-2017, where V₃ gave the maximum (2.93) and V₁ gave the minimum (2.71) value of effective tiller/plant. In the second year 2017-2018, varietal performance was presented as like previous year and had the greater value of effective tiller/plant from V₃ (3.06) and V₂ (2.92). In connection with the preceding year during the 2018-2019 cropping season, V₃ and V₂ also produced the highest (3.25) and lowest (2.98) number of effective tiller/plant with statistical identity in between V₂ (3.07) and V₁ (2.98).

Under this study, recorded results on effective tiller/plant showed significant difference by the influence of treatments. Out of different soil amendments, chemical fertilizer (T₈) and poultry manure (T₄) performed as superior to the rest of the treatments mainly unamend (T₀) all over the research period (Table 25). On the basis of the above statement, T₈ was produced the maximum number of effective tiller/plant (3.49) followed by T₄ (3.40), whereas the minimum value (2.07) was given by T₀ in 2016-2017. During 2017-2018, treatments were exhibited approximately equal performance with slight increase of effective tiller/plant obtained by treatment T₈ and T₀. In the last year 2018-2019, the maximum value on this parameter was found from T₈ (3.89) and T₄ (3.69) and then gone downward as followed T₅ (3.36) >T₇ (3.22) >T₃ (3.09) >T₂ (2.91) >T₆ (2.84) >T₁ (2.73) over T₀ (2.16).

The attempts to evaluate the combined effects of variety and various soil amendments on effective tiller/plant had insignificant. But along the experimental period, as V_3 and T_8 individually presented the greater value thus their interaction (V_3T_8) produced the maximum effective tiller/plant (4.20) in 2018-2019 and oppositely minimum (1.87) effective tiller/plant noticed from V_2T_0 in 2016-2017 (Table 26).

	Plar	nt height (cm)	То	otal plant/	m ²	То	tal tiller/pl	ant	Effe	ctive tiller	/plant
Variety	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
V	80.24	81.68	84.42	177.00	185.11	189.19	2.86	2.95	3.16	2.71	2.81	2.98
V_1	$\pm 1.07c$	± 1.19	± 1.23c	$\pm 3.01b$	± 3.20a	$\pm 3.67b$	$\pm 0.07c$	$\pm 0.08c$	$\pm 0.08c$	$\pm 0.08b$	$\pm 0.08c$	$\pm 0.08b$
V	81.62	82.48	85.64	173.00	179.93	181.37	2.98	3.08	3.27	2.80	2.92	3.07
V_2	± 1.09b	± 1.23	± 1.33b	$\pm 2.78b$	$\pm 2.80b$	± 3.06c	$\pm 0.10b$	$\pm 0.10b$	$\pm 0.11b$	$\pm 0.10b$	$\pm 0.10b$	$\pm 0.11b$
V	83.16	84.37	86.88	182.22	187.67	193.56	3.11	3.19	3.52	2.93	3.06	3.25
V ₃	± 1.14a	± 1.18	± 1.36a	± 2.67a	± 3.07a	± 3.86a	± 0.09a	± 0.10a	± 0.11a	± 0.10a	± 0.11a	± 0.12a
LS	**	NS	**	**	**	**	**	**	**	**	**	**
CV (%)	1.77	5.42	1.50	5.21	4.20	3.27	5.03	6.02	5.83	6.37	6.56	5.92

Table 24: Effect of variety on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI-28, V_2 = BARI-29, V_3 = BARI-30, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

	Plant height (cm)			Total plant/m ²			Total tiller/plant			Effective tiller/plant		
Treatment	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
T_0	75.76	74.63	78.45	158.67	163.89	159.00	2.24	2.32	2.33	2.07	2.10	2.16
10	± 1.05g	$\pm 1.05 f$	$\pm 1.46g$	± 3.11g	± 2.59f	± 2.29f	$\pm 0.07g$	$\pm 0.04g$	$\pm 0.07 g$	$\pm 0.08e$	$\pm 0.06 f$	$\pm 0.07 f$
T_1	77.61	78.77	81.43	164.89	170.22	173.44	2.62	2.73	2.98	2.47	2.58	2.73
11	± 1.31f	± 1.38ef	± 1.54f	± 3.17fg	± 2.17ef	± 1.69e	$\pm 0.05 f$	$\pm 0.06 f$	$\pm 0.07 f$	$\pm 0.05d$	$\pm 0.05e$	$\pm 0.05e$
T_2	79.35	80.57	83.44	171.44	177.00	179.00	2.82	2.87	3.11	2.67	2.73	2.91
12	± 1.28e	± 1.36de	± 1.71e	± 2.59ef	± 3.09de	± 2.35e	$\pm 0.06e$	$\pm 0.06 \text{ef}$	$\pm 0.06 \text{ef}$	$\pm 0.07c$	± 0.06de	$\pm 0.05e$
T_3	80.89	82.06	85.63	175.44	182.22	187.22	2.89	3.02	3.29	2.71	2.87	3.09
13	± 1.46d	\pm 1.28cde	$\pm 1.74d$	± 3.46de	± 2.33 cd	± 2.20d	$\pm 0.06e$	± 0.05 de	$\pm 0.07 de$	$\pm 0.05c$	± 0.06cd	± 0.06d
T_4	87.19	88.68	91.29	191.56	202.89	209.89	3.51	3.62	3.84	3.40	3.53	3.69
14	± 1.48b	± 1.41ab	$\pm 1.92b$	$\pm 2.86ab$	$\pm 2.53a$	± 3.53b	$\pm 0.06b$	$\pm 0.08b$	$\pm 0.09b$	$\pm 0.08a$	$\pm 0.08a$	$\pm 0.09b$
T_5	84.08	85.63	88.38	185.00	193.78	199.78	3.24	3.33	3.62	3.07	3.18	3.36
15	± 1.73c	± 1.57bc	± 1.94c	± 3.44bc	$\pm 2.00b$	± 2.56c	$\pm 0.09c$	$\pm 0.08c$	$\pm 0.10c$	$\pm 0.08b$	$\pm 0.09b$	$\pm 0.08c$
T_6	78.46	79.93	82.89	169.44	173.67	177.67	2.76	2.84	3.13	2.56	2.71	2.84
16	± 1.13ef	± 1.61de	± 1.50e	± 2.74ef	± 2.32e	± 1.76e	± 0.04ef	$\pm 0.07 ef$	$\pm 0.08 \text{ef}$	± 0.06 cd	± 0.05 de	$\pm 0.07e$
т	82.08	83.92	85.50	181.11	186.67	190.22	3.09	3.18	3.40	2.91	3.05	3.22
T_7	± 1.14d	$\pm 1.29 bcd$	± 1.78d	± 2.69 cd	± 2.26bc	$\pm 2.66d$	$\pm 0.07 d$	$\pm 0.07 cd$	$\pm 0.08 d$	$\pm 0.07 b$	$\pm 0.07 bc$	$\pm \ 0.07 cd$
m	89.62	91.38	93.81	199.11	207.78	216.11	3.67	3.80	4.13	3.49	3.64	3.89
T_8	± 1.34a	± 1.32a	$\pm 2.09a$	$\pm 2.65a$	$\pm 3.25a$	$\pm 4.17a$	$\pm 0.08a$	$\pm 0.09a$	$\pm 0.11a$	$\pm 0.09a$	$\pm 0.10a$	± 0.11a
LS	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	1.77	5.42	1.50	5.21	4.20	3.27	5.03	6.02	5.83	6.37	6.56	5.92

Table 25: Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

Voriety 0		Plant height (cm)			Total plant/m ²			Total tiller/plant			Effective tiller/plant		
	Variety & Treatment		2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
Treat	iment	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
	T_0	75.12	74.21	77.00	158.00	161.33	159.67	2.27	2.27	2.33	2.13	2.07	2.20
		± 1.33	± 2.08	± 2.93	± 7.23	± 6.64	± 4.33	± 0.07	± 0.07	± 0.07	± 0.18	± 0.07	± 0.12
	T_1	76.18	77.88	80.40	164.00	171.33	175.00	2.60	2.67	2.87	2.40	2.53	2.67
		± 2.46	± 2.64	± 2.97	± 7.81	± 4.47	± 3.22	± 0.12	± 0.13	± 0.07	± 0.12	± 0.07	± 0.07
	T_2	77.97	79.21	82.90	171.33	177.33	179.33	2.73	2.80	3.00	2.53	2.67	2.87
		± 1.55	± 2.37	± 3.39	± 5.55	± 8.41	± 6.01	± 0.07	±0.12	± 0.12	± 0.07	± 0.13	± 0.07
	T_3	79.40	80.69	84.70	174.33	183.00	188.33	2.80	2.93	3.20	2.73	2.80	3.00
	13	± 2.28	± 2.48	± 2.93	± 6.12	± 5.29	± 1.76	± 0.12	± 0.07	± 0.12	± 0.13	± 0.12	± 0.12
\mathbf{V}_1	T_4	85.94	87.27	90.00	190.67	204.67	212.33	3.33	3.47	3.60	3.20	3.33	3.47
v ₁	14	± 2.91	± 2.60	± 3.33	± 6.39	± 2.60	± 3.71	± 0.07	± 0.13	± 0.12	± 0.12	± 0.07	± 0.07
	T_5	82.13	84.45	86.40	186.00	194.00	200.00	3.00	3.13	3.33	2.87	2.93	3.13
		± 3.96	± 3.77	± 3.48	± 6.43	± 4.51	± 1.53	± 0.12	± 0.07	± 0.13	± 0.07	± 0.07	± 0.07
	T_6	76.83	78.61	82.00	168.33	174.33	177.67	2.67	2.73	3.07	2.47	2.67	2.80
		± 2.01	± 3.09	± 2.97	± 4.91	± 4.41	± 1.20	± 0.07	± 0.07	± 0.13	± 0.07	± 0.07	± 0.12
	T_7	80.96	82.78	83.78	180.67	189.33	190.33	2.93	3.00	3.20	2.80	2.87	3.07
		± 2.85	± 3.12	± 2.50	± 5.78	± 1.76	± 4.33	± 0.13	± 0.12	± 0.12	± 0.12	± 0.13	± 0.07
	T ₈	87.62	89.99	92.60	199.67	208.67	220.00	3.40	3.53	3.80	3.27	3.40	3.60
		± 2.73	± 2.39	± 3.12	± 4.41	± 6.64	± 4.04	± 0.12	± 0.07	± 0.12	± 0.07	± 0.12	± 0.12
	T_0	75.87	74.55	78.60	156.33	160.33	153.00	2.07	2.20	2.13	1.87	1.93	2.00
	10	± 2.42	± 2.31	± 2.70	± 5.90	± 2.03	± 2.52	± 0.07	± 0.12	± 0.07	± 0.07	± 0.07	± 0.12
	T_1	77.71	78.29	81.69	160.33	166.00	169.00	2.67	2.73	2.93	2.53	2.60	2.73
	11	± 2.63	± 2.83	± 2.73	± 5.24	± 3.79	± 2.31	± 0.07	± 0.07	± 0.13	± 0.07	± 0.12	± 0.07
	T_2	79.03	80.06	83.29	167.00	175.00	176.67	2.80	2.87	3.13	2.67	2.73	2.93
		± 2.84	± 2.18	± 3.17	± 4.36	± 5.13	± 3.53	± 0.12	± 0.07	± 0.07	± 0.13	± 0.07	± 0.07
	T_3	80.38	81.23	85.48	172.33	178.33	182.00	2.87	3.00	3.27	2.73	2.87	3.13
	13	± 2.69	± 2.05	± 3.948	± 8.95	± 2.73	± 3.46	± 0.07	± 0.12	± 0.13	± 0.07	± 0.07	± 0.07
V_2	T_4	86.94	88.47	91.87	188.00	196.67	199.00	3.53	3.67	3.80	3.40	3.53	3.67
• 2		± 2.51	± 3.13	± 4.01	± 5.13	± 3.48	± 4.04	± 0.07	± 0.07	± 0.12	± 0.12	± 0.07	± 0.18
	T ₅	84.70	85.63	88.62	179.33	191.00	193.00	3.33	3.40	3.60	3.00	3.20	3.33
		± 2.93	± 2.79	± 3.84	± 7.54	± 3.22	± 3.51	± 0.13	± 0.12	± 0.12	± 0.12	± 0.12	± 0.07

Table 26: Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat.

Contd....

Variety &		Plant height (cm)			Total plant/m ²			Total tiller/plant			Effective tiller/plant		
Treatm		2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
Treatin	em	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
	T_6	78.50	79.16	83.12	164.67	170.33	172.67	2.73	2.80	3.00	2.60	2.67	2.73
	1 ₆	± 2.28	± 3.77	± 3.11	± 4.06	± 4.63	± 2.73	± 0.07	± 0.12	± 0.12	± 0.12	± 0.07	± 0.07
	T_7	82.14	83.87	84.42	176.00	180.33	183.67	3.07	3.20	3.40	2.87	3.07	3.20
	17	± 1.85	± 2.29	± 3.81	± 4.36	± 4.06	± 3.33	± 0.07	± 0.12	± 0.12	± 0.13	± 0.07	± 0.12
	T_8	89.29	91.05	93.63	193.00	201.33	203.33	3.73	3.87	4.20	3.53	3.67	3.87
	18	± 2.42	± 2.77	± 4.05	± 4.04	± 5.24	± 4.33	± 0.07	± 0.13	± 0.12	± 0.13	± 0.18	± 0.18
	T_0	76.30	75.13	79.76	161.67	168.00	164.33	2.40	2.27	2.53	2.20	2.20	2.27
	10	± 2.27	± 1.81	± 2.76	± 4.63	± 4.16	± 2.33	± 0.12	± 0.07	± 0.07	± 0.12	± 0.12	± 0.07
	T_1	78.95	80.15	82.20	170.33	173.33	176.33	2.60	2.80	3.13	2.47	2.60	2.80
		± 2.36	± 2.54	± 3.35	± 2.60	± 2.40	± 1.86	± 0.12	± 0.12	± 0.13	± 0.07	± 0.12	± 0.12
	T_2	81.04	82.44	84.12	176.00	178.67	181.00	2.93	2.93	3.20	2.80	2.80	2.93
		± 2.58	± 2.98	± 3.64	± 3.22	± 3.71	± 3.61	± 0.13	± 0.13	± 0.12	± 0.12	± 0.12	± 0.13
	T_3	82.90	84.26	86.70	179.67	185.33	191.33	3.00	3.13	3.40	2.67	2.93	3.13
		± 3.11	± 2.38	± 3.35	± 3.38	± 4.10	± 4.49	± 0.12	± 0.07	± 0.12	± 0.07	± 0.13	± 0.13
V ₃	T_4	88.67	90.29	92.00	196.00	207.33	218.33	3.67	3.73	4.13	3.60	3.73	3.93
v ₃	14	± 3.08	± 2.20	± 3.98	± 3.79	± 5.21	± 4.63	± 0.07	± 0.18	± 0.07	±0.12	± 0.18	± 0.07
	т	85.42	86.80	90.13	189.67	196.33	206.33	3.40	3.47	3.93	3.33	3.40	3.60
	T_5	± 2.96	± 2.49	± 3.85	± 4.10	± 3.18	± 4.41	± 0.12	± 0.13	± 0.07	± 0.07	± 0.12	± 0.12
	T_6	80.05	82.02	83.55	175.33	176.33	182.67	2.87	3.00	3.33	2.60	2.80	3.00
		± 1.87	± 1.95	± 2.80	± 4.49	± 3.76	± 1.86	± 0.07	± 0.12	± 0.13	± 0.12	± 0.12	± 0.12
	T_7	83.13	85.12	88.29	186.67	190.33	196.67	3.27	3.33	3.60	3.07	3.20	3.40
		± 1.68	± 1.94	± 3.36	± 2.33	± 3.38	± 3.53	± 0.07	± 0.07	± 0.12	± 0.07	± 0.12	± 0.12
	т	91.95	93.09	95.20	204.67	213.33	225.00	3.87	4.00	4.40	3.67	3.87	4.20
	T_8	± 1.85	± 2.25	± 4.75	± 3.76	± 4.33	± 6.66	± 0.07	± 0.12	± 0.12	± 0.18	± 0.18	± 0.12
LS		NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
CV (%	6)	1.77	5.42	1.50	5.21	4.20	3.27	5.03	6.02	5.83	6.37	6.56	5.92

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, CV = Co-efficient of variation.

4.4.5 Awn length

Variety responded significantly on awn length of wheat and almost alike development was reported by the variety in every year under the research period. The gradual increase in awn length of variety was noticed as descending order by $V_3 > V_1 > V_2$. Hence, V_3 (6.72 cm) and V_1 (6.61 cm) demonstrated the higher value with statistical identity of awn length compared to V_2 (6.41 cm) in 2016-2017. Such developments were continued in the second year (2017-2018) and third year (2018-2019) of the study and thus the maximum (6.86 cm) and the minimum (6.51 cm) length of awn recorded from V_3 and V_2 in 2018-2019 (Table 27).

A significant difference was found in the awn length of wheat among the various treatments in this study (Table 28). In 2016-2017, treatment T₈ performed well (7.15 cm) to produce awn length, while T₀ resulted in less (6.03 cm). Rest of the treatments were influenced the awn length by descending order as T₄ (6.95 cm) >T₅ (6.82 cm) >T₇ (6.66 cm) >T₃ (6.59 cm) >T₂ (6.40 cm) >T₆ (6.36 cm) >T₁ (6.03 cm), though T₈, T₄ and T₅ had the statistical similarity. As like previous year, T₈ and T₀ treatment showed the highest (7.29 cm) and lowest (5.65 cm) awn length in 2017-2018. But statistical similarity was found among T₈ and T₄, T₄ and T₅ treatments. Compared with the previous year, the awn length was increased in 2018-2019. Further, the maximum (7.38 cm) and minimum (5.36 cm) awn length was noticed from T₈ and T₀, respectively in this period (Table 28).

The insignificant interaction between variety and soil amendments was observed for awn length mostly with gradual increase. The numerical values of this parameter considering the maximum (7.50 cm) and the minimum (5.23 cm) length were obtained from V_3T_8 and V_2T_0 in 2018-2019 (Table 29).

4.4.6 Spike length

There were significant differences among the varieties in respect of spike length in the whole research period (Table 27). In 2017-2018, variety V₃ showed the largest spike length (7.89 cm) which had similar statistical equality with V₁ (7.85 cm), the least spike length was observed in V₂ (7.63 cm). Similar results were noticed by the varieties in 2017-2018, at this time V₃ also produced the higher spike length (8.00 cm) followed by V₁ (7.98 cm)

and V_2 (7.85 cm). In the last year of study, each variety showed a slight increase in spike length compared to previous year. But statistical identity was found between V_3 (8.10 cm) and V_1 (7.98 cm).

There were significant differences among the sources soil amendment in terms of spike length in every year of the study (Table 28). Chemical fertilizer (T_8) was resulted the largest spike length than other soil amendments. Among the organic soil amendments, poultry manure produced larger spike length. During 2016-2017, T_8 demonstrated the highest (8.60 cm) spike length statistically at par with T_4 (8.12 cm), while the smallest spike length reported from T_0 (7.31 cm). In contrast, in 2017-2018, similar effects were recorded from the treatments where the largest (8.41 cm) and the smallest (7.12 cm) spike length were also recorded from T_8 and T_0 , correspondingly. There the application of poultry manure (8.29 cm) was comparable with chemical fertilizer (8.41 cm) in respect of this parameter. The spike length for all the treatments involving soil amendment was marginally higher in 2018-2019 than previous years. At this point, poultry manure (T_4) was revealed the second most ranked next to chemical fertilizer (T_8) having the values (8.43 cm) and (8.60 cm), respectively (Table 28).

There was no significant variation among the treatment combinations on spike length of wheat in the complete study period. The lengths of spike were ranged as the lowest and the highest from 6.89 to 8.73 cm in research time and that were presented by V_2T_0 and V_3T_8 in 2018-2019 (Table 29).

4.4.7 Spikelet/spike

Spikelet/spike was significantly affected by the selected varieties of wheat under this experiment period. The maximum value (16.34) of this parameter was observed with V_3 and lowers one by V_2 (15.75). However, performance of V_3 was closely followed by V_1 (16.04) on this parameter in 2016-2017. Statistically significant difference was found among the varieties in 2017-2018. Where V_3 was produced higher number (16.52) spikelet/spike followed by V_1 (16.17), but V_1 was statistically at par with V_2 (16.03). Similar performance of wheat variety was noticed in this regard during 2018-2019.

As also previous, V_3 was produced the highest (16.67) spikelet/spike, while V_1 and V_2 were presented the next values 16.36 and 16.20 with no significant differences between two varieties (Table 27).

Application of different soil amendments influenced spikelet/spike significantly in each year of the study. From the recorded result in the table 28, it is noticed that spikelet/spike varied significantly from 13.80 (T_0) to 17.97 (T_8) and this was also the maximum and the minimum value for this parameter under the study time. But among the treatments, the same statistical interrelation was observed as with (i) T_8 and T_4 (ii) T_5 , T_7 and T_3 (iii) T_6 , T_2 and T_1 all over the experimental period.

There was no significant variation among the treatment combinations of variety and soil amendments concerning the spikelet/spike at harvesting from the first year to the last year of the study. Nevertheless, the upper limit spikelet/spike (18.20) was recorded from the treatment combination V_3T_{8} , while the lower limit of spikelet/spike (13.56) was obtained from the treatment combination V_2T_0 (Table 29).

4.4.8 Fertile spikelet/spike

In the present experiment, varietal effect on fertile spikelet/spike showed significant difference in each year. The fertile spikelet/spike under variety V_3 was higher compared to V_1 and V_2 in the whole research period. In 2016-2017, the better fertile spikelet/spike (15.78) was observed with V_3 followed by V_1 (15.48) and V_2 (15.19). As compared to the first year, a little bit improvement of fertile spikelet/spike was recorded in the second and the third year of investigation. Thus, among the varieties V_3 produced the maximum (16.06) fertile spikelet/spike during 2018-2019, while the minimum value (15.19) was found from V_2 in 2016-2017 (Table 27).

Data shows in Table 28 indicated that fertile spikelet/spike of wheat was positively enhanced by various soil amendments at all round the research. Further, data demonstrated that incorporation of chemical fertilizer (T_8) and poultry manure (T_4) was at par to each other with superior significance over the rest of the treatments. However, in 2016-2017, the maximum (16.89) and the minimum (14.00) result yielded on this parameter from T_8 and T_0 . Further evaluation showed that probiotics acted as the third most significant transporter of the fertile spikelet/spike. Related records were also reported for this parameter during 2017-2018 and 2018-2019. Although the parallel trend of fertile spikelet/spike from respective treatments recorded, but 17.24 were the biggest figure and 13.22 was the lowest one obtained from T_8 and T_0 during 2018-2019 (Table 28).

Data shows in Table 29 indicated that there was no significant difference among variety and various treatment combinations in terms of their effect on fertile spikelet/spike over the study period. Overall, 4.20 was the high-grade result on the contrary, 1.87 was the low-grade result for fertile spikelet/spike under the experimental treatment combinations V_3T_8 and V_2T_0 in 2018-2019 and 2017-2018, respectively.

	Aw	vn length (c	cm)	Spi	ke length (cm)	S	pikelet/spik	te	Ferti	le spikelet/	′spike
Variety	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
V	6.61	6.69	6.76	7.75	7.853	7.98	16.04	16.17	16.36	15.48	15.66	15.77
\mathbf{V}_1	± 0.08a	$\pm 0.09b$	$\pm 0.11b$	± 0.08ab	± 0.09ab	± 0.10ab	± 0.18b	$\pm 0.22b$	$\pm 0.24b$	$\pm 0.18b$	± 0.19b	$\pm 0.25b$
V_2	6.41	6.53	6.62	7.63	7.76	7.85	15.75	16.03	16.20	15.19	15.31	15.47
• 2	$\pm 0.07b$	± 0.10c	$\pm 0.11c$	$\pm 0.09b$	$\pm 0.09b$	$\pm 0.10b$	$\pm 0.17c$	$\pm 0.23b$	$\pm 0.25b$	± 0.19c	$\pm 0.21c$	$\pm 0.23c$
V ₃	6.72	6.80	6.86	7.89	8.00	8.10	16.34	16.52	16.67	15.78	15.92	16.06
	± 0.08a	± 0.10a	± 0.12a	± 0.08a	± 0.09a	± 0.09a	± 0.18a	± 0.22a	± 0.24a	± 0.18a	± 0.18a	± 0.22a
LS	**	**	**	*	*	**	**	**	**	**	**	**
CV (%)	3.46	3.07	2.15	4.58	4.03	2.93	2.76	3.23	3.23	2.42	2.00	2.61

Table 27: Effect of variety on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. $V_1 = BARI-28$, $V_2 = BARI-29$, $V_3 = BARI-30$, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

	Av	wn length (cr	m)	Sp	ike length (c	m)	S	Spikelet/spik	e	Fert	ile spikelet/s	pike
Treatment	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019
T ₀	6.03 ± 0.07g	5.65 ± 0.09g	$\begin{array}{c} 5.36 \\ \pm \ 0.05 f \end{array}$	7.31 ± 0.16e	7.12 ± 0.10f	7.00 ± 0.05e	14.66 ± 0.21d	14.20 ± 0.20d	13.80 ± 0.15d	14.00 ± 0.20g	13.78 ± 0.17f	13.22 ± 0.19f
T_1	$\begin{array}{c} 6.26 \\ \pm \ 0.09 f \end{array}$	6.37 ± 0.08f	6.50 ± 0.06e	7.45 ± 0.11de	7.55 ± 0.11e	7.63 ± 0.08d	15.33 ± 0.14c	15.50 ± 0.26c	15.77 ± 0.17c	14.74 ± 0.22f	14.86 ± 0.13e	15.08 ± 0.22e
T_2	6.40 ± 0.09ef	6.54 ± 0.07ef	6.63 ± 0.07e	7.62 ± 0.12de	7.73 ± 0.11de	7.84 ± 0.08d	15.60 ± 0.12c	15.86 ± 0.20c	16.10 ± 0.14c	15.20 ± 0.12de	15.32 ± 0.10d	$\begin{array}{c} 15.51 \\ \pm \ 0.13 \text{d} \end{array}$
T_3	6.59 ± 0.09de	6.71 ± 0.07de	6.82 ± 0.08d	7.77 ± 0.09bcd	7.99 ± 0.12bcd	8.18 ± 0.10c	$\begin{array}{c} 16.10 \\ \pm \ 0.14b \end{array}$	$\begin{array}{c} 16.40 \\ \pm \ 0.23b \end{array}$	16.63 ± 0.16b	15.50 ± 0.14cd	15.72 ± 0.12c	15.93 ± 0.08c
T_4	6.95 ± 0.09ab	7.17 ± 0.05ab	7.27 ± 0.07a	8.12 ± 0.12ab	8.29 ± 0.11ab	8.43 ± 0.08ab	17.16 ± 0.13a	17.59 ± 0.17a	17.81 ± 0.16a	16.60 ± 0.16a	16.78 ± 0.09a	17.00 ± 0.13a
T_5	6.82 ± 0.07bc	6.99 ± 0.09bc	7.12 ± 0.06b	7.91 ± 0.10bc	8.12 ± 0.07abc	8.25 ± 0.06bc	16.43 ± 0.14b	16.68 ± 0.16b	$\begin{array}{c} 16.88 \\ \pm \ 0.21b \end{array}$	15.90 ± 0.16b	16.16 ± 0.15b	$\begin{array}{c} 16.40 \\ \pm \ 0.17b \end{array}$
T_6	6.36 ± 0.10f	6.53 ± 0.06ef	6.62 ± 0.07e	7.54 ± 0.10cde	7.66 ± 0.08de	7.77 ± 0.08d	15.51 ± 0.15c	15.78 ± 0.20c	16.01 ± 0.17c	14.90 ± 0.18ef	15.10 ± 0.17de	15.41 ± 0.15de
T_7	6.66 ± 0.06cd	6.83 ± 0.07cd	6.97 ± 0.06c	7.79 ± 0.12bcd	7.95 ± 0.09cd	8.08 ± 0.06c	16.26 ± 0.19b	16.46 ± 0.11b	16.70 ± 0.19b	15.62 ± 0.15bc	15.89 ± 0.13bc	16.12 ± 0.16bc
T_8	7.15 ± 0.06a	7.29 ± 0.06a	7.38 ± 0.07a	8.27 ± 0.08a	8.41 ± 0.09a	8.60 ± 0.10a	17.36 ± 0.17a	17.68 ± 0.16a	17.97 ± 0.17a	16.89 ± 0.13a	17.06 ± 0.10a	17.24 ± 0.18a
LS	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	3.46	3.07	2.15	4.58	4.03	2.93	2.76	3.23	3.23	2.42	2.00	2.61

Table 28: Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

V		Av	wn length (c	cm)	Spi	ike length (cm)	S	Spikelet/spik	æ	Ferti	ile spikelet/s	spike
	ariety & eatment	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
11	eatment	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
	T	6.04	5.69	5.39	7.29	7.13	7.00	15.00	14.27	13.87	14.20	14.00	13.17
	T_0	± 0.14	± 0.05	± 0.07	± 0.18	± 0.13	± 0.07	± 0.42	± 0.50	± 0.24	± 0.35	± 0.21	± 0.46
	т	6.28	6.38	6.50	7.42	7.51	7.63	15.30	15.47	15.70	14.67	14.77	14.93
	T_1	± 0.19	± 0.16	± 0.09	± 0.22	± 0.19	± 0.11	± 0.21	± 0.39	± 0.36	± 0.33	± 0.09	± 0.41
	т	6.47	6.59	6.66	7.59	7.71	7.85	15.53	15.83	16.07	15.13	15.27	15.47
	T_2	± 0.20	± 0.14	± 0.14	± 0.17	± 0.11	± 0.10	± 0.18	± 0.27	± 0.12	± 0.07	± 0.13	± 0.27
	т	6.63	6.71	6.81	7.76	8.00	8.20	16.00	16.30	16.53	15.40	15.70	15.87
	T_3	± 0.15	± 0.12	± 0.13	± 0.11	± 0.12	± 0.21	± 0.12	± 0.27	± 0.24	± 0.27	± 0.17	± 0.15
V	т	6.96	7.15	7.26	8.15	8.29	8.45	17.20	17.53	17.83	16.60	16.80	17.03
\mathbf{V}_1	T_4	± 0.15	± 0.07	± 0.09	± 0.13	± 0.17	± 0.23	± 0.23	± 0.48	± 0.26	± 0.27	± 0.06	± 0.20
	т	6.85	7.01	7.15	7.90	8.12	8.25	16.33	16.50	16.77	15.93	16.33	16.60
	T_5	± 0.09	± 0.08	± 0.09	± 0.11	± 0.12	± 0.13	± 0.35	± 0.29	± 0.38	± 0.24	± 0.18	± 0.23
	т	6.41	6.52	6.63	7.49	7.56	7.75	15.47	15.67	15.90	14.83	15.03	15.30
	T_6	± 0.19	± 0.14	± 0.13	± 0.27	± 0.12	± 0.10	± 0.26	± 0.24	± 0.21	± 0.22	± 0.19	± 0.25
	T_7	6.67	6.87	7.03	7.80	7.95	8.09	16.13	16.23	16.57	15.63	15.97	16.30
	17	± 0.08	± 0.09	± 0.09	± 0.18	± 0.18	± 0.04	± 0.49	± 0.18	± 0.43	± 0.19	± 0.15	± 0.15
	T_8	7.16	7.29	7.39	8.32	8.40	8.59	17.40	17.70	17.97	16.93	17.07	17.27
	18	± 0.11	± 0.09	± 0.11	± 0.06	± 0.23	± 0.26	± 0.27	± 0.32	± 0.29	± 0.35	± 0.12	± 0.37
	T_0	5.90	5.49	5.23	7.20	7.02	6.89	14.30	13.97	13.53	13.47	13.20	12.97
	10	± 0.12	± 0.23	± 0.08	± 0.37	± 0.19	± 0.06	± 0.44	± 0.26	± 0.29	± 0.29	± 0.15	± 0.09
	T_1	6.11	6.25	6.39	7.35	7.43	7.48	15.00	15.20	15.47	14.50	14.57	14.73
	11	± 0.06	± 0.14	± 0.09	± 0.12	± 0.19	± 0.18	± 0.12	± 0.42	± 0.29	± 0.27	± 0.22	± 0.43
	T_2	6.19	6.42	6.50	7.53	7.65	7.70	15.33	15.53	15.87	15.10	15.10	15.30
	12	±0.09	± 0.10	± 0.10	± 0.28	± 0.28	± 0.22	± 0.12	± 0.35	± 0.18	± 0.06	± 0.15	± 0.21
	T_3	6.35	6.55	6.72	7.67	7.86	8.07	15.83	16.10	16.40	15.30	15.53	15.80
	13	± 0.09	± 0.13	± 0.13	± 0.24	± 0.24	± 0.20	± 0.15	± 0.59	± 0.31	± 0.15	± 0.24	± 0.12
V_2	T_4	6.80	7.05	7.13	7.95	8.17	8.31	16.80	17.43	17.60	16.37	16.53	16.73
• 2	14	± 0.23	± 0.10	± 0.140	± 0.30	± 0.24	± 0.06	± 0.12	± 0.23	± 0.38	± 0.41	± 0.12	± 0.18
	T_5	6.66	6.85	7.00	7.78	8.03	8.15	16.23	16.67	16.80	15.50	15.67	15.83
	15	± 0.09	± 0.26	± 0.12	± 0.22	± 0.09	± 0.09	± 0.15	± 0.24	± 0.32	± 0.21	± 0.18	± 0.09
	T_6	6.13	6.41	6.52	7.45	7.58	7.63	15.17	15.50	15.73	14.57	14.80	15.13
	16	± 0.08	± 0.05	± 0.14	± 0.07	± 0.16	± 0.15	± 0.15	± 0.29	± 0.29	± 0.22	± 0.35	± 0.26
	T_7	6.54	6.62	6.79	7.61	7.79	7.93	16.07	16.43	16.67	15.23	15.47	15.67
	1 /	± 0.10	± 0.12	± 0.07	± 0.26	± 0.13	± 0.13	± 0.23	± 0.15	± 0.33	± 0.15	± 0.12	± 0.23
	T_8	7.00	7.16	7.25	8.12	8.28	8.47	17.03	17.40	17.73	16.70	16.90	17.10
	18	± 0.07	± 0.11	± 0.08	± 0.19	± 0.08	± 0.13	± 0.24	± 0.31	± 0.41	± 0.17	± 0.21	± 0.27

Table 29: Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat.

Contd	,

Vo	riety &	Av	wn length (c	cm)	Spi	ike length (em)	S	pikelet/spik	te	Ferti	le spikelet/s	spike
	•	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
110	eatment	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
	T_0	6.15	5.78	5.47	7.45	7.20	7.13	14.67	14.37	14.00	14.33	14.13	13.53
	10	± 0.09	± 0.13	± 0.09	± 0.34	± 0.26	± 0.06	± 0.24	± 0.32	± 0.23	± 0.19	± 0.19	± 0.34
	T_1	6.40	6.49	6.60	7.58	7.72	7.79	15.70	15.83	16.13	15.07	15.23	15.57
	11	± 0.19	± 0.10	± 0.13	± 0.24	± 0.23	± 0.11	± 0.21	± 0.60	± 0.19	± 0.58	± 0.20	± 0.23
	T_2	6.52	6.59	6.75	7.73	7.84	7.97	15.93	16.20	16.37	15.37	15.60	15.77
	12	± 0.11	± 0.12	± 0.14	± 0.21	± 0.22	± 0.10	± 0.18	± 0.44	± 0.35	± 0.38	± 0.12	± 0.17
	т	6.79	6.86	6.94	7.88	8.11	8.27	16.47	16.80	16.97	15.80	15.93	16.13
	T ₃	± 0.16	± 0.03	± 0.16	± 0.14	± 0.27	± 0.15	± 0.29	± 0.31	± 0.23	± 0.25	± 0.18	± 0.09
V	T_4	7.10	7.31	7.43	8.25	8.40	8.52	17.47	17.80	18.00	16.83	17.00	17.23
V_3	14	± 0.06	± 0.05	± 0.09	± 0.16	± 0.20	± 0.08	± 0.18	± 0.17	± 0.27	± 0.15	± 0.12	± 0.27
	T_5	6.94	7.13	7.22	8.05	8.20	8.33	16.73	16.87	17.07	16.27	16.47	16.77
	15	± 0.13	± 0.08	± 0.05	± 0.21	± 0.14	± 0.08	± 0.18	± 0.35	± 0.48	± 0.24	± 0.18	± 0.15
	т	6.53	6.66	6.72	7.67	7.85	7.94	15.90	16.17	16.40	15.30	15.47	15.80
	T_6	± 0.18	± 0.09	± 0.13	± 0.18	± 0.10	± 0.11	± 0.21	± 0.44	± 0.32	± 0.38	± 0.30	± 0.12
	T_7	6.78	6.99	7.10	7.96	8.12	8.21	16.57	16.70	16.87	16.00	16.23	16.40
	17	± 0.13	± 0.08	± 0.06	± 0.22	± 0.12	± 0.06	± 0.23	± 0.17	± 0.35	± 0.25	± 0.15	± 0.29
	т	7.28	7.42	7.50	8.39	8.55	8.73	17.63	17.93	18.20	17.03	17.20	17.37
	T_8	± 0.04	± 0.07	± 0.14	± 0.13	± 0.13	± 0.10	± 0.35	± 0.20	± 0.21	± 0.15	± 0.21	± 0.38
	LS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
С	V (%)	3.46	3.07	2.15	4.58	4.03	2.93	2.76	3.23	3.23	2.42	2.00	2.61

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, CV = Co-efficient of variation.

4.4.9 Grains/spike

A significant variation in grains/spike of wheat was noticed due to varietal influences appearing in the study length. In 2016-2017, the grains/spike varied from 39.81 to 38.56 that were presented by V_3 and V_2 . In the next two years of experiment, V_3 also demonstrated higher efficiency than the other two varieties. Thus with the related performance of the varieties in 2018-2019, V_3 marked as the best producer of grains/spike (42.14), whereas in 2017-2018 V_2 reported as the lower producer with the value of 39.70 (Table 30).

In comparison to the control plot, development of grains/spike was significantly higher under different treatment combinations round the research. The maximum number of grains/spike (42.74) was noted in T_8 and the minimum value (36.88) obtained by T_0 in 2016-2017. There the treatments were ranked in order $T_8>T_4>T_5>T_7>T_3>T_6>T_2>T_1>T_0$ for grains/spike production. The grain in each spike was also varied with different soil amendments in 2017-2018. Where T_8 was produced the utmost grains/spike (44.41) which had statistical equality with T_4 (43.50) and the least value recorded from T_0 (35.04). In 2018-2019, the grains/spike was affected greatly by T_8 with the best value 45.73, oppositely 33.67 was the poorest value observed from T_0 (Table 31).

Treatment combination effects did not vary significantly for grains/spike except in 2018-2019. With statistical variation V_3T_8 combination was reported the highest number of grains/spike (47.60) followed by V_3T_4 (46.00) and the combination V_2T_0 (33.20) was noted as the lower one during 2018-2019 (Table 32).

4.4.10 Deformed grains/spike

In the three years of the study, nearly equal number of deformed grains/spike was produced by the selected wheat varieties with significant deviation. In 2016-2017, the higher (2.90) and the lower number (2.70) of deformed grains/spike recorded from V_1 and V_2 , respectively. In the next two years, alike results were found in respect of deformed grains/spike, where 2.90 was the maximum and 2.65 was the minimum value for recorded by V_1 and V_2 in 2018-2019 (Table 30). The number of deformed grains/spike varied significantly by the treatment factor all over the experimental duration. Most of the soil amendments exhibited a similar trend in production of deformed grains/spike. Through the control treatment (T_0) was denoted the highest score followed by T_8 , but the lowest was found in T_5 followed by T_7 in each year. The chronological higher values of the deformed grains/spike were recorded by T_0 (3.36, 3.49 and 3.72) and lower in T_5 (2.38, 2.29 and 2.27) collected in the sequential year of 2016-2017, 2017-2018 and 2018-2019, respectively (Table 31). The deformed grains/spike did not influence significantly by the interaction of variety coupled with soil amendments in the study period. But numerical differences were recorded from the treatment combinations where V_1T_0 (4.00) and V_2T_5 (2.17) in 2018-2019 (Table 32).

4.4.11 Grain weight/spike

The significant variation of grain weight/spike was recorded by the varietal influence in each sampling year. The grain weight/spike was ranged from 1.80-1.89 g in 2016-2017, where the maximum weight was observed by V_3 (1.89 g) which was at par with V_1 (1.86 g) and the minimum (1.80 g) was noted at V_2 . There was statistical equality among the varieties in the rest years of the experiment. Between the last two years, 2018-2019 had demonstrated a slight increase of grain weight/spike over 2017-2018. So, comparatively the higher (2.02 g) and the lower (1.85 g) grain weight/spike were recorded by V_3 and V_2 (Table 30).

Grain weight/spike at harvest varied significantly due to different soil amendments during the study period (Table 31). The highest value of grain weight/spike was found in T₈ (2.07 g) and the minimum result was recorded by T₀ (1.66 g) in 2016-2017. Where the next highest value was noticed from T₄ (1.98 g) and T₅ (1.93 g) had the statistical likeness. In the year 2017-2018, grain weight/spike exhibited a similar trend reported in 2016-2017. There the highest (2.15 g) grain weight/spike produced by T₈ and the second most in T₄ (2.10 g), but the lowest grain weight/spike (1.62 g) was noticed in T₀. Among the different soil amendments, T₈ (2.20 g) was performed superior besides T₄ (2.14 g) to other treatments in 2018-2019. Rest of the treatments were showed also significant difference ranked as descending by T₅ (2.06 g), T₇ (2.01 g), T₃ (1.99 g), T₂ (1.93 g), T₆ (1.89 g), T₁ (1.86 g), while the lowest one in T₀ (1.60 g). The objective to evaluate the combined effects of variety and various soil amendments on grain weight/spike was insignificant in 2016-2017 and in 2017-2018, while significant results were found in 2018-2019. In the experimental period 2018-2019, as V_3 and T_8 individually presented the greater value on this parameter thus their interaction (V_3T_8) produced the maximum grain weight/spike (2.25 g) and oppositely the minimum (1.57 g) grain weight/spike noticed from V_2T_0 in the same period. Furthermore, data in Table 32 clearly demonstrated the next greater grain weight/spike by V_1T_8 (2.20 g) with statistical akin by V_3T_4 (2.18 g).

4.4.12 1000-grain weight

The response of 1000-grain weight of selected wheat varieties in the research period is shown in the table 30. There this factor was significant in 2016-2017 and 2018-2018 other than 2017-2018. A bit increase in 1000-grain weight was reported in the consecutive years of the experiment. In 2016-2017, the maximum 1000-grain weight (51.86 g) was noticed in V_3 at the same time the minimum value (51.06 g) of 1000-grain weight obtained from V_3 . Though not significant differences were observed for 1000-grain weight during 2017-2018, but the trend of increase was parallel with the previous year. Here V_3 was superior (52.22 g), while V_2 had the minimum (51.72 g) weight in respect of 1000-seed of wheat. At the end year (2018-2019) of the experiment, significantly higher (52.52 g) and lower (51.89 g) 1000-grain weight was obtained from V_3 and V_2 , respectively.

The reaction of 1000-grain weight of wheat to different soil amendments showed significantly positive results in every year (Table 31). In 2016-2017, wheat plants that received chemical fertilizer (T₈) had the highest 1000-grain weight (53.28 g) followed by poultry manure (T₄), while plants in the control plot had the lowest 1000-grain weight (49.52 g). In 2017-2018, 1000-grain weight differed significantly under various treatments. In this case, 1000-grain weight was also higher (53.81 g) under T₈, but statistically interlinked with T₄ (53.51 g), T₅ (52.98 g), T₇ (52.51 g) and T₃ (52.34 g), while T₀ had the lower mean of 1000-grain weight (48.61 g). During 2018-2019, chemical fertilizer (T₈) was superior with a mean of 1000-grain weight (54.12 g), while the plants that did not receive any soil amendments (T₀) had the lowest mean (48.78 g).

Experimental results revealed that there were not significant differences among treatment combinations during 2016-2017 and 2017-2018, but there was significant variation in 2018-2019 with regard to 1000-grain weight (Table 32). Although the first and the second year were presented not statistical variation among the treatments, but numerically the leading 1000-grain weight (54.02 g) was recorded from V_3T_8 , whereas the inferior (48.55 g) was noticed in V_1T_0 . The value 54.45 g was remarked as the utmost, while the value 48.21 g denoted as the lowest exhibited by V_3T_8 and V_1T_0 , correspondingly during 2018-2019. At the same time, V_1T_8 was demonstrated the second higher 1000-grain weight (54.21 g) and this was statistically similar with V_3T_4 (54.14 g) and V_1T_4 (54.03 g).

	(Grains/spik	te	Defor	med grains	/spike	Grain	weight/spi	ke (g)	1000-	grain weig	ht (g)
Variety	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019	2017	2018	2019
V ₁	39.19	40.06	41.19	2.90	2.89	2.90	1.86	1.94	1.99	51.56	52.07	52.25
\mathbf{v}_1	$\pm 0.40b$	$\pm 0.52b$	$\pm 0.83b$	± 0.08a	± 0.09a	± 0.11a	± 0.03a	$\pm 0.03b$	$\pm 0.04b$	$\pm 0.32ab$	± 0.37	$\pm 0.38b$
V_2	38.56	39.70	40.62	2.71	2.66	2.65	1.80	1.85	1.89	51.06	51.72	51.89
• 2	± 0.37c	± 0.51b	± 0.81c	$\pm 0.07b$	± 0.08c	± 0.09c	$\pm 0.03b$	± 0.03c	$\pm 0.04c$	± 0.26b	± 0.46	± 0.36c
V_3	39.81	41.04	42.14	2.83	2.80	2.79	1.89	1.97	2.02	51.86	52.22	52.52
• 3	± 0.48a	± 0.63a	± 0.90a	± 0.07a	$\pm 0.08b$	$\pm 0.10b$	± 0.03a	± 0.03a	± 0.04a	± 0.28a	± 0.34	± 0.35a
LS	**	**	**	**	**	**	**	**	**	*	NS	**
CV (%)	1.52	2.55	1.09	6.12	5.85	4.80	3.69	2.90	1.48	1.90	3.12	1.23

Table 30: Effect of variety on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. $V_1 = BARI-28$, $V_2 = BARI-29$, $V_3 = BARI-30$, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

Treatment		Grains/spike	;	Defo	rmed grains/	/spike	Grair	n weight/spil	ke (g)	1000-grain weight (g)			
Treatment -	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	
T ₀	36.88 ± 0.30f	35.04 ± 0.30e	33.67 ± 0.87i	3.36 ± 0.07a	3.49 ± 0.07a	3.72 ± 0.09a	1.66 ± 0.03g	1.62 ± 0.02g	1.60 ± 0.04g	49.52 ± 0.44f	48.61 ± 0.51e	48.78 ± 0.41i	
T_1	$\begin{array}{c} 36.98 \\ \pm \ 0.22 f \end{array}$	$\begin{array}{c} 38.42 \\ \pm \ 0.23 d \end{array}$	$\begin{array}{c} 39.53 \\ \pm \ 0.88 h \end{array}$	$\begin{array}{c} 2.80 \\ \pm \ 0.07 c \end{array}$	2.69 ± 0.07c	2.63 ± 0.05d	$\begin{array}{c} 1.74 \\ \pm \ 0.02 f \end{array}$	$\begin{array}{c} 1.80 \\ \pm \ 0.03 f \end{array}$	$\begin{array}{c} 1.86 \\ \pm \ 0.04 f \end{array}$	$\begin{array}{c} 50.03 \\ \pm \ 0.37 f \end{array}$	$\begin{array}{c} 51.05 \\ \pm \ 0.30d \end{array}$	51.30 ± 0.31h	
T_2	38.11 ± 0.21e	$\begin{array}{c} 39.36 \\ \pm \ 0.26d \end{array}$	$\begin{array}{c} 40.67 \\ \pm \ 0.96f \end{array}$	2.67 ± 0.06cd	2.62 ± 0.06c	$\begin{array}{c} 2.56 \\ \pm \ 0.05 d \end{array}$	$\begin{array}{c} 1.80 \\ \pm \ 0.03 ef \end{array}$	$\begin{array}{c} 1.88 \\ \pm \ 0.03 de \end{array}$	1.93 ± 0.05e	$\begin{array}{c} 51.19 \\ \pm \ 0.33 de \end{array}$	51.84 ± 0.59bcd	$\begin{array}{c} 51.95 \\ \pm \ 0.33f \end{array}$	
T ₃	$\begin{array}{c} 38.76 \\ \pm \ 0.21 d \end{array}$	40.51 ± 0.39c	41.58 ± 1.02e	$\begin{array}{c} 2.52 \\ \pm \ 0.07 de \end{array}$	2.47 ± 0.06d	2.42 ± 0.06e	$\begin{array}{c} 1.83 \\ \pm \ 0.02 de \end{array}$	$\begin{array}{c} 1.91 \\ \pm \ 0.03 d \end{array}$	1.99 ± 0.04d	51.77 ± 0.19cd	52.34 ±0.34a-d	52.49 ± 0.34e	
T_4	42.11 ± 0.40b	43.50 ± 0.59a	44.80 ± 1.13b	3.14 ± 0.05b	3.07 ± 0.06b	2.94 ± 0.04c	$\begin{array}{c} 1.98 \\ \pm \ 0.02b \end{array}$	$\begin{array}{c} 2.10 \\ \pm \ 0.02b \end{array}$	2.14 ± 0.05b	52.94 ± 0.27ab	53.51 ± 0.47ab	53.90 ± 0.39b	
T ₅	40.17 ± 0.22c	41.59 ± 0.43b	43.14 ± 1.08c	2.38 ± 0.04e	2.29 ± 0.05e	$\begin{array}{c} 2.27 \\ \pm \ 0.05 f \end{array}$	1.93 ± 0.03bc	2.02 ± 0.02c	2.06 ± 0.05c	52.27 ± 0.21bc	52.98 ± 0.90abc	53.19 ± 0.38c	
T_6	37.20 ± 0.20f	38.71 ± 0.27d	40.23 ± 0.97g	2.73 ± 0.07c	2.66 ± 0.07c	2.57 ± 0.06d	1.77 ± 0.03ef	1.84 ± 0.03ef	1.89 ± 0.05f	50.48 ± 0.40ef	51.39 ± 0.31cd	51.60 ± 0.34g	
T_7	39.73 ± 0.25c	40.83 ± 0.42bc	42.50 ± 1.07d	2.47 ± 0.04e	2.38 ± 0.06de	2.34 ± 0.03ef	1.89 ± 0.02cd	1.99 ± 0.02c	$\begin{array}{c} 2.01 \\ \pm \ 0.05 d \end{array}$	51.97 ± 0.14bcd	52.51 ± 0.26a-d	52.64 ± 0.36d	
T ₈	42.74 ± 0.37a	44.41 ± 0.53a	45.73 ± 1.04a	3.23 ± 0.06a	3.37 ± 0.06a	3.53 ± 0.06b	2.07 ± 0.03a	2.15 ± 0.03a	2.20 ± 0.05a	53.28 ± 0.30a	53.81 ± 0.29a	54.12 ± 0.40a	
LS CV (%)	**	**	**	** 6.12	**	** 4.80	** 3.69	**	**	**	** 3.12	**	

Table 31: Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

V	Variety &		Grains/spik	e	Defo	rmed grains/	spike	Grai	n weight/spil	ke (g)	100	0-grain weig	nt (g)
Т	Treatment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	T ₀	37.53	34.93	33.80	3.47	3.67	4.00	1.65	1.61	1.59	49.20	48.55	48.21
	10	± 0.55	± 0.18	$\pm 1.56qr$	± 0.12	± 0.15	± 0.12	± 0.02	± 0.03	$\pm 0.07 op$	± 1.16	± 0.85	± 0.53p
	T_1	36.87	38.53	39.40	2.93	2.80	2.73	1.75	1.83	1.91	50.01	51.05	51.33
	1	± 0.35	± 0.41	± 1.480p	± 0.13	± 0.12	± 0.09	± 0.02	± 0.03	± 0.071	± 0.69	± 0.23	$\pm 0.62 mn$
	T_2	38.00	39.13	40.60	2.77	2.73	2.67	1.83	1.91	1.95	51.25	51.84	51.99
	12	± 0.23	± 0.44	± 2.00 k-n	± 0.12	± 0.09	± 0.09	± 0.02	± 0.02	± 0.09 jkl	± 0.72	± 1.56	± 0.65 hi
	T ₃	38.87	39.53	41.13	2.60	2.57	2.53	1.86	1.93	2.01	52.11	52.38	52.57
	13	± 0.24	± 0.48	± 1.78jk	± 0.12	± 0.12	± 0.07	± 0.02	± 0.03	± 0.08 hi	± 0.26	± 0.21	$\pm 0.66 f$
V_1	T_4	42.00	42.93	44.50	3.20	3.13	3.00	1.97	2.10	2.14	53.10	53.64	54.03
v ₁	14	± 0.21	± 0.94	± 2.44	± 0.12	± 0.13	± 0.06	± 0.03	± 0.03	± 0.10 cd	± 0.34	± 0.77	$\pm 0.75b$
	T ₅	40.03	41.87	43.40	2.43	2.40	2.33	1.92	2.04	2.09	52.38	53.11	53.36
	15	± 0.09	± 0.85	$\pm 1.88 ef$	± 0.09	± 0.06	± 0.09	± 0.04	± 0.04	$\pm 0.08 efg$	± 0.58	± 0.41	$\pm 0.73d$
	T_6	37.10	38.87	40.00	2.87	2.77	2.70	1.79	1.88	1.94	50.49	51.56	51.81
	16	± 0.21	± 0.18	± 1.96 mno	± 0.12	± 0.12	± 0.06	± 0.02	± 0.02	± 0.061	± 0.82	± 0.64	± 0.66ij
	T_7	39.57	40.90	42.70	2.50	2.43	2.40	1.90	2.00	2.05	52.16	52.57	52.75
	17	± 0.29	± 0.87	± 2.34 fgh	± 0.12	± 0.12	± 0.06	± 0.02	± 0.03	± 0.10 fgh	± 0.14	± 0.13	± 0.69ef
	T_8	42.73	43.80	45.20	3.37	3.50	3.70	2.08	2.16	2.20	53.34	53.92	54.21
	18	± 0.09	± 0.53	± 1.83c	± 0.09	± 0.12	± 0.06	± 0.04	± 0.01	$\pm 0.08b$	± 0.56	± 0.81	$\pm 0.78b$
	T ₀	36.37	34.27	33.20	3.27	3.37	3.47	1.59	1.58	1.57	49.18	48.25	48.11
	10	± 0.32	± 0.55	$\pm 1.82r$	± 0.12	± 0.09	± 0.09	± 0.03	± 0.03	$\pm 0.07 p$	± 0.60	± 1.10	± 0.53p
	T_1	36.53	38.13	39.00	2.67	2.53	2.50	1.68	1.71	1.75	49.63	50.94	51.16
	1	± 0.24	± 0.58	± 1.80p	± 0.12	± 0.09	± 0.06	± 0.03	± 0.03	$\pm 0.05n$	± 0.39	± 0.69	$\pm 0.60n$
	T_2	37.67	39.33	40.30	2.52	2.47	2.40	1.73	1.78	1.84	50.62	51.63	51.72
	12	± 0.18	± 0.68	± 1.51 k-n	± 0.06	± 0.12	± 0.06	± 0.06	± 0.03	$\pm 0.09 m$	± 0.26	± 0.80	± 0.64 jk
	T ₃	38.13	40.60	41.00	2.40	2.37	2.33	1.76	1.82	1.90	51.15	52.11	52.31
	13	± 0.24	± 0.70	± 1.78jkl	± 0.12	± 0.09	± 0.09	± 0.03	± 0.03	± 0.071	± 0.20	± 1.10	$\pm 0.66g$
V_2	T_4	41.00	42.47	43.90	3.07	2.97	2.90	1.95	2.06	2.10	52.58	53.05	53.54
• 2	14	± 0.64	± 0.66	± 2.15de	± 0.09	± 0.09	± 0.12	± 0.04	± 0.03	$\pm 0.10 def$	± 0.33	± 1.27	± 0.77 cd
	T ₅	39.73	40.67	42.13	2.33	2.20	2.17	1.90	1.95	1.99	51.89	52.58	52.75
	15	± 0.18	± 0.47	± 2.43 hi	± 0.09	± 0.06	± 0.09	± 0.03	± 0.02	± 0.06 ijk	± 0.17	± 0.90	$\pm 0.72 ef$
	T_6	36.70	38.67	39.90	2.60	2.50	2.43	1.70	1.75	1.77	50.13	51.23	51.43
	16	± 0.15	± 0.77	± 1.61no	± 0.12	± 0.06	± 0.12	± 0.06	± 0.02	$\pm 0.07 n$	± 0.79	± 0.71	± 0.65 lm
	T_7	39.43	40.03	41.70	2.40	2.30	2.27	1.86	1.92	1.94	51.54	52.20	52.28
	17	± 0.09	± 0.70	± 2.05 ij	± 0.06	± 0.12	± 0.03	± 0.01	± 0.05	$\pm 0.07 kl$	± 0.25	± 0.76	$\pm 0.70g$
	T_8	41.50	43.10	44.40	3.10	3.20	3.37	2.02	2.09	2.13	52.82	53.49	53.69
	18	± 0.25	± 0.25	$\pm 1.92d$	± 0.10	± 0.06	± 0.09	± 0.05	± 0.05	± 0.10 cde	± 0.41	± 0.47	± 0.78c

Table 32: Interaction effects of variety and soil amendments on yield and yield contributing characters of wheat.

Contd...

Va	ariety &		Grains/spike	e	Defo	rmed grains/	spike	Grai	n weight/spik	te (g)	100	0-grain weigl	ht (g)
Tre	eatment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	T ₀	36.73	35.93	34.00	3.33	3.43	3.70	1.73	1.65	1.63	50.19	49.02	50.01
	10	± 0.55	± 0.24	$\pm 1.77q$	± 0.12	± 0.09	± 0.06	± 0.05	± 0.02	± 0.060	± 0.53	± 0.99	± 0.560
	T ₁	37.53	38.60	40.20	2.80	2.73	2.67	1.78	1.85	1.93	50.45	51.16	51.42
	11	± 0.35	± 0.46	\pm 1.861-o	± 0.12	± 0.12	± 0.09	± 0.02	± 0.02	± 0.081	± 0.94	± 0.75	± 0.63 lm
	т	38.67	39.60	41.10	2.73	2.67	2.60	1.84	1.94	2.00	51.70	52.05	52.15
	T_2	± 0.41	± 0.31	$\pm 2.02 jk$	± 0.09	± 0.07	± 0.06	± 0.02	± 0.02	± 0.06	± 0.66	± 1.02	$\pm 0.68 gh$
	T ₃	39.27	41.40	42.60	2.57	2.47	2.40	1.88	1.98	2.05	52.04	52.53	52.59
	13	± 0.24	± 0.46	$\pm 2.34 gh$	± 0.12	± 0.07	± 0.12	± 0.03	± 0.02	± 0.08	± 0.16	± 0.35	$\pm 0.71 f$
V_3	т	43.33	45.10	46.00	3.17	3.10	2.93	2.00	2.14	2.18	53.14	53.85	54.14
v ₃	T_4	± 0.35	± 0.91	± 1.86b	± 0.09	± 0.06	± 0.07	± 0.07	± 0.05	$\pm 0.08 bc$	± 0.74	± 0.56	$\pm 0.77b$
	T ₅	40.73	42.23	43.90	2.37	2.27	2.30	1.95	2.07	2.09	52.53	53.25	53.47
	15	± 0.52	± 0.76	± 1.90de	± 0.07	± 0.09	± 0.06	± 0.07	± 0.03	$\pm 0.10 efg$	± 0.25	± 0.69	± 0.72d
	т	37.80	38.60	40.80	2.73	2.70	2.57	1.82	1.88	1.95	50.83	51.37	51.55
	T ₆	± 0.23	± 0.46	± 2.12 klm	± 0.09	± 0.12	± 0.03	± 0.05	± 0.04	± 0.07 jkl	± 0.68	± 0.43	± 0.67 kl
	т	40.20	41.57	43.10	2.50	2.40	2.37	1.91	2.03	2.05	52.21	52.75	52.90
	T ₇	± 0.70	± 0.50	$\pm 1.87 efg$	± 0.06	± 0.06	± 0.03	± 0.06	± 0.02	$\pm 0.06 gh$	± 0.12	± 0.38	± 0.72e
	т	44.00	46.33	47.60	3.23	3.40	3.53	2.13	2.21	2.26	53.66	54.02	54.45
	T ₈	± 0.12	± 0.41	± 1.79a	± 0.03	± 0.06	± 0.09	± 0.07	± 0.08	$\pm 0.10a$	± 0.65	± 0.16	$\pm 0.77a$
	LS	NS	NS	**	NS	NS	NS	NS	NS	**	NS	NS	**
С	CV (%)	1.52	2.55	1.09	6.12	5.85	4.80	3.69	2.90	1.48	1.90	3.12	1.23

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

4.4.13 Grain yield

The varietal response on grain yield was significant all over the study period. From the analyzed records, the highest mean grain yield of 3.38 t/ha was produced from the V₃, whereas the minimum grain yield of 3.17 t/ha was reported from the V₁ in the year 2016-2017 (Fig. 21). Like previous year similar trends were noticed in 2017-2018, where with slight increment varietal performance was ranked as ascending order V₂ (3.30 t/ha) >V₁ (3.37 t/ha) >V₃ (3.47 t/ha) (Fig. 22). At the end of three years study, the data showed an increased trend in the last year over the previous two years. The maximum grain yield (3.59 t/ha) was found in V₃ oppositely the least grain yield (3.40 t/ha) had in V₂ (Fig. 23).

Application of different soil amendments under this investigation was generated significant variation on the grain yield of wheat (Fig. 24, 25 and 26). The beneficial effects of different treatments on the grain yield was expressed in the range 4.08-2.67 t/ha in 2016-2017, here the maximum value was obtained from T₈ (4.08 t/ha) followed by T₄ (3.77 t/ha) and the least value was found in T₀ (2.67 t/ha). In 2017-2018, with the minor enhancement of grain yield to previous year treatments was demonstrated. There grain yield 4.13 t/ha was the superior, on the other hand, grain yield 2.54 t/ha was the inferior derived from T₈ and T₀, respectively. The production of grain yield under various soil amendments went upwards in the last year of study. In the last year, influences of different treatments on grain yield were found to be ranked in the following descending order T₈ (4.25 t/ha) >T₄ (4.09 t/ha) >T₅ (3.85 t/ha) >T₇ (3.64 t/ha) >T₃ (3.60 t/ha) >T₂ (3.34 t/ha) >T₆ (3.27 t/ha)

In grain production there was a non-significant interaction among the treatment combinations at harvest, except final year of the investigation. Though there was no significant variation among the treatment combinations in 2016-2017, but the most efficient treatment combination for higher grain yield was found in V_3T_8 (4.10 t/ha). Without major change, minor increase in grain yield as like tendency was observed from the treatment combinations in 2017-2018. Here the treatment combination V_3T_8 yielded the maximum grain (4.17 t/ha), while the minimum was recorded in V_3T_0 (2.55 t/ha). The treatment combination V_3T_8 was produced the highest grain yield (4.35 t/ha) at the last year of the research which was the best treatment combination for grain yield.

The treatment combination V_1T_8 was defined as the second major (4.25 t/ha) followed by V_3T_4 (4.14 t/ha) on the basis of grain production records, whereas V_3T_0 had the least (2.34 t/ha) grain production shown in Table 33.

4.4.14 Straw yield

The Fig. 21, 22 and 23 represent that the data related to straw yield was affected significantly by the different wheat varieties. Among the different varieties, the maximum straw yield (4.76 t/ha) was recorded in V_3 and it was statistically different from V_1 (4.69 t/ha) and V_2 (4.64 t/ha) in 2016-2017. The straw yield of wheat resulted in a little increase in 2017-2018 and that was recorded the greater and the lesser from V_3 (4.84 t/ha) and V_1 (4.75 t/ha). As in 2017-2018, the parallel varietal effect on straw yield was revealed in 2018-2019 and had the maximum production at V_3 (4.96 t/ha) which was followed by V_2 (4.89 t/ha) and V_1 (4.87 t/ha), respectively.

Soil amendments application significantly influenced the straw yield all over the research period (Fig. 24, 25 and 26). The positive effect of organic amendments in straw yield was greater (5.26 t/ha) in T₈ than other treatments, even though the lowest straw yield was obtained from T₀ (4.30 t/ha) in the period of 2016-2017. The straw yield also varied considerably under the treatments studied in 2017-2018. The maximum (5.36 t/ha) straw yield was recorded by T₈, but the minimum (4.17 t/ha) straw yield was reported to T₀. Among the different soil amendments, T₈ (5.40 t/ha) was performed superior besides T₄ (5.31 t/ha) to other treatments in 2018-2019. Rest of the treatments were showed also significant difference ranked as descending in T₅ (5.14 t/ha), T₇ (5.05 t/ha), T₃ (4.93 t/ha), T₆ (4.83 t/ha), T₂ (4.73 t/ha), T₁ (4.60 t/ha), while the lowest one was found in T₀ (4.18 t/ha).

The straw yield showed significant variations at various combinations of treatments and varieties during the research period (Table 33). Increasing development of straw yield was noticed under different treatment combinations in 2016-2017, where the combination V_2T_8 (5.53 t/ha) was produced greater straw yield followed by V_1T_8 (5.22 t/ha), V_3T_8 (5.20 t/ha) and V_2T_4 (5.16 t/ha), while the lowest was observed in V_2T_0 (4.20 t/ha). On the other hand, with statistical significance the maximum straw yield was recorded in V_2T_8 (5.48 t/ha) followed by V_1T_8 (5.34 t/ha) and V_3T_8 (5.26 t/ha) in 2017-2018. Comparable results were found in 2018-2019 with the previous two years regarding straw yield. But the increased straw yield value was documented in case of V_2T_8 (5.50 t/ha) and oppositely the decreased value was observed in case of V_1T_0 (4.18 t/ha).




Fig. 21: Effect of variety on grain and straw yield of wheat in 2016-2017.




Fig. 22: Effect of variety on yield and straw yield of wheat in 2017-2018.




Fig. 23: Effect of variety on grain and straw yield of wheat in 2018-2019.




Fig. 24: Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2016-2017.




Fig. 25: Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2017-2018.




Fig. 26: Effect of organic and probiotic soil amendments on grain and straw yield of wheat in 2018-2019.

4.4.15 Harvest index

A summary of varietal influences of this study on harvest index was described in Fig. 27. The highest harvest index (41.38%) was noted by the variety V_3 and this was statistically at par with the V_1 (41.27%) in 2016-2017, at the same time the smallest harvest index (40.47%) was obtained in V_2 . Likewise previous year, the harvest index of the second year varied significantly with the variety factor. But the scale of increase in harvest index was a little bit more, where the V_3 was reported the maximum harvest index (41.62%) and the lowest (40.80%) in V_2 . During 2018-2019, the maximum harvest index (41.74%) was recorded with V_3 , though the V_2 demonstrated the minimum index (40.81%).

The three years harvest index trend of wheat varied under different soil amendments. In 2016-2017, application of chemical fertilizer (T₈) gave the greater harvest index (43.28%) and poultry manure (T₄) had the second highest value (42.60%) which had statistical equality (42.02%) with T₅. On the other hand, the least result (38.36%) was obtained from T₀. In the next year, treatments were further produced with the similar effect on harvest index with the utmost value (43.53%) and the lowest value (37.93%) by T₈ and T₀, respectively. Extension of the study and further evaluation of data, revealed the highest value 44.05% followed by 43.48% and the lowest one 36.13% was derived from T₈, T₄ and T₀ during 2018-2019 (Fig. 28).

Results in Table 33 indicate that harvest index by different treatment combinations were non-significant in 2016-2017, but significant in 2017-2018 and 2018-2019. The most effective treatment combination for higher harvest index was recorded from V_3T_8 (44.08%), but the minimum was noted from V_3T_8 (38.29%) for the year 2016-2017. As previous year, the treatment combination V_3T_8 resulted significantly the maximum harvest index (44.25%) after that V_1T_8 (43.64%) and V_3T_4 (43.28%), while the minimum was recorded in V_2T_0 (37.69%) from the year 2017-2018. In the last year of the research, the treatment V_3T_8 was produced the highest harvest index (45.02%) that was the best treatment combination for harvest index. The treatment V_1T_8 was defined as the second major (44.14%) followed by V_3T_4 (43.91%) and V_1T_4 (43.72%) on the basis of harvest index records, whereas V_2T_0 had the least harvest index (35.91%).




Fig. 27: Effect of variety on harvest index of wheat in 2016-2017, 2017-2018 and 2018-2019.




Fig. 28: Effect of organic and probiotic soil amendments on harvest index of wheat in 2016-2017, 2017-2018 and 2018-2019.

Variaty	& Treatment	(Grain yield (t/ha	.)	S	straw yield (t/ha	a)	Н	arvest index (%)
vallety e		2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	T ₀	2.68	2.55	2.36	4.32	4.21	4.12	38.29	37.75	36.41
	10	± 0.12	± 0.11	± 0.10p	± 0.20 klm	$\pm 0.21o$	± 0.171	± 0.61	± 0.72 lm	± 0.071
	T_1	2.88	2.93	3.01	4.48	4.52	4.59	39.47	40.02	39.64
	11	± 0.09	± 0.09	$\pm 0.10o$	$\pm 0.22hi$	± 0.20 lm	± 0.23i	± 0.52	± 0.91j	$\pm 0.39k$
	T_2	3.11	3.22	3.33	4.35	4.47	4.63	41.68	41.93	41.83
	12	± 0.13	± 0.13	± 0.14 kl	± 0.16 jkl	± 0.19 mn	$\pm 0.17i$	± 0.54	$\pm 0.85 f$	± 0.07 hi
	T ₃	3.36	3.45	3.57	4.63	4.72	4.85	42.04	42.58	42.39
	13	± 0.14	± 0.14	± 0.15i	± 0.16g	± 0.19 hi	$\pm 0.17 f$	± 0.17	± 0.65 de	± 0.17 fg
\mathbf{V}_1	T_4	3.77	3.82	4.10	5.00	5.05	5.28	43.04	43.06	43.72
v ₁	14	± 0.13	± 0.19	± 0.15 cd	± 0.20d	$\pm 0.22e$	± 0.22c	± 0.15	$\pm 0.53 bcd$	$\pm 0.15 bc$
	T ₅	3.57	3.66	3.85	4.87	4.94	5.12	42.29	42.88	42.93
	15	± 0.14	± 0.15	$\pm 0.15 f$	$\pm 0.17 ef$	$\pm 0.24 fg$	$\pm 0.24d$	± 0.11	± 0.72 cd	± 0.16de
	т	2.95	3.10	3.30	4.42	4.57	4.85	40.05	40.45	40.51
	T_6	± 0.12	± 0.13	± 0.13 klm	± 0.22ij	± 0.19 kl	$\pm 0.24 f$	± 0.21	± 0.79ij	$\pm 0.21 j$
	т	3.42	3.51	3.65	4.90	4.95	5.02	41.10	41.51	42.09
	T_7	± 0.16	± 0.16	± 0.17 hi	$\pm 0.21 \text{ef}$	$\pm 0.20 fg$	± 0.22e	± 0.62	$\pm 0.37 fgh$	± 0.07 ghi
	т	4.01	4.13	4.25	5.22	5.34	5.38	43.46	43.64	44.14
	T_8	± 0.16	± 0.17	$\pm 0.18b$	± 0.24b	± 0.23b	$\pm 0.25b$	± 0.14	$\pm 0.92b$	$\pm 0.12b$
	т	2.57	2.46	2.34	4.20	4.08	4.18	37.99	37.69	35.91
	T_0	± 0.10	± 0.09	± 0.09p	$\pm 0.21n$	± 0.17p	± 0.21 kl	± 0.27	$\pm 0.68m$	$\pm 0.27m$
	т	2.73	2.85	2.93	4.27	4.42	4.51	38.98	39.26	39.40
	T_1	± 0.14	± 0.09	$\pm 0.10o$	$\pm 0.18 mn$	± 0.20n	± 0.19j	± 0.26	$\pm 0.70k$	$\pm 0.24k$
	T_2	2.90	3.16	3.24	4.30	4.55	4.72	40.28	41.01	40.69
	12	± 0.13	± 0.14	± 0.14 lmn	± 0.20 klm	± 0.181	$\pm 0.18h$	± 0.55	± 0.84 hi	$\pm 0.14j$
	T ₃	3.25	3.38	3.46	4.56	4.79	4.83	41.56	41.41	41.75
	13	± 0.15	± 0.14	± 0.14j	± 0.12 gh	$\pm 0.21h$	± 0.22fg	± 0.46	± 0.88 fgh	± 0.13 hi
	T_4	3.70	3.76	4.02	5.16	5.19	5.37	41.78	42.02	42.80
V_2	14	± 0.17	± 0.17	± 0.19de	± 0.21 bc	± 0.20cd	$\pm 0.22b$	± 0.14	± 0.90ef	± 0.14ef
	T ₅	3.37	3.58	3.77	4.86	5.00	5.16	40.98	41.77	42.22
	15	± 0.12	± 0.13	± 0.14 fg	± 0.24ef	$\pm 0.23 efg$	$\pm 0.19d$	± 0.35	± 0.84 fg	$\pm 0.02 gh$
	T_6	2.81	2.97	3.15	4.24	4.44	4.76	39.87	40.11	39.84
	16	± 0.11	± 0.12	$\pm 0.13n$	$\pm 0.20 mn$	$\pm 0.18n$	$\pm 0.22 gh$	± 0.14	$\pm 0.79 j$	$\pm 0.14k$
	T_7	3.29	3.45	3.57	4.81	4.93	5.00	40.64	41.20	41.67
	17	± 0.12	± 0.13	$\pm 0.14i$	$\pm 0.21 f$	$\pm 0.20 fg$	± 0.22e	± 0.15	± 0.74 gh	± 0.11i
	T_8	3.91	4.08	4.15	5.35	5.48	5.50	42.21	42.70	42.98
	18	± 0.19	± 0.20	± 0.20c	± 0.22a	± 0.25a	± 0.23a	± 0.19	± 0.60cd	± 0.18de

Table 33: Interaction effects of variety and soil amendments on grain yield, straw yield and harvest index of wheat.

Variety & 7	Frantmant	(Grain yield (t/ha)	5	Straw yield (t/ha	a)	Н	arvest index (%)
vallety & I	reatment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	T ₀	2.77	2.61	2.39	4.37	4.21	4.24	38.80	38.34	36.05
	10	± 0.12	± 0.11	$\pm 0.10 p$	$\pm 0.20 jk$	$\pm 0.17o$	$\pm 0.20 k$	± 0.61	± 0.841	$\pm 0.07 lm$
	T_1	2.92	3.07	3.22	4.33	4.58	4.71	40.29	40.18	40.62
	11	± 0.11	± 0.12	$\pm 0.12 mn$	± 0.19 j-m	$\pm 0.21 kl$	$\pm 0.20 h$	± 0.16	$\pm 0.86j$	$\pm 0.12 j$
	т	3.18	3.29	3.44	4.56	4.64	4.83	41.10	41.52	41.61
	T_2	± 0.11	± 0.11	$\pm 0.12 j$	$\pm 0.18 gh$	± 0.19jk	$\pm 0.20 fg$	± 0.16	$\pm 0.70 fgh$	$\pm 0.13i$
	т	3.43	3.55	3.76	4.87	4.92	5.11	41.28	41.94	42.38
V ₃	T ₃	± 0.15	± 0.15	$\pm 0.16 fg$	$\pm 0.14 ef$	$\pm 0.21g$	$\pm 0.21 d$	± 0.35	$\pm 0.95 f$	$\pm 0.07 fg$
	т	3.85	3.94	4.14	5.10	5.17	5.29	43.02	43.28	43.91
v ₃	14	± 0.15	± 0.16	$\pm 0.17c$	$\pm 0.22c$	$\pm 0.21 d$	± 0.23c	± 0.62	$\pm 0.84 bc$	$\pm 0.06b$
	T ₄ T ₅	3.63	3.78	3.94	4.85	5.01	5.15	42.78	43.01	43.33
	15	± 0.18	± 0.19	± 0.19e	$\pm 0.23 ef$	$\pm 0.20 \text{ef}$	$\pm 0.23 d$	± 0.51	$\pm 0.69 bcd$	± 0.19cd
	T ₅ T ₆	3.05	3.16	3.35	4.64	4.69	4.87	39.66	40.31	40.75
	1 ₆	± 0.13	± 0.14	$\pm 0.15k$	$\pm 0.19g$	$\pm 0.23ij$	$\pm 0.20 f$	± 0.51	$\pm 0.49 j$	$\pm 0.07 j$
	т	3.50	3.62	3.71	4.95	5.05	5.13	41.45	41.79	41.98
	T ₇	± 0.12	± 0.13	$\pm 0.15 gh$	± 0.24 de	$\pm 0.22e$	$\pm 0.25 d$	± 0.33	$\pm 0.78 fg$	± 0.24gh
	т	4.10	4.17	4.35	5.20	5.26	5.31	44.08	44.25	45.02
	T_8	± 0.19	± 0.19	± 0.20a	$\pm 0.23b$	$\pm 0.26 bc$	$\pm 0.23 bc$	± 0.55	± 0.62a	$\pm 0.07a$
LS	5	NS	NS	*	**	**	**	NS	*	**
CV (%)	1.39	1.55	1.57	1.10	0.93	.084	1.61	0.89	0.65

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, * = 5% Level of significance, CV = Co-efficient of variation.

4.5 Seed quality

4.5.1 Germination

Seed germination differed significantly due to variety factor under the whole research period (Table 34). Significantly the higher value of germination was noticed from V₃ (82.42%), on the contrary, the lesser value was found in V₁ (78.83%) in the period of 2016-2017. The similar movement as previous year was reported by V₃ and that was presented the maximum germination (83.41%) followed by V₁ (82.04%) with statistical identity in 2017-2018. However in the final year, germination of seed was ranked as the descending order V₃ (85.79%) >V₁ (84.24%) >V₂ (83.34%).

In the present experiment, soil amendments influenced the germination of seed with statistical variation. The percentage of seed germination in control (T_0) was 70.50% which increased considerably with addition of the different soil amendments in the first year (2016-2017). There the highest germination was observed in T_4 (88.27%) and subsequently in T_5 (84.75%) and T_7 (83.59%) in comparison with the rest of the treatments. In the second year (2017-2018), equivalent results were derived from the various treatments, where 89.93% was the best performance and 72.53% was the lowest one provided by T_4 and T_5 , respectively. With the advancement of germination percentage, the treatment T_4 highly enhanced (92.08%) the germination followed by T_5 (88.89%), oppositely the treatment T_0 had lower effect (74.78%) on germination in the final year of the study (Table 35).

Though the interaction of variety and soil amendment treatment produced significant variation in the first year, but non-significant result was obtained from the next two years. The treatment V_3T_4 resulted in the maximum germination (90.67%) and subsequently in V_1T_4 (88.38%), while the minimum were found in V_2T_0 (68.93%) in 2016-2017. In the second year, interactions demonstrated the same tendency with minor increase in germination as the previous year showing greater (91.58%) and lowest (70.33%) was found in V_3T_4 and V_2T_0 . In the final year, the V_3T_4 was also demonstrated the highest value (94.42%) followed by V_1T_4 (92.00%) on the basis of germination percentage, whereas V_2T_0 showed the lowest (72.50%) germination percentage in Table 36.

4.5.2 Vigor index

The Table 34 revealed that the records related to vigor indices were affected significantly by the function of different wheat varieties. Among the different varieties, the greater vigor index (31.85) was recorded in V₁ and it was statistically different from V₃ (31.65) and V₂ (31.15) in 2016-2017. The vigor index of wheat was found to increase in 2017-2018 and that was reported the higher and the lower from V₃ (35.93) and V₂ (33.85), respectively. As in 2017-2018, the parallel varietal effect on vigor index was recorded in 2018-2019, where the maximum value (37.75) was recorded in V₃ which was followed by V₁ (36.36) and V₂ (35.20).

The data of vigor index of wheat under different soil amendments showed significantly positive result in the consecutive year of experiment. In 2016-2017, wheat plants that treated with poultry manure (T₄) had the highest vigor index (35.99) followed by *Trichoderma harzianum* (T₅) with the value 34.57, while plants in the control plot had the lowest vigor index (24.70). In 2017-2018, a trend of enhancement was noted against various treatments on vigor index. At that time, the highest vigor index (39.48) was noted under T₄ and the second highest (37.30) in T₅ which was statistically interlinked with T₃ (36.84), while T₀ had the lowest mean (27.49). Comparable trend of vigor index from respective treatment showed that 41.54 were higher subsequently 39.26 and 28.84 was the lowest one obtained in T₄, T₅ and T₀, respectively in 2018-2019 (Table 35).

There was a significant interaction effect of treatments on vigor index except in the final year of the investigation (Table 36). The major efficient treatment for higher vigor index was recorded from V_1T_4 (37.44) followed by V_1T_5 (35.65) and V_3T_4 (35.44), but the minimum was noted from V_2T_0 (24.52) for the year 2016-2017. In 2017-2018, the V_3T_4 resulted in the maximum vigor index (42.47) after that ranked V_1T_4 (38.56) and V_3T_5 (38.42), while the minimum was recorded in V_2T_0 (26.58). In the last year of the research, the treatment combination did not show any significant effect. Though, the combination V_3T_4 produced the highest vigor index (43.79), whereas V_2T_0 had the least vigor index (27.69).

4.5.3 Total soluble protein content

Effect of wheat variety on protein content was affected significantly all over the study period (Table 34). From the experimental findings, it was revealed that the protein content of seed in the period of 2016-2017 showed as descending V₁ (84.62 mg/g FW), V₃ (83.67 mg/g FW) and V₂ (83.19 mg/g FW). Among the varieties, the maximum protein content was noticed from V₃ (86.31 mg/g FW), but the minimum was found in V₂ (83.97 mg/g FW) for the production year 2017-2018. A little bit advancement in seed protein content was recorded in the year 2018-2019, where the greater value was noted by V₃ (86.91 mg/g FW) and the minor was in V₂ (85.73 mg/g FW).

The results presented in Table 35 demonstrate a significant variation on the protein content of seed by the different soil amendments in the whole research time. The obtained results in 2016-2017 showed that the protein content of seed varied significantly from 77.31 to 93.32 mg/g FW. Conversely, the higher protein content of seed (93.32 mg/g FW) was recorded in T₈ followed by T₄ (88.17 mg/g FW), while the lesser protein content of seed was noticed in T₀ (77.31 mg/g FW). The data concerned with protein content of seed in 2017-2018 clearly explained that the maximum protein content (93.80 mg/g FW) was observed with T₈, while the least in control (77.47 mg/g FW). In the last year of study, the protein content of seed under various soil amendments was to have slightly increased. Influences of different treatments on protein content of seed were found to be highest in T₈ (96.43 mg/g FW) followed by T₄ (92.56 mg/g FW), whereas the lowest was found in control (79.10 mg/g FW).

The protein content of wheat seed was responded significantly due to interaction of varieties and soil amendments throughout experimental duration (Table 36). In general, as the T₈ individually presented the greater value of protein content thus the V₁T₈ interaction produced the maximum protein content of seed (95.28 mg/g FW) which was statistical identity with V₃T₈ (94.19 mg/g FW) and oppositely the minimum (77.06 mg/g FW) protein content of seed noticed from V₂T₀ in the season 2016-2017. In the case of season 2017-2018, the treatment combination V₃T₈ resulted in the maximum protein content of seed (96.07 mg/g FW), while the minimum was recorded in V₁T₀ (76.72 mg/g FW). In the final year (2018-2019) of this study, the treatment combination V₃T₈ was produced the highest protein content of seed (100.12 mg/g FW), whereas V₂T₀ had the least protein content (78.16 mg/g FW).

Variety	G	ermination (%	6)		Vigor index		Soluble protein content (mg/g FW)			
	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	
	78.83	82.04	84.24	31.85	34.40	36.36	84.62	85.04	86.86	
V_1	± 1.04c	± 1.07a	± 1.21b	± 0.64a	$\pm 0.59b$	$\pm 0.75b$	± 0.33a	$\pm 0.53b$	± 0.42a	
N7	79.34	80.41	83.34	31.15	33.85	35.20	83.19	83.97	85.73	
V_2	± 1.09b	$\pm 1.18b$	± 1.04c	$\pm 0.72c$	$\pm 0.70b$	$\pm 0.60c$	$\pm 1.01b$	± 1.08c	± 1.06b	
V	82.42	83.41	85.79	31.65	35.93	37.75	83.67	86.31	86.91	
V_3	± 1.08a	± 1.08a	± 1.05a	$\pm 0.74b$	± 0.76a	± 0.66a	±1.05b	± 1.13a	± 1.23a	
LS	**	**	**	**	**	**	**	**	**	
CV (%)	0.82	3.34	1.79	0.97	3.16	2.91	1.58	1.61	1.42	

 Table 34: Effect of variety on seed quality of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI-28, V_2 = BARI-29, V_3 = BARI-30, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.

Treatment	C	Germination (%	b)		Vigor index		Soluble protein content (mg/g FW)		
Heatment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
T ₀	70.50	72.53	74.78	24.70	27.49	28.84	77.31	77.47	79.10
0	± 0.92h	$\pm 0.93g$	$\pm 0.85g$	$\pm 0.42h$	± 0.40f	$\pm 0.27 f$	± 0.28e	$\pm 0.58e$	$\pm 0.42e$
T_1	75.31	77.54	79.67	28.20	32.52	34.39	78.20	79.68	79.58
1	± 1.08g	± 1.02f	± 1.18f	± 0.55g	± 0.54e	± 0.36e	± 0.45e	± 1.05d	± 0.52e
T_2	78.14	80.88	83.67	31.69	34.42	35.93	80.25	80.97	82.98
12	$\pm 0.82 f$	± 0.98 de	± 0.96d	$\pm 0.85e$	± 0.96d	$\pm 0.38d$	± 0.46d	± 0.73 cd	± 0.39d
т	81.08	83.86	86.17	32.60	36.84	38.36	80.20	81.98	82.24
T_3	$\pm 0.85d$	± 0.92bc	± 0.94c	± 0.43d	± 0.49b	± 0.34b	± 1.02d	± 1.44c	± 1.28d
T	88.27	89.93	92.08	35.99	39.48	41.54	88.17	93.30	92.56
T_4	± 1.05a	± 1.01a	± 1.24a	± 0.73a	± 0.84a	$\pm 0.58a$	± 1.89b	± 0.90a	$\pm 0.70b$
т	84.75	86.52	88.89	34.57	37.30	39.26	87.59	88.30	91.03
T_5	± 1.06b	± 1.15b	$\pm 0.84b$	$\pm 0.52b$	$\pm 0.58b$	$\pm 0.46b$	$\pm 0.59b$	$\pm 0.66b$	$\pm 0.99c$
т	78.76	78.40	81.83	30.21	34.68	35.37	81.51	81.31	82.62
T_6	± 0.89e	± 1.46ef	± 1.42e	$\pm 0.46f$	$\pm 0.33d$	± 0.33 de	$\pm 0.72c$	$\pm 0.63c$	± 0.70d
т	83.59	83.33	86.28	33.56	35.78	36.98	87.88	89.17	91.99
T_7	± 0.93c	± 1.05 cd	± 0.96c	$\pm 0.43c$	$\pm 0.38c$	$\pm 0.42c$	$\pm 0.58b$	$\pm 0.83b$	$\pm 0.67 bc$
т	81.37	84.57	86.76	32.46	34.03	37.27	93.32	93.80	96.43
T_8	$\pm 0.86d$	$\pm 1.05bc$	$\pm 01.16c$	$\pm 0.50 d$	$\pm 0.34d$	$\pm 0.48c$	± 1.08a	± 1.20a	± 1.14a
LS	**	**	**	**	**	**	**	**	**
CV (%)	0.82	3.34	1.79	0.97	3.16	2.91	1.58	1.61	1.42

Table 35: Effect of organic and probiotic soil amendments on seed quality of wheat.

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.


Variety & Treatment		Germination (%)				Vigor index		Soluble protein content (mg/g FW)		
variety o	2 Treatment	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
T ₀		70.17	72.99	75.25	25.24	27.50	28.95	77.20	76.72	80.17
	10	$\pm 1.42m$	± 1.48	± 1.52	± 0.360	± 0.401	± 0.59	± 0.24ij	± 1.791	± 0.71jk
	T_1	72.65	77.58	79.17	27.32	33.59	34.98	77.58	78.34	80.02
	11	± 1.681	± 1.85	± 1.83	$\pm 0.55n$	± 0.44 hij	± 0.81	± 0.91 hij	± 0.47 jkl	± 0.94jk
	т	76.23	80.32	83.67	32.33	33.31	36.20	81.35	81.85	84.01
	T_2	±1.46j	± 1.74	± 0.97	$\pm 0.75h$	± 0.67ij	± 0.42	$\pm 0.02 efg$	± 0.97 gh	$\pm 0.75h$
	T_3	78.62	83.74	86.33	33.31	36.01	37.51	82.66	85.53	86.58
	13	± 1.14i	± 2.18	± 1.67	± 0.39g	± 0.42 c-f	± 0.54	± 0.26e	± 0.23ef	$\pm 0.26g$
V	т	88.38	90.16	92.00	37.44	38.56	41.02	93.70	95.11	92.28
\mathbf{V}_1	T_4	$\pm 2.04b$	± 1.56	± 1.59	± 0.97a	$\pm 0.33b$	± 0.95	$\pm 0.82a$	± 1.94ab	± 0.82 cde
	т	84.57	85.44	88.08	35.65	36.10	38.82	86.02	87.39	90.39
	T_5	± 2.20de	± 1.97	± 2.03	± 0.72b	± 0.52 c-f	± 1.01	± 0.02d	$\pm 0.02e$	± 0.53ef
	т	74.27	79.48	81.65	29.37	34.77	35.77	82.10	81.22	83.96
	T_6	± 1.50k	± 1.72	± 1.65	± 0.421	± 0.50 f-i	± 0.52	± 1.67ef	± 1.18 ghi	± 1.23h
	T_7	82.73	82.37	84.82	34.52	35.86	36.89	85.66	86.32	90.06
		± 1.43fg	± 1.80	± 1.71	$\pm 0.80e$	± 0.83 c-g	± 0.85	$\pm 0.48d$	$\pm 0.51e$	± 1.31f
	т	81.84	86.28	87.17	31.48	33.87	37.12	95.28	92.93	94.32
	T ₈	± 2.13g	± 1.91	± 1.76	± 0.55ij	± 0.59g-j	± 0.64	± 0.81a	± 2.16bc	± 0.26bc
	T_0	68.95	70.33	72.50	24.52	26.58	27.69	77.06	79.32	78.98
	10	± 1.39n	± 1.02	± 1.05	± 0.35p	± 0.151	± 0.97	± 0.90ij	± 0.21ijk	$\pm 0.70k$
	T_1	74.58	75.98	78.33	28.45	30.74	32.84	77.62	76.60	78.57
	11	± 1.08k	± 1.54	± 2.04	$\pm 0.66m$	$\pm 0.71k$	± 0.99	± 0.21 hij	± 1.121	$\pm 0.46k$
	T_2	77.82	80.11	82.58	31.21	32.19	33.05	78.48	79.01	81.89
	12	± 1.80i	± 1.85	± 1.43	± 0.63j	± 0.93jk	± 0.86	± 0.21 hij	± 0.24 ijkl	± 0.25 hij
	T_3	82.29	83.22	84.92	32.24	36.86	38.39	76.69	76.87	78.26
	13	± 1.43g	± 1.44	± 1.72	$\pm 0.37h$	± 1.17b-e	± 0.78	± 1.34j	± 1.79kl	$\pm 1.37k$
V_2	T_4	85.76	88.04	89.83	35.08	37.42	39.81	89.79	91.42	94.61
• 2	▲ 4	± 1.65c	± 2.03	± 1.30	± 0.91 cd	$\pm 0.76bcd$	± 0.58	$\pm 0.24b$	±0.81cd	$\pm 0.57b$
	T_5	83.53	85.28	87.92	33.28	37.37	38.53	87.06	87.39	89.81
	- 5	± 0.97ef	± 1.88	± 1.71	± 0.58g	± 1.08bcd	± 0.89	± 0.77 cd	± 0.49e	$\pm 0.27 f$
	T_6	80.29	74.17	80.45	30.62	34.12	34.00	82.44	82.72	81.36
	- 0	± 1.62h	± 2.14	± 0.98	$\pm 0.71k$	± 0.79f-j	± 0.79	± 0.93ef	± 1.21g	± 1.65ij
	T_7	82.64	82.38	86.19	32.96	35.47	36.36	89.09	89.94	93.29
	- /	± 1.91fg	± 1.80	±1.03	± 0.48g	± 0.51 d-h	± 0.53	±0.02bc	± 1.06d	$\pm 0.25bc$
1	T_8	78.19	84.15	87.34	32.00	33.90	36.17	90.50	92.41	94.85
	0	± 1.13i	± 1.85	±1.04	± 0.65 hi	± 0.69g-j	± 0.73	± 2.11b	± 2.42cd	± 2.21b

 Table 36: Interaction effects of variety and soil amendments on seed quality of wheat.

Contd	
-------	--

Variaty &	Traatmant	Germination (%)				Vigor index		Soluble protein content (mg/g FW)		
Variety & Treatment		2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019	2016-2017	2017-2018	2018-2019
	T ₀	72.39	74.29	76.58	24.33	28.38	29.88	77.67	83.00	78.16
	10	± 1.251	± 1.78	± 1.55	± 0.63p	± 0.981	± 0.52	$\pm 0.02 hij$	$\pm 0.02g$	$\pm 0.24 k$
	т	78.69	79.06	81.50	28.81	33.23	35.35	79.39	77.46	80.17
	T_1	± 1.82i	± 2.05	± 1.41	$\pm 0.50 m$	$\pm 0.58ij$	± 0.92	± 0.70 ghi	± 1.36kl	± 1.17jk
	т	80.36	82.21	84.75	31.53	37.76	38.53	80.93	82.05	83.04
	T_2	± 1.16h	± 1.90	± 1.96	± 0.64ij	$\pm 0.98 bc$	± 0.45	$\pm 0.21 efg$	$\pm 1.45 gh$	$\pm 0.25 hi$
Υ.	T ₃	82.32	84.63	87.25	32.24	37.67	39.19	81.25	83.53	81.89
		± 1.66g	± 1.71	± 1.26	$\pm 0.84h$	$\pm 0.83 bc$	± 0.79	$\pm 0.95 efg$	$\pm 0.98 fg$	± 0.49 hij
	T_4	90.67	91.58	94.42	35.44	42.47	43.79	81.01	93.36	90.80
V_3		± 2.36a	± 1.59	± 1.91	$\pm 0.82 bc$	± 0.86a	± 1.01	$\pm 0.24 efg$	± 1.36bc	$\pm 1.06 def$
	т	86.16	88.85	90.67	34.77	38.42	40.44	89.69	90.11	92.89
	T ₅	± 1.00c	± 2.05	± 2.04	$\pm 0.50 de$	± 1.11b	± 0.58	$\pm 0.24b$	± 1.58d	$\pm 2.97 bcd$
	т	81.72	81.56	83.39	30.65	35.14	36.35	80.00	79.99	82.53
	T_6	± 1.89g	± 1.88	± 1.93	$\pm 0.35k$	± 0.41 e-i	± 0.42	± 0.94 fgh	± 0.02 hij	± 0.02 hi
	т	85.39	85.25	87.82	33.18	36.01	37.69	88.91	91.25	92.63
	$T_7 \pm 1$	± 1.73cd	± 1.97	± 2.03	$\pm 0.77g$	± 0.83 c-f	± 0.87	$\pm 0.27 bc$	± 0.54 cd	± 0.82 bcd
	т	84.07	83.29	85.75	33.90	34.32	38.53	94.19	96.07	100.12
	T_8	± 1.21e	± 1.92	± 1.98	$\pm 0.69 f$	± 0.69 f-i	± 0.78	± 1.65a	± 1.68a	± 0.59a
LS		**	NS	NS	**	**	NS	**	**	**
CV (%)		0.82	3.34	1.79	0.97	3.16	2.91	0.90	1.58	1.61

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per DMRT. V_1 = BARI wheat-28, V_2 = BARI wheat-29, V_3 = BARI wheat-30, T_0 = Control, T_1 = Rice straw + vermicompost + green manure, T_2 = Cow dung + vermicompost + green manure, T_3 = Compost + vermicompost + green manure, T_4 = Poultry manure + vermicompost + green manure, T_5 = *T. harzianum* + vermicompost + green manure, T_6 = Mung bean residue + vermicompost + green manure, T_7 = *T. viride* + vermicompost + green manure, T_8 = Chemical fertilizer, DAS = Days after sowing, NS = Non-significant difference between initial and final values, LS = Level of significance, ** = 1% Level of significance, CV = Co-efficient of variation.



5. DISCUSSION

This section represents the discussion of results of the investigation in relation to the effect of soil amendments on physical and chemical properties of soil, seedling sustainability, crop growth, yield and yield contributing components and seed quality of wheat through utilization of probiotic and organic manures. Throughout the research, parallel results were shown for T_8 (Chemical fertilizer) and T_4 (Poultry manure + vermicompost + green manure) with few exceptions. Chemical fertilizer can give instant results, but its long term effect is not good for crops and soil. On the contrary, organic soil amendments have many positive effects on soil and crops. Therefore, organic soil amendment has given emphasis in this finding.

5.1 Effects of soil amendments on soil properties

5.1.1 Physical properties

As a physical component, soil moisture has an important role on plant nutrient availability and uptake. At the first cropping season of study, the moisture content of the amended soil presented the negative value over initial value. Then moisture content of soil was found as an increasing trend in the next two years of study except no amendments (T_0) and with chemical fertilizer (T_8). In the study period, the maximum decrease of moisture was recorded from T_8 (-3.33%) in 2016-2017 and the maximum moisture increase was found in T_3 (+0.87%) during 2018-2019. It might be due to the effect of the nature and decomposition rate of organic matter added to the soil. As also may be due to improvement of soil physical property like aggregation of soil particles and water holding capacity as a result of organic waste applications. The findings regarding soil moisture content was also consistent with the findings of Vengadaramana and Jashothan (2012) and Desta (2015).

Bulk density is a vital factor to define the soil structure which also explains the soil quality, productivity and porosity (Almendro-Candel 2018). From this study, decrease of soil bulk density was recorded under organic inputs, added plots and increase in chemical fertilizer treated plots. Considering the initial and final value of bulk density, the higher significant

deviation (-0.08 g/cc) and significant increase (+0.10 g/cc) was recorded from treatment T_3 and T_8 , respectively in the last year of the study. The reason behind this result may be for the potentiality of compost in improving soil physical characteristics such as aggregation of soil particles. This statement was at par with Mbah and Onweremadu (2009) findings. Additionally, Zink and Allen (1998) also mentioned the improvement of soil aggregation through organic amendments with compost. Brown and Cotton (2011) also stated that organic fractions reduce the total weight of soil which results in minimal bulk density.

Particle density helps to understand the physical property of a soil sample which indicates presence of organic matter (Abdalmoula and Alias 2016). During the study, it has been observed that addition of organic manure decreases the particle density of soil. From the comparison of different amendment treatments, T_2 and T_3 showed the higher capability to reduce particle density (-0.02 g/cc) of soil round the study period. It may be due to higher organic matter content of manure and its impact on soil physical properties improvement. This strong association was previously notified by Celik et al. (2004) and Rasoulzadeh and Yaghoubi (2010).

Indeed, porosity is a pathway of air and water movement and residence of the root which is an essential part of soil structure (Nimmo 2004). However, the porosity of soil was increased by the application of organic amendments in this research. The highest positive change of porosity (+2.83%) was recorded from the application of T_3 and the second most (+1.82%) from T_2 . It might be due the fact of addition of organic manure which increases the organic matter status of soil including particle binding agents and thus led to improved soil porosity. These results are in agreement with the findings of Kirchmann and Gerzabek (1999), Vasilica (2009) and Woignier et al. (2016). Furthermore, Marinari et al. (2000) was reported that addition of organic fertilizers and compost increased total soil porosity.

Different soil particles aggregate *viz*. sand, silt and clay form the texture of soil. Inherent soil texture may be influenced by any physical disorder but changing of soil texture needs a long time. From the analysis of soil particles, it was observed that a little change has been made over initial value through the study period. Analyzed data presented the decrease of

sand and clay particles, whereas increase of silt particles by all the organic amendments and greater in treatment T_3 . Such kind of variation may be due the result of organic matter addition which prohibits the dispersion of particles and helps to improve soil structure and texture. Although there was variation in particles content percentage, but soil texture remained unchanged. It may take long time to change the soil texture. But within short time, the texture can not be improved using organic matters that have been occurred in our study. It was reviewed by many researchers that organic manure improves the soil structure and texture. This finding is in line with the opinion of Cooperband (2002), Adesodun et al. (2005) and Mahmood et al. (2017).

5.1.2 Chemical properties

Soil pH that relates with soil acidity or alkalinity is an important issue for soil fertility which showed deviation from initial alkaline soil under all the treatments applied in this study. The soil pH was declined maximum (-0.10 unit) with the amendments of T_2 and T_3 after three years of study. This decline was possibly associated with the application of organic manures which may produce organic acids and pushes the change to neutral (Wakene et al. 2005). In this respect, Sumner (2009) stated that to create hospitable soil environments organic matter slow down acidification and buffer soil pH. Brautigan et al. (2014) treated the organic amendments as a means of reducing soil pH in alkaline soil.

Application of organic inputs slightly increased the organic matter content of soil over initial in the study time. The gain of organic matter may be for the cumulative action of different organic amendment inputs which were led for such increased accumulation. Here the maximum significant accumulation (+0.14%) was noted in T₃. Many researchers were stated the similar result on organic matter enhancement by adding organic manures *viz*. Pattanayak et al. (2001), Sarwar et al. (2003) and Bhogal et al. (2018). Furthermore, organic matter declining is associated with less application or less availability of organic manure, continuous cropping, removal of crop residues and excessive tillage (Ladha et al. 2004). While a sustainable cropping system with reasonable nutrient management practices might be adopted by organic amendment of soil like compost (Makinde 2007).

Except the first year, the rest two years of the study produced increments of total N by organic amendments of soil. There most pronounced (+0.013%) treatment was poultry manure + vermicompost + green manure (T_4). It might be due addition of biomass (Ali et al. 2018) and also biological nitrogen fixation capability (Bista and Dahal 2018) of the treatment component. This was also credited by adequate supply of organic matter and higher percentage of N by poultry manure. The presence of N helps microbial activity for decomposition of organic manures that residual effect was retained in the soil. This result is in line with the findings of Iqbal et al. (2008), Ogunbanjo (2010) and Omara et al. (2019).

Chemical fertilizers and organic manures are the dominant source of K input to soil. The Table 12 showed that K value was reduced at the beginning of the study by all the amendments practice, while increased in the next two cropping seasons. Among the treatments maximum increase (+0.015 cmol (+)/kg) of K was noted by poultry manure (T₄) application. There all organic amendments including chemical fertilizer were found to be a positive supplement of K to soil. This might be due to combined application of each organic amendment and recommended dose of chemical fertilizer which releases K and reduction of K fixation that retained exchangeable K. Some earlier workers mentioned the similar findings as found in the study (Bharadwaj and Omanwar 1994, Ali et al. 2009). Abubakar and Ali (2018) also evaluated the effect of poultry manure as the alternate refill of NPK.

Phosphorus (P) as a major plant nutrient acted on stem elongation and hardening, disease prevention and development of the root system which resulted in better crop growth and yield. At the end of first year of study, all the applied treatments demonstrated negative P balance over initial value. It may be due to less presence of P and higher uptake by plants. But after the first cropping season i.e. in the next two years, the presence of P reported as positive by the most applied amendments except control. Where the maximum (+5.42 μ g/g) P balance was found at the second cropping season as well as in third season (+4.05 μ g/g) by T₃. Such increment of P may be due to residual effect of organic manures and recommended doses of fertilizer to soil. The outcome is in harmony with the results by Shen et al. (2011).

Beside this, organic amendment with compost, cow dung and poultry manure played a vital role of retaining P in available form and improving P fertility status in soil (Reddy et al. 2000, Eckhardt et al. 2018).

Sulphur (S) is an important micro nutrient for plants. Plant growth, seed production, protein and chlorophyll content are linked with S in wheat production (Girma et al. 2005). In this study, most of the organic amendments including chemical fertilizer had the positive value in S retention in soil. Among organic amendments compost (+1.45 μ g/g), cow dung (+1.11 μ g/g) and poultry litter (+1.31 μ g/g) showed slight enhanced values. This might be due to the reason for higher organic matter, microbial decomposition and mobilization of S derived from organic manure. Such increment of S was also reported by Paulsen (2005), Eriksen (2009), Forster et al. (2012) and Gao et al. (2017).

Zn is an essential micronutrient for wheat and organic outputs are constructive to agroenvironment (Ali et al. 2019). In the study, changes of Zn exhibited negative value over initial under all the amendments during 2016-2017 cropping year. But positive change of Zn was reported by all the amendments excluding control in the sequential two years, where the treatment T_4 (+0.06 µg/g) followed by T_3 (+0.05 µg/g) added higher balance. This was detectably attributed to addition of organic manures and its efficacy to deliver micronutrients with solubilizing capacity of this nutrient (Behera et al. 2011). Furthermore, there is also some evidence for such findings presented by Antil and Singh (2007) and Ru et al. (2012).

Soluble salts concentration of growing media and ability to conduct electrical current is expressed by electrical conductivity (EC) (Schaefer et al. 2007). In this study, all amendments performed as vehicle of EC enhancer over initial but not in control. The maximum influence (+13 μ s/cm) from starting (145 μ s/cm) to end value (181 μ s/cm) of the study was recorded by the application of poultry manure (T₄) followed by compost combination (T₃) amendment (+11 μ s/cm). Such increase of EC can be occurred due the application of organic manures for soil amendment that leads water holding capacity and available selected soil nutrients with its reasonable mineralization and sorption of ions that prevail in the soil (USDA 2011). These results are in harmony with the findings of Zhao et al. (2014), Carmo et al. (2016) and Bhatt et al. (2019).

Organic manures applied in the soil need to be decomposed for release nutrients by soil microbes. The C:N ratio is varied from 8-10:1 in soil and the ratio is an indicator of nitrogen availability in soil. Microorganisms break down the applied organic manures in soil to get carbon and then release nitrogen for plants (Brown 2015). The high C:N ratio (>30) content materials may create deficiency of nitrogen to plants due slow break down of organic manures. In this study, most of the amended materials had a C:N ratio not more than 30. The data collected from the research revealed that positive balance of C:N ratio in the first year was exhibited by all treatments, but negative in the rest two years except in T_0 and T_8 (Table 17). There the lowest (8:1) and the highest (23:1) C:N ratio was noticed from T_4 and T_0 , respectively. It might be due to deficiency of nitrogen in the first year and lower decomposition, but in the next two seasons higher decomposition with available nitrogen reduced the C:N ratio. Thus organic soil amendment practices directed such change of C:N ratio and that was reported by many researchers (Huang et al. 2001, Hachicha et al. 2006, Yaacob et al. 2019).

5.2 Effects of soil amendments on growth and yield of wheat

5.2.1 Seedling survivability

A significant variation of seedling infection was noticed all over the research period. The trend of infection was found to decline with the progress of the study due to application of probiotic and organic manure. The highest rate of seedling infection was found in T_0 (2.32%), while the minimum rate of infection was recorded in T_5 (0.32%) from 2018-2019. It might be due to the definite role of probiotic fortified-organic manures that exerted antagonistic effect against the fungi causing seedling infection. Furthermore, the addition of organic matter may create a hospitable environment for growth and development of fungal antagonists resulting in enhanced biocontrol activity (Sarkar et al. 2002). This finding was in harmony with the reports Tewari and Singh (2005).

Seedling infection later on converted into seedling blight. Like seedling infection, this parameter was also varied significantly under various treatments all over the study period. The control activity against seedling blight was highly observed in T_5 (0.87%) and the lowest exhibited from check in T_0 (4.74%) during 2018-2019. Such result may be obtained for the efficacy of the prohibitory and biocontrol action of the treatment. This efficacy of *Trichoderma* fortified-organic manures was evaluated to reduce the pre-emergence and post-emergence seedling mortality, diseases of stem and root of chickpea by Talukder et al. (2017). Similar control activity of *Trichoderma* spp. against plant pathogen and seedling mortality was also reported by Bhuiyan et al. (2007) and Ha (2010).

5.2.2 Leaf chlorophyll content

Chlorophyll content of leaf from diverse treatments was provided significant dissimilarities from beginning to end of the study. On the focus of organic amendments, T_4 (48.37 SPAD) next to T_8 (51.48 SPAD) had the highest accumulation of chlorophyll and the smaller was noticed in T_0 (40.33 SPAD). It may probably relate with the decomposition rate of organic manures that happened in case of T_4 treatment. This result was supported by Khan et al. (2005) with the statement that N supply in organic treatment is generally restricted and slows N mineralization as compared to crop demand of N. Chlorophyll content is closely related to the presence of N and poultry manure (T_4) had the higher ability of supplying N. This result is an agreement with the findings of Sims and Wolf (1992), Khanam et al. (2001), Ru et al. (2012) and Krishna (2013).

5.2.3 Growth attributes

The soil amendments demonstrated a remarkable significant variation on TDM at all DAS. Generally, with the advancement of time total dry matter (TDM) accumulation of wheat plants positively progressed during the research time. This progress of TDM at later stages might be used for the development of a large number of late tillers (Balyan 1992, Rahman et al. 2006) resulted in increased TDM. The accumulation of TDM was commonly exhibited the higher in T₈ (767.14 g/m²) and it was statistically identical with T₄ (750.91 g/m²) at 85 DAS in 2018-2019. The improvement of dry matter accumulation possibly due to high content of essential nutrients present in poultry manure which

accelerates vegetative growth (Muhammad et al. 2018). This finding was also supported by Bakhtiar et al. (2017) by the statement that N in poultry manure promoted leaf, stem and spike growth which enhanced plant vegetative growth.

Leaf area index (LAI) expresses the efficacy of photosynthetic processes which is related with crop vegetative growth as also crop yield. The leaf area had an ascending tendency up to 55 DAS, whereas descending nature exhibited in the next 70 to 85 DAS during the three years of experiment. At the last sampling date, T_8 (3.73) and T_4 (3.39) showed the statistical identical value of LAI and a similar tendency was observed in the consecutive year 2017-2018. It might be due to slow and continuous release of plant nutrients from organic manures which directed more leaf production resulting in increased leaf area. Ogundare et al. (2012) and Kareem et al. (2017) confirmed the judgments of the research outcome. Furthermore, Al-Rawashdeh and Abdel-Ghani (2008) supported the decline of leaf area index due to losses of nitrogen (gaseous form) at the end of vegetative stage.

Generally, crop growth rate (CGR) was demonstrated lower increase at the initial stage of wheat growth and later it reached the peak at 55-70 DAS and then declined. Most of the sampling dates revealed that T_8 had the highest activity in CGR followed by T_4 but at the end of CGR sampling (70-85 DAS) T_4 showed the greater influence. Thus, the greater CGR value 21.24 g/m²/d was followed by 19.89 g/m²/d in T_8 and T_4 , respectively. This slow-fast-slow growth rate trend might be due to the slow availability of nutrients that produced less dry matter at the early stage with poultry manure and later stage produced more dry matter almost similar to chemical fertilizer for releasing more nutrients. This observation may be linked with the findings of Geng et al. (2019). Afterwards the decrease of CGR was noticed due to leaf senescence and tiller drying. This result is an agreement with the observation of Boateng et al. (2006) and Ram (2017).

5.2.4 Yield and yield contributing attributes

Throughout the research duration, T_8 and T_4 showed the greater influence on plant height. During the study, the maximum (93.81 cm) and closely the next higher (91.29 cm) plant height was noticed from T_8 and T_4 , respectively. It might be due to rich and balanced nutrients (N, P, K, Ca and Mg) and least C:N of poultry manure which encouraged quick nutrient uptake and sustainable performance. Furthermore, Rehman et al. (2010) stated that organic manure increases plant cell division which results in taller plant production. These findings are in line with the results of Umanah et al. (2003), Gowda et al. (2010) and Shahid et al. (2015).

The parameter plant/m² is one of the major vegetative growth indicators of wheat. The best values of plant/m² were obtained from T_8 (216.11) and then T_4 (202.81) where the lowest at T_0 (158.66). The ability of such enhancement plant/m² due to application of poultry manure might contain high amounts of nutrients and organic matter that improved the soil structure and environment which aids density of plant. Result in the study was in the same manner previously notified by Hassan (2002), Dauda et al. (2008) and Ismaeil et al. (2012).

Tillering is one of the important growth stages of wheat which provides the necessary stalks for satisfactory production. Tiller number and its survival may be influenced by weather and soil nutrient status. A gradual advancement of tillering was observed under the soil amendments and significantly higher in T_8 (4.13) and T_4 (3.84). This result is in agreement with the findings of Enujeke (2013) who reported that readily available and easily absorbable nutrients lead to faster growth and development and ultimately to more tillers by application of poultry manure. Besides this, addition of organic matter improved the soil physical condition which influenced the tillering. This result is in accordance with the observation of Bhuiya and Akhand (1983) and Rajput et al. (1988). Though tillers are lateral branches which emerge from the main stem, but not all produce an ear. Generally, the first two-three tillers are noted as productive tillers of wheat and production is related to environment and nutrition supply. In this investigation, a greater number of productive tiller/plant was demonstrated in T_8 (3.89) and T_4 (3.69). This advancement of effective tiller might be due to optimum moisture and available nutrients provided by poultry manure which may have created the favorable condition for wheat growth. This was supported by Simpson (1990), Eck and Stewart (1995), Mitchell and Tu (2005) and Sistani et al. (2008). The result of this investigation was related to the observations of Belefant-Miller (2007), Abbasi and Khaliq (2016).

The long typically needle like appendage of the lemma in spikelet in wheat known as awn. It has relation to photosynthesis and improvement of grain yield and quality (Elbaum et al. 2007, Yuo et al. 2012). Applied treatments as a source of variance significantly influenced the awn length of wheat. Among the treatments, T_8 produced the higher awn length (7.38 cm) and it was statistically at par with T_4 (7.27 cm), whereas the lowest in T_0 (5.36 cm). It might be due to creating soil healthier and nutrient restore for vegetative as well reproductive growth of wheat which was obtained by soil amendment. This finding is partially related in accordance with the observation of Kowsar et al. (2015).

Data regarding spike length was affected significantly by various soil amendments round the research seasons. Results showed that among the treatments, T_8 was produced the utmost spike length (8.60 cm) and then T_4 exhibited the nearer one (8.43 cm) and it was statistically at par with T_8 . It might be due to higher nutrient reserve and moisture retention capacity of poultry manure than other organic manures. This result is correlated with Hussain (2001) and Ahmed et al. (2002). Nevertheless, green manure fortified poultry manure keeps the N balance which enhances the length of spike of wheat. The above statement was also supported by Nguyen et al. (1995).

Spike properties no doughty affect the grain yield. Spikelet/spike treated as an important element of wheat grain yield (Sabaghina et al. 2014). Spikelet/spike in this study revealed variation between 13.80 (T₀) to 17.97 (T₈). These variations are in line with Tahir et al. (2011) and Sarwar et al. (2007) who reported that yield components of wheat significantly influenced organic manures over control. Considering the mean values of spikelet/spike to evaluate the performance of different treatments, the maximum value (17.97) was observed in T₈ and it was statistically identical with T₄ (17.81) all over the research period. It might be due to supply of nitrogen rich organic manures which positively affected the growth and yield attributes of wheat. These findings are partially correlated with the observation of Siavoshi et al. (2011). With the acceptance of the above assumption of N supply, especially during spikelet differentiation and development, T₈ produced the higher fertile spikelet/spike (17.24) and T₄ (17.00) having statistical equality. Similar observations were made by Patel et al. (1995) and Rahman (2004).

Grains/spike is a common yield component for cereals (Slafer et al. 1996, Thiry et al. 2002). Under this experiment the tendency of grains/spike was notified higher from T_8 which was followed by T_4 . These two treatments were shown to have the greater value 45.73 and 44.40 with significant variation compared with the rest of the treatments. Considering the organic amendment, presence of greater N in poultry manure produced more dry matter and partitioning it towards grains which enhanced the grains/spike. This result is confirmatory with the outcomes of Shah et al. (2010), Yildirim et al. (2016) and Mukhtiar et al. (2018).

As a yield component, grains weight/spike of wheat was measured and analyzed by some researchers (Frederick et al. 2001, Peltonen-Sainio et al. 2007). Application of different soil amendment treatment significantly affected the grains weight/spike over the study period. Data across various amendments demonstrated that the maximum weight of grains/spike was noticed in T_8 (2.20 g) afterwards in T_4 (2.14 g), while the least in T_0 (1.60 g). The advancement of grain weight/spike from organic manures treated with poultry manure that may have the balanced nutrient stock during the grain filling stage. Similar reports were also presented by Sevaram et al. (1998), Garg and Bahla (2008).

For cereal crops, 1000-grain weight is an important trait to determine the yield and yield attributes (Metho et al. 1998). The data concerning 1000-grain weight of wheat clearly indicated that the applied treatments T_8 (54.12 g) and T_4 (53.90 g) significantly increased the weight than others. The reason for such kind of nearer advancement by T_4 (poultry manure) may be due to availability of nutrients throughout the growing season and improvement of soil characters. This outcome was agreed with Ma et al. (1999), Haris et al. (2002) and Ahmed et al. (2017).

Different major yield components like spike/m², number and weight of grains/spike and 1000-grain weight mathematically functioned on yield of cereal crop (Thiry et al. 2002). All the soil amendments proved its significant effect on grain yield of wheat. From the mean comparison of different treatments it notified that the maximum grain yield was noted for T₈ (4.25 t/ha) followed by T₄ (4.09 t/ha) and the minimum of it (2.36 t/ha) belonged to T₀. The production of yield by T₄ may be for the reasons of enhancement soil

fertility and improvement of physical conditions, reduction of nutrient losses and preservation of soil biota which contributes to higher grain yield. This result is in conformity with the observation of Sharma and Madan (1988), Shanggan et al. (2000), Khaliq et al. (2004), Islam et al. (2009), Naseri et al. (2010) and Zhang et al. (2016).


Wheat straw yield was responded significantly from the application of various soil amendmentsover the research period. The maximum straw yield (5.40 t/ha) was produced by chemical fertilizer (T_8) and the nearest result (5.31 t/ha) was obtained by poultry manure (T_4). This performance of organic manures might be due to capability of abundant nutrients supply and creation of a favorable growth environment. This turn on increased photosynthetic activity, cell elongation and vegetative growth that was reflected in straw yield. Similar cause and effect was also noted by Auti et al. (1999), Usman et al. (2003), Akhtar et al. (2011), Kumar and Abraham (2018).

Harvest index is an important reproductive efficiency of crops in increasing grain yield. In the present study, the harvest index of wheat was significantly affected by the various soil amendments. Regarding this parameter, the highest value (44.05%) was recorded with the application of treatment T_8 followed by T_4 (43.48%), whereas the least value (36.13%) was obtained from T_0 . The performance of poultry manure on boosting harvest index might be continued supply of nutrients and higher dry matter distribution towards grain. This result is in agreement with the investigation of Swarup and Yaduvanshi (2000), Rees and Castle (2002), Krishnan et al. (2003), Delfine et al. (2005), Kabesh et al. (2009) and Khan et al. (2018).

5.2.5 Seed quality

Introducing designed soil amendments not only developed growth and yield of wheat but also enriched the parameters of seed quality. Significantly greater mean of seed germination was noticed in T_4 (92.08%), whereas the least mean was calculated from T_0 (74.78%). This might be due to adequate supply of nutrients which helped to build up carbohydrate like seed elements thus producing vigorous seeds. The outcome of the investigation is the conformity of previous results of researchers Gowda et al. (2010), Aslam et al. (2011), Raissi et al. (2012) and Peerzada et al. (2016). Vigor index is an imperative quality of seed that indicates the potentiality of constant emergence of seedlings (Pradeep 2018). In the present study, it was observed that soil amendments effectively increased vigor index of seed over control. This upturn was dominantly (41.54) found from the plot which consumed poultry manure (T_4), while the minimum (24.70) was found at control plot (T_0). The speed of germination i.e. vigor index might be significantly differ for providing the substantial macro and micro nutrient in the grain filling stage and stored in seed. This result is in agreement with the reflection of several researchers like Page-Dumroese et al. (1991), Farhad et al. (2011) and Sheoran et al. (2017).

Among the major cereals, wheat is the prominent source of vegetal protein (13%). Protein content in wheat seed showed the variable responses to different soil amendments where finally T_8 produced greater total soluble protein (96.43 mg/g FW) followed by T_4 (92.56 mg/g FW) and the least at T_0 (79.10 mg/g FW). One important part of the study is organic amendment of soil therefore; special emphasis had been given in this case. The increased protein content might be due to availability of high NPK that satisfy the plant requirement. The accumulated N in the leaf mobilized to seed components which leads the protein production. This result is in pact with the findings of Abedi et al. (2010) and Rasul et al. (2015). Moreover, the improvement of protein content might be also due to green manure based crop rotation and poultry manure application which enriched N availability and N uptake by wheat plant. Related findings are reflected by Alam et al. (2018).



6. SUMMARY

The study was carried out for three consecutive years 2016-2017, 2017-2018 and 2018-2019, with a view to find out the effect of probiotic and organic soil amendment on sustainable improvement of wheat (*Triticum aestivum* L.) growth and yield. The experiment was laid out in randomized complete block design (RCBD) with three replications where three wheat varieties *viz*. BARI wheat-28, BARI wheat-29 and BARI wheat-30 and nine soil amendments *viz*. T₀ = Control (without treatment), T₁ = Rice straw + vermicompost + green manure, T₂ = Cow dung + vermicompost + green manure, T₃ = Compost + vermicompost + green manure, T₄ = Poultry manure + vermicompost + green manure, T₅ = *Trichoderma harzianum* + vermicompost + green manure, T₆ = Mung bean residue + vermicompost + green manure, T₇ = *Trichoderma viride* + vermicompost + green manure and T₈ = Chemical fertilizer (recommended dose) were used. The plant growth parameters, yield and yield contributing attributes and seed quality of wheat were studied in the research. Besides these, soil properties in respect of physical and chemical characters were also considered.

From the investigation, it has been found that moisture content of soil was greatly influenced by different amendment practices. There negative moisture balance was also found by all the amendments and -2.50% was the highest deviation occured by treatment T_7 during 2016-2017. After two years, maximum balance of moisture of soil was recorded in T₃. Two closely related physical properties *viz*. bulk density and particle density showed negative changes under different amendments round the research period, but it was positive in case of control. The higher change in bulk density (-0.08 g/cc) and particle density (-0.02 g/cc) was noticed in T₃. Based on the value of bulk density and particle density, the estimated porosity of soil also exhibited significantly better results (+2.83%) found in T₃ treatment. Though texture is a stable property of soil, but it may be changed by disruption of soil particles over a long period. Hence, no change was recorded in texture from initial to final study period but changes were recorded in respect of sand, silt and clay particles. Thus at the end of the study, the maximum significant deviation of sand (-2.33%)

was recorded in T_3 and equally value (-0.67%) deviation for clay particles was found by the T_4 , T_1 and T_6 . Whereas significant increase (+2.67%) of silt particles was recorded by T_3 amendment.

Inclusion of different organic amendments did not significantly influence the soil pH in comparison with initial reading. But increasing numerically trends were observed in case of T₀, T₄ and T₈, while the higher decreased reading was -0.10 unit and that was observed in T₃ and T₂. Applied amendments exhibited the building up trend of organic matter content in the experimental plots except control and the maximum accumulation (+0.14%)was noticed in T₃. Plants macronutrients like total N, available P, exchangeable K and available S were influenced positively by addition of amendment materials except control. There was significantly higher value for total N (+0.013%) and exchangeable K (+0.015 (+)/kg) presented in T₄. On the other hand, under amendment with T₃, available P and available S revealed the higher value as +5.42 and $+1.45 \ \mu g/g$, separately. In this study, secondary nutrients of plants like Zn significantly showed the positive enhancement at the end year of cropping except control. There the T_4 was presented the greater balance $(+0.06 \ \mu g/g)$ of Zn in comparison with other amendments. In respect of electrical conductivity of soil, it was found to be increased over initial value by organic amendments. The most pronounced value (+13 μ s/cm) of electrical conductivity was obtained in T₄. Although the carbon nitrogen ratio presented a positive balance in the starting year but it was found as negative balance in the next two years. At the end year of the study, C:N ratio was observed as higher decline (-0.60) in T₄.

The results revealed that seedling survivability against seedling infection and seedling blight were higher in the V_3 than that of other two varieties in 2017-2018. The survivability of seedling was found greater under the T_5 than other organic soil amendment treatments in the season 2018-2019. The result indicated significantly lower seedling infection and seedling blight. The combination of V_3T_5 exhibited better performance for reducing seedling infection and seedling blight during 2018-2019. In the context of leaf chlorophyll content, V_3 showed greater achievement than other varieties in 2018-2019. Among the organic amendment treatments, T_4 provided more leaf chlorophyll content, especially at the last cropping of the study. Thus the interaction of variety and soil amendments produced insignificantly the greater leaf chlorophyll from the arrangement in V_3T_4 from 2018-2019.

Growth attributes of wheat like total dry matter, leaf area index and crop growth rate were significantly affected by the selected variety. General progressive increase of TDM was recorded at all sampling dates, but the greater TDM was noticed at 85 DAS in V₃ from 2018-2019. The leaf area index and crop growth rate increased up to 55 DAS and then decreased where the V₃ produced the maximum LAI and CGR during 2018-2019. These parameters were found significantly better by the application of various organic soil amendments round the research period, however, the performance was pronounced in T₄ from 2018-2019. Though no significant variation was recorded on the maximum TDM production from the combination V_3T_4 in the season 2018-2019, but the LAI and CGR was significantly influenced by the combination and reached the peak of increase at the last year of the study.

Excluding the year 2017-2018, plant height of wheat at harvest significantly affected by the varietal influence where V_3 showed better height round the research period. Other plant characters of wheat like total plant/m², total tiller/plant and effective tiller/plant also provided the maximum efficiency derived in V_3 during 2018-2019. Among the treatment sets of organic amendments, poultry manure (T₄) demonstrated the best output in relation to the aforesaid parameters in 2018-2019. Furthermore, the combination of variety factor and treatment factor did not affect the above stated parameters significantly, but V_3T_4 prearrangement exposed the greater numerical value all over the research seasons.

All the spike characters as awn length, spike length, spikelet/spike and fertile spikelet/ spike were significantly influenced by varietal factors. Among the wheat varieties, V_3 was superior against the aforesaid spike characters from the first year to last year of the research. On the other hand, T_8 and T_4 showed statistically similar performance in relation to the above spike morphology in the three seasons of the study. But the organic soil amendment T_4 denoted as the best performer with higher values for the mentioned parameters in the period 2018-2019. The selected variety coupled with designated soil amendments had no significant variation on the spike parameters. But from the combination of variety and organic treatments, the combination of V₃T₄ gave the greater numerical results on these spike characters in the year 2018-2019.

Some grain related parameters like grains/spike, grain weight/spike and 1000-grain weight significantly achieved the higher value with V_3 but deformed grains/spike was high with V_1 . In general, under the organic amendment treatment, T_4 significantly enhanced the grain parameters except deformed grains/spike by T_0 during 2018-2019. These parameters were found significantly higher in V_3T_4 arrangement excluding deformed grains/spike at V_3T_0 .

The result exposed that variety had a significant effect on grain yield, straw yield and harvest index of wheat. The above mentioned sequential parameters recorded with the best values as 3.59 t/ha, 4.96 t/ha and 41.74% under V₃ in 2018-2019. All the soil amendment packages also significantly promoted the grain yield, straw yield and harvest index. It appeared that the expressed parameters were greatly influenced by the T₄ in comparison with the organic components of amendment. The major improvement was notified under T₄ regarding grain yield (4.09 t/ha), straw yield (5.31 t/ha) and harvest index (43.48%) in 2018-2019. This development of the yield was approximately 73%, 27% and 20% more over respective control. Due to conjunctive use of variety and organic amendment treatments, the highest grain yield (4.14 t/ha), straw yield (5.37 t/ha) and harvest index (43.91%) were produced by the arrangement V₃T₄, V₂T₄ and V₃T₄, respectively.


The seed quality parameters such as germination percentage, vigor index and total soluble protein content were positively increased. These parameters significantly affected the varietal performance where V_3 treated as the higher producer of germination, vigor index and protein content in 2018-2019. All the treatments of soil amendment significantly influenced the above mentioned seed quality. There was a considerable higher performance noticed by T_4 for all the seed quality parameters studied in this research. The relationship of variety and soil amendments showed non-significant movement in case of germination

and vigor index, whereas significant progress was found in protein content at the final year of study. These relations were higher in case of germination and vigor index in V_3T_4 , but the protein content showed better in V_2T_4 .

The experiment has made known that the application of probiotic and various organic manures improved the soil physical and chemical properties. In this investigation, manure acted as a vital source of organic matter and essential plant nutrients reservoir which influenced the physical characters of soil like moisture, particle density, bulk density and porosity as also build up positive changes in chemical properties like pH, C:N ratio and electrical conductivity of soil as well as plant growth. The above mentioned characters were efficiently modified by the application of green manure and vermicompost fortified poultry manure (T_4). Hence for improvement and sustenance of soil fertility, poultry manure may be considered as an imperative component.

There was a potentiality for sustainable improvement of wheat growth, yield and seed quality with soil amendment through probiotic and organic manures. In that term, inclusion of green manure, vermicompost and poultry manure may be a worthy alternative against chemical fertilizers. The capacity of surprising decomposition and nutrient release of those organic manures could be used to synchronize the nutrient demand of wheat. Organic manures particularly poultry manure was a healthier source for improvement of growth, yield and seed quality of wheat as compared to other manures.

Thus the organic manures i.e. poultry manure combination (T_4) would be an important substitute of chemical fertilizer for its easy availability, less cost, positive residual effects and soil fertility storage that directed to sustainable environmental improvement. So, the above declared manures may be suggested to use for soil amendment and sustaiable wheat production in the studied region.



7. REFERENCES

- Abbasi MK and Khaliq A (2016). Nitrogen mineralization of a loam soil supplemented with organic-inorganic amendments under laboratory incubation. Frontiers in Plant Science, 7: 1-9.
- Abdallahi MM and N'dayegamiye A (2000). Effects of two incorporations of green manures on the yield and nitrogen nutrition of wheat (*Triticum aestivum* L.), as well as on the physical and biological properties of the soil. Canadian Journal of Soil Science, 80(1): 81-89.
- Abdalmoula MM and Alias MA (2016). Investigation of soil physical properties under *Shorea peltata* sym. in Tengaroh Forest Reserve, Johor Bahru, Malaysia. Journal of the Faculty of Forestry, Istanbul University, 66(1): 198-210.
- Abedi T, Alemzadeh A and Kazemeini SA (2010). Effect of organic and inorganic fertilizers on grain yield and protein banding pattern of wheat. Australian Journal of Crop Science, 4(6): 384-389.
- Abubakar ZA and Ali AD (2018). Screening effect of organic manure on the vegetative growth of maize (*Zea mays* L.). Journal of Bioscience and Agriculture Research, 16(2): 1356-1364.
- Adeniyan ON, Ojo AO, Akinbode OA and Adediran JA (2011). Comparative study of different organic manures and NPK fertilizer for improvement of soil chemical properties and dry matter yield of maize in two different soils. Journal of Soil Science and Environmental Management, 2(1): 9-13.
- Adesodun JK, Mbagwu JSC and Oti N (2005). Distribution of carbon, nitrogen and phosphorus in water stable aggregates of an organic waste amended Ultisol in southern Nigeria. Bioresource Technology, 96(4): 509-516.
- Agbede TM, Ojeniyi SO and Adeyemo AJ (2008). Effect of poultry manure on soil physical and chemical properties, growth and grain yield of sorghum in southwest, Nigeria. America-Eurasian Journal of Sustainable Agriculture, 2(1): 72-77.

- Ahmed BEAM, Ishag AA, Hassan MK and Ahmed MO (2017). Response of two wheat cultivars (*Triticum aestivum* L.) to amended nitrogen fertilizer on yield and grain quality in Halfa Elgadidah area. Medcrave Online Journal of Biology and Medicine, 1(5): 133-136.
- Ahmed SI, Abbasi MK and Rasool G (2002). Integrated plant nutrition system (IPNS) in wheat under rain fed conditions of Rawalakot Azad Jammu and Kashmir [Pakistan]. Pakistan Journal of Soil Science, 21(1&2): 79-86.
- Ahsan E and Karim Z (1986). Soil fertility and management research on upland soils in Bangladesh. *In*: Proceedings of the International Conference on the Management and Fertilization on Upland Soils in the Tropics and Subtropics, Nanjing, China, pp. 247-252.
- Akhtar M, Asif N, Javed A, Bokhari SA and Wajid I (2011). Improvement in nutrient uptake and yield of wheat by combined use of urea and compost. Soil in Environmental Science, 30(1): 45-49.
- Akhtar N, Ramani VB, Yunus M and Femi V (2018). Effect of different nutrient management treatments on growth, yield attributes, yield and quality of wheat (*Triticum aestivum* L.). International Journal of Current Microbiology and Applied Sciences, 7: 3473-3479.
- Akter N (2011). Evaluation of soil fertility status of 10 dominant soil series of Jamalpur Sadar Upazila. MS Thesis, Department of Soil Science, Bangladesh Agricultural University, Mymensingh, Bangladesh.
- Alam MM, Karim MR and Ladha JK (2013). Integrating best management practices for rice with farmers' crop management techniques: A potential option for minimizing rice yield gap. Field Crops Research, 144: 62-68.
- Alam MR, Ali MA, Molla MSH, Omin MAM and Annan MAM (2009). Evaluation of different levels of potassium on the yield and protein content of wheat in the High Ganges River Floodplain soil. Bangladesh Journal of Agricultural Research, 34(1): 97-104.

- Alam MS, Haydar FMA and Hoque A (2020). Effects of planting method on yield and yield attributing characters in chickpea (*Cicer arietinum* L.). Journal of Bio-Science, 28(20): 105-110.
- Alam MZ, Lynch DH, Tremblay G, Gillis-Madden R and Vanasse A (2018). Optimizing combining green manures and pelletized manure for organic spring wheat production. Canadian Journal of Soil Science, 98(4): 638-649.
- Ali AM, Ibrahim SM and Bijay-Singh (2020). Wheat grain yield and nitrogen uptake prediction using atLeaf and GreenSeeker portable optical sensors at jointing growth stage. Information Processing in Agriculture, 7(3): 375-383.
- Ali G, Rao CP, Rao AS and Rani YA (2018). Effect of *in situ* incorporation green manures on soil organic carbon, pH, bulk density and economics involved in its incorporation. International Journal of Current Microbiology and Applied Sciences, 7(9): 62-67.
- Ali ME, Islam MR and Jahiruddin M (2009). Effect of integrated use of organic manures with chemical fertilizers in the rice-rice cropping system and its impact on soil health. Bangladesh Journal of Agricultural Research, 34(1): 81-90.
- Ali MM, Rahman MH, Khan MR and Khan MK (2012). Evaluation of soil characters for assessment of land degradation situation in the Ganges River Floodplain area. *In*: Proceedings Research review and planning workshop on soils program of NARS, Dhaka, Bangladesh.
- Ali MM, Saheed SM and Kubota D (1997) Soil degradation during the period 1967-1995 in Bangladesh. II. Selected chemical characters. Soil Science Plant Nutrition, 43(4): 870-890.
- Ali W, Ali M, Iqbal J, Khalil MK and Ahmad Z (2019). Impact of zinc and organic manure on wheat productivity and soil fertility. Advances in Crop Science and Technology, 7(1): 1-4.
- Almendro-Candel MB, Lucas IG, Navarro-Pedreno J and Zorpas AA (2018). Physical properties of soils affected by the use of agricultural waste. Agricultural Waste and Residues, 2: 9-27.

- Al-Rawashdeh YA and Abdel-Ghani AH (2008). Effect of nitrogen application timing on dry matter and nitrogen assimilation and partitioning in six wheat cultivars under rain-fed conditions of Jordan. Archives of Agronomy and Soil Science, 54(2): 149-162.
- Altomare C, Norvell WA, Bjorkman T and Harman GE (1999). Solubilization of phosphates and micronutrients by the plant-growth-promoting and biocontrol fungus *Trichoderma harzianum* Rifai 1295-22. Applied and Environmental Microbiology, 65(7): 2926-2933.
- Antil RS and Singh M (2007). Effects of organic manures and fertilizers on organic matter and nutrients status of the soil. Archives of Agronomy and Soil Science, 53(5): 519-528.
- Aslam M, Khan MA, Awan IU, Khan EA, Khan AA and Jilani G (2011). Effect of single and combined use of various organic amendments on wheat grown over green manured soil: growth and yield attributes. Pakistan Journal of Nutrition, 7(10): 640-646.
- Auti AK, Wadile SC and Pawar VS (1999). Yield, quality and nutrient removal of wheat (*Triticum aestivum* L.) as influenced by levels and sources of fertilizer. Indian Journal of Agronomy, 44(1): 119-120.
- Azizi S, Thomas T and Rao S (2016). Effect of different levels of chemical fertilizers on soil physico-chemical properties of incept solutions. International Journal of Multidisciplinary Research and Development, 3(8): 29-32.
- Bakht J, Shafi M, Jan MT and Shah Z (2009). Influence of crop residue management, cropping system and N fertilizer on soil N and C dynamics and sustainable wheat (*Triticum aestivum* L.) production. Soil and Tillage Research, 104(2): 233-240.
- Bakhtiar M, Afridi MZ, Munsif F, Arshad IU, Rehman S, Khan S, Adil M, Khan MR, Khan MN and Farooq M (2017). Dry matter partitioning and grain yield of wheat as affected by sources, methods and timing of nitrogen application. Pure and Applied Biology, 6(4): 1198-1215.

- Balyan JS (1992). Effect of organic recycling on yield attributes, nitrogen concentration and uptake in wheat (*Triticum aestivum* L.). Indian Journal of Agronomy, 37(4): 701-704.
- Banglapedia MS (2014). National Encyclopedia of Bangladesh, Asiatic Society of Bangladesh.
- Baque MA, Karim MA, Hamid A and Tetsushi H (2006). Effects of fertilizer potassium on growth, yield and nutrient uptake of wheat (*Triticum aestivum* L.) under water stress condition. South Pacific Studies, 27(1): 25-35.
- BARC (2005). Bangladesh Agricultural Research Council, Fertilizer Recommendation Guide, Farmgate, Dhaka-1215, Bangladesh.
- BARC (2012). Bangladesh Agricultural Research Council, Fertilizer Recommendation Guide, Farmgate, Dhaka-1215, Bangladesh.
- BARI (2011). Bangladesh Agricultural Research Institute, Krishi Projukti Hatboi (Handbook on Agro-technology), 5th edition, Gazipur-1701, Bangladesh.
- BARI (2014). Bangladesh Agricultural Research Institute, Krishi Projukti Hatboi (Handbook on Agro-technology), 6th edition, Gazipur-1701, Bangladesh.
- Barlas NT, Conkeroglu B, Unal G and Belliturk K (2018). The effect of different vermicompost doses on wheat (*Triticum vulgaris* L.) nutrition. Journal of Tekirdag Agriculture Faculty, 15(2): 1-4.
- Basic F, Kisic I, Mesic M, Nestroy O and Butorac A (2004). Tillage and crop management effects on soil erosion in central Croatia. Soil and Tillage Research, 78: 197-206.
- BBS (2008). Bangladesh Bureau of Statistics. Statistical Year Book Bangladesh. Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka, Bangladesh.
- BBS (2017). Bangladesh Bureau of Statistics. Statistical Year Book Bangladesh. Statistics and Informatics Division, Ministry of Planning, Government of the People's Republic of Bangladesh, Dhaka, Bangladesh.

- Behera SK, Singh MV, Singh KN and Todwal S (2011). Distribution variability of total and extractable zinc in cultivated acid soils of India and their relationship with some selected soil properties. Geoderma, 162(3&4): 242-250.
- Behera UK and Pandey HN (2002). The importance of poultry manure in sustaining wheat productivity in the wheat-soybean system. Indian Farming, 51(12): 21-22.
- Belefant-Miller H (2007). Poultry litter induces tillering in rice. Journal of Sustainable Agriculture, 31(1): 151-160.
- Berg G (2009). Plant-microbe interactions promoting plant growth and health: perspectives for controlled use of microorganisms in agriculture. Applied Microbiology and Biotechnology, 84(1): 11-18.
- Beri V, Sidhu BS, Bahl GS and Bhat AK (1995). Nitrogen and phosphorus transformations as affected by crop residue management practices and their influence on crop yield. Soil Use and Management, 11(2): 51-54.
- Bharadwaj V and Omanwar PK (1994). Long term effects of continuous rotational cropping and fertilization on crop yields and soil properties-II. Effects on EC, pH, organic matter and available nutrients of soil. Journal of the Indian Society of Soil Science, 42(3): 387-392.
- Bhatt M, Singh AP, Singh V, Kala DC and Kumar V (2019). Long-term effect of organic and inorganic fertilizers on soil physico-chemical properties of a silty clay loam soil under rice-wheat cropping system in Tarai region of Uttarakhand. Journal of Pharmacognosy and Phytochemistry, 8(1): 2113-2118.
- Bhogal A, Nicholson FA, Rollett A, Taylor M, Litterick A, Whittingham MJ and Williams JR (2018). Improvements in the quality of agricultural soils following organic material additions depend on both the quantity and quality of the materials applied. Frontiers in Sustainable Food Systems, 2: 1-13.
- Bhuiya MR and Akhand MS (1983). Effect of different sources of organic materials along and in combination with two fertilizers levels on growth parameter and composition of rice. Bangladesh Journal of Agricultural Research, 7: 32-39.

- Bhuiyan MK, Ahmed I and Begum F (2007). Integrated management of seedling mortality of cauliflower (*Brassica oleraceae* var. capitata) caused by *Sclerotium rolfsii*. Journal of Agriculture, Science and Technology, 8(1&2): 79-86.
- Bista B and Dahal S (2018). Cementing the organic farming by green manures. International Journal of Applied Sciences and Biotechnology, 6(2): 87-96.
- Black CA (1965). Methods of soil analysis. American Society of Agronomy, Madison, Winscousin, USA, pp. 1-770.
- Boateng SA, Zickerman J and Kernaharens M (2006). Poultry manure effect growth and yield of maize. West African Journal of Applied Ecology, 9: 1-11.
- Bokhtiar SM, Roksana S and Moslehuddin AZM (2015). Soil fertility and productivity of sugarcane influenced by enriched pressmud compost with chemical fertilizers. SAARC Journal of Agriculture, 13(2): 183-197.
- Bolton JH, Elliot LF, Papendick RI and Bezdicek DF (1985). Soil microbial biomass and selected soil enzyme activities: Effect of fertilization and cropping practices. Soil Biology and Biochemistry, 17(3): 297-302.
- Bonde AN, Karl BG, Deshmukh MS, Tekale KU and Patil NP (2004). Effect of different organic residues on physico-chemical properties of soil in cotton-soybean intercropping in Vertisol. Journal of Soils and Crops, 14(1): 122-125.
- Bouyoucos GJ (1936). Directions for making mechanical analyses of soils by the hydrometer method. Soil Science, 42(3): 225-230.
- Brautigan DJ, Rengasamy P and Chittleborough DJ (2014). Amelioration of alkaline phytotoxicity by lowering soil pH. Crop and Pasture Science, 65(12): 1278-1287.
- Bremner JM and Mulvaney CS (1982). Nitrogen-Total *In*: Methods of Soil Analysis Part 2 (2nd Editions). American Society of Agronomy, Madison Wisconsin, USA, pp. 595-623.
- Brown S and Cotton M (2011). Changes in soil properties and carbon content following compost application: results of on-farm sampling. Compost Science and Utilization, 19(2): 87-96.

- Buarach K, Thongjoo C, Udomprasert N and Amkha S (2014). Effects of tillage system and soil organic matter amendment on growth, yield of pathumthani 80 rice and carbon sequestration in paddy soil. Modern Applied Science, 8(4): 1-7.
- Carmo DL, Lima LB and Silva CA (2016). Soil fertility and electrical conductivity affected by organic waste rates and nutrient inputs. Revista Brasileira de Ciência do Solo, 40: 1-17.
- Cassman KG, Peng S, Olk DC, Ladha K, Richardt W, Dobermann A and Singh U (1998). Opportunities for increased nitrogen use efficiency from improved resource management in irrigated rice systems. Field Crops Research, 56(1&2): 7-39.
- Celik I, Ortas I and Kilic S (2004). Effects of compost, mycorrhiza, manure and fertilizer on some physical properties of a chromoxerert soil. Soil and Tillage Research, 78(1): 59-67.
- Chaudhary S, Dheri GS and Brar BS (2017). Long-term effects of NPK fertilizers and organic manures on carbon stabilization and management index under rice-wheat cropping system. Soil and Tillage Research, 166: 59-66.
- Choudhary VP and Kumar S (2013). Maize production economics and soil productivity under different organic source of nutrients in eastern Himalayan region, India. International Journal of Plant Production, 7(2): 167-186.
- Cooperband L (2002). Building soil organic matter with organic amendments. Centre for Integrated Agricultural Systems, College of Agricultural and Life Sciences, University of Wisconsin, Madison, pp. 16.
- Dahri IA, Tagar AA, Adamowski J, Leghari N, Shah AR and Soomro SA (2018). Influence of straw incorporation-to-planting interval on soil physical properties and maize performance. International Agrophysics, 32(3): 341-347.
- Dandotiya P and Agrawal OP (2014). A simple method for improvement of soil fertility through vermicomposting. Octa Journal of Environmental Research, 2(2):139-147.
- Das M, Singh BP and Prasad RN (1992). Response of maize (*Zea mays* L.) to phosphorus enriched manures growing in P-deficient alfisols on terraced land in Meghalaya. Indian Journal of Agricultural Sciences, 61(6): 383-388.

- Dauda SN, Ajayi FA and Ndor E (2008). Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. Journal of Agriculture and Social Science, 4(3): 121-124.
- Debnath C, Kader MA and Islam N (2014). Effect of nitrogen and boron on the performance of wheat. Environmental Science and Natural Resources, 7(1): 105-110.
- Delfine S, Tognetti R, Desiderio E and Alvino A (2005). Effect of foliar application of nitrogen and humic acid on growth and yield of durum wheat. Agronomy for Sustainable Development, 25(2): 183-191.
- Deshpande H and Devasenapathy P (2010). Influence of green manure and different organic sources of nutrients on yield and soil chemical properties of rice (*Oryza sativa* L.) grown under lowland condition. International Journal of Agricultural Science, 6(2): 433-438.
- Desta HA (2015). Effects of organic and inorganic fertilizers on selected soil properties after harvesting maize at Antra Catchment, Northwestern Ethiopia. International Invention Journal of Agricultural and Soil Science, 3(5): 68-78.
- Dhar D, Datta A, Basak N, Paul N, Badole S and Thomas T (2014). Residual effect of crop residues on growth, yield attributes and soil properties of wheat under rice-wheat cropping system. Indian Journal of Agricultural Research, 48(5): 373-378.
- Diacono M and Montemurro F (2009). Long-term effects of organic amendments on soil fertility: A review. Agronomy for Sustainable Development, 30: 401-422.
- Dominguez J, Edwards CA and Subler S (1997). A comparison of vermicomposting and composting. Biocycle, 38: 57-59.
- Doni F, Isahak A, Zain CRCM and Yusoff WMW (2014). Physiological and growth response of rice plants (*Oryza sativa* L.) to *Trichoderma* spp. inoculants. AMB Express, 4(45): 1-7.
- Dubey L, Dubey M and Jain P (2015). Role of green manure in organic farming. Plant Archives, 15(1): 23-26.

- Dubey V, Patel AK, Shukla A, Shukla S and Singh S (2012). Impact of continuous use of chemical fertilizer. International Journal of Engineering Research and Development, 3(11): 13-16.
- Eck HV and Stewart BA (1995). Manure in soil amendments and environmental quality. Lewis publication, Boca Raton, Florida, pp. 169-98.
- Eckhardt DP, Redin M, Santana NA, Conti LD, Dominguez J, Jacques RJ and Antoniolli ZI (2018). Cattle manure bioconversion effect on the availability of nitrogen, phosphorus, and potassium in soil. Revista Brasileira de Ciência do Solo, 42: 1-10.
- Eghball B, Ginting D and Gilley JE (2004). Residual effects of manure and compost applications on corn production and soil properties. American Society of Agronomy, 96(2): 442-447.
- Elbaum R, Zaltzman L, Burgert I and Fratzl, P (2007). The role of wheat awns in the seed dispersal unit. Science, 316(5826): 884-886.
- Enujeke EC (2013). Effects of poultry manure on growth and yield of improved maize in asaba area of delta state, Nigeria. Journal of Agriculture and Veterinary Science, 4(5): 24-30.
- Eriksen J (2009). Soil sulfur cycling in temperate agricultural systems. Advances in Agronomy, 102: 55-89.
- FAO (Food and Agriculture Organization) and UNDP (United Nations Development Program) (1994). Land Degradation in South Asia: It's Severity Causes and Efforts upon the People, FAO World Resources Report No. 78.
- Farhad W, Saleem MF, Cheema MA and Hammad HM (2009). Effect of poultry manure levels on the productivity of spring maize (*Zea mays* L). The Journal of Animal and Plant Sciences, 19(3): 122-125.
- Farhad W, Saleem MF, Cheema MA, Khan HZ and Hammad HM (2011). Influence of poultry manure on the yield and quality of spring maize. Crop Environment, 2: 6-10.

- Farooq N, Iqbal M, Zahir ZA and Farooq M (2018). Integration of allelopathic crop residues and NPK fertilizer to mitigate residue-phytotoxicity, improve soil fertility and wheat growth under different moisture conditions. Planta Daninha, 36: 1-11.
- Forster S, Welp G and Scherer HW (2012). Sulfur specification in bulk soil as influenced by long-term application of mineral and organic fertilizers. Plant, Soil & Environment, 58(7): 316-321.
- Frederick JR, Bauer PJ and Busscher WJ (2001). Grain yield and yield components of double cropped winter wheat as affected by wheat and previous soybean production practices. Crop Science, 41: 778-784.
- Gaind S and Nain L (2007). Chemical and biological properties of wheat soil in response to paddy straw incorporation and it's biodegradation by fungal inoculants. Biodegradation, 18(4): 495-503.
- Gao M, Lu X, Huang Y, Liu N and Yang J (2017). Impact of long-term application fertilizer on soil total sulphur and valid sulphur. *In*: IOP Conference Series: Materials Science and Engineering, 207(1): 1-6.
- Garcia MCV, Estrella FS, Lopez MJ and Morino J (2008). Influence of compost amendment on soil biological properties and plants. Dynamic Soil, Dynamic Plant, 2(1): 1-9.
- Garg S and Bahla GS (2008). Phosphorus availability to maize as influenced by organic manure and fertilizer P associated phosphorus activity in soil. Bioresource Technology, 99(13): 5773-5777.
- Geng Y, Cao G, Wang L and Wang S (2019). Effects of equal chemical fertilizer substitutions with organic manure on yield, dry matter, and nitrogen uptake of spring maize and soil nitrogen distribution. PLoS One, 14(7): 1-16.
- Ghassan JZ, Zakaria W, Shaari AR and Mohammud CH (2017). Mung bean residue incorporated at different incubation periods affect soil chemical properties. Journal of Agricultural and Biological Science, 12(7): 230-235.

- Ghosh AB, Bajaj JC, Hasan R and Singh D (1983). Soil and Water Testing Methods. A Laboratory Manual, Division of Soil Science and Agricultural Chemistry. IARI, New Delhi, pp. 36-45.
- Gil MV, Carballo MT and Calvo LF (2007). Fertilization of maize with compost from cattle manure supplemented with additional mineral nutrients. Waste Management, 28(8): 1432-1440.
- Girma K, Mosali J, Freeman KW, Raun WR, Martin KL and Thomason WE (2005). Forage and grain yield response to applied sulfur in winter wheat as influenced by source and rate. Journal of Plant Nutrition, 28(9): 1541-1553.
- Gomez KA and Gomez AA (1984). Statistical Procedures for Agricultural Research (2nd edition). International Rice Research Institute. John Wiley and Sons, a Wiley-Inter Science Publication, pp. 1-680.
- Gopinath KA, Saha S, Mina BL, Pande H, Kundu S and Gupta HS (2008). Influence of organic amendments on growth, yield and quality of wheat and on soil properties during transition to organic production. Nutrient Cycling in Agroecosystems, 82(1): 51-60.
- Gowda C, Biradar Patil NK, Patil BN, Awakknavar JS and Hunje R (2010). Effect of organic manures on growth, seed yield and quality of wheat. Karnataka Journal of Agricultural Sciences, 21(3): 366-368.
- Guong VT, Hien NX and Minh D (2010). Effect of fresh and composted organic amendment on soil compaction and soil biochemical properties of citrus orchards in the Mekong Delta, Vietnam. 19th World Congress of Soil Science, Soil Solutions for a Changing World, 1-6 August 2010, Brisbane, Australia, pp. 76-78.
- Gurpreet S, Jalota, SK and Yadvinder S (2007). Manure and residue management effects on physical properties of a soil under the rice-wheat system in Punjab, India. Soil and Tillage Research, 94(1): 229-238.
- Guy C, Haskell D, Neven L, Klein P and Smelser C (1992). Hydration-state-responsive proteins link cold and drought stress in spinach. Planta, 188(2): 265-270.

- Ha TN (2010). Using *Trichoderma* species for biological control of plant pathogens in Viet Nam. International Society for Southeast Asian Agricultural Sciences, 16(1): 17-21.
- Hachicha S, Chtourou M, Medhioub K and Ammar E (2006). Compost of poultry manure and olive mill wastes as an alternative fertilizer. Agronomy for Sustainable Development, 26(2): 135-142.
- Hajieghrari B and Mohammadi M (2016). Growth-promoting activity of indigenous *Trichoderma* isolates on wheat seed germination, seedling growth and yield. Australian Journal of Crop Science, 10(9): 1339-1347.
- Hammad HM, Khaliq A, Ahmad A, Aslam M, Malik AH, Farhad W and Laghari KQ (2011). Influence of different organic manures on wheat productivity. International Journal of Agriculture and Biology, 13(1): 137-140.
- Harris F (2002). Management of manure in farming systems in semi-arid West Africa. Experimental Agriculture, 38(2): 131-148.
- Hassan EA (2002). Effect of chicken manure and season on the performance and HCN Content of two forage sorghum cultivar. PhD Thesis, Faculty of Agriculture, University of Khartoum, Sudan.
- Hermosa R, Viterbo A, Chet I and Monte E (2012). Plant-beneficial effects of *Trichoderma* and of its genes. Microbiology, 158(1): 17-25.
- Hossaen MA, Shamsuddoha ATM, Paul AK, Bhuiyan MSI and Zobaer ASM (2011). Efficacy of different organic manures and inorganic fertilizer on the yield and yield attributes of *Boro* rice. The Agriculturists, 9(1&2): 117-125.
- Hossain MM, Hossain I and Khalequzzaman KM (2016). Biological control of leaf blight of wheat caused by *Bipolaris sorokinian*. Bulletin of the Institute of Tropical Agriculture, Kyushu University, 39(1): 43-51.
- Hossain S (2011). Effect of cow dung and nitrogen (urea) on the growth, yield and yield contributing characters of aromatic rice cv. BRRI *Dhan* 50. MS Thesis, Department of Agronomy, Bangladesh Agricultural University, Mymensingh.

- Huang GF, Fang M, Wu QT, Zhou LX, Liao XD and Wong JW (2001). Co-composting of pig manure with leaves. Environmental Technology, 22(10): 1203-1212.
- Hue NV and Silva JA (2000). Organic soil amendments for sustainable agriculture: organic sources of nitrogen, phosphorus, and potassium. Plant nutrient management in Hawaii's soils, approaches for tropical and subtropical agriculture. College of Tropical Agriculture and Human Resources, University of Hawaii, Manoa, pp. 133-144.
- Hussain T (2001). Manures and organic wastes. *In*: Soil Science (2nd Ed.), National Book Foundation, Islamabad, Pakistan, pp. 387-403.
- Ibrahim M, Hassan A, Iqbal M and Valeem EE (2008). Response of wheat growth and yield to various levels of compost and organic manure. Pakistan Journal of Botany, 40(5): 2135-2141.
- Imran NJ, Ashfaq A and Asad AK (2017). Grain yield, yield attributes of wheat and soil physico chemical characteristics influenced by biochar, compost and inorganic fertilizer application. Agricultural Research and Technology: 10(4): 1-6.
- Iqbal J, Hayat K and Hussain S (2012). Effect of seeding rates and nitrogen levels on yield and yield components of wheat (*Triticum aestivum* L.). Pakistan Journal of Nutrition, 11(7): 531-536
- Iqbal T, Jilani G, Chaudhry AN and Zahid A (2008). Studies on the residual effect of poultry litter application under wheat-maize cropping system. Presented and abstracted in the 12th Congress of Soil Science Society of Pakistan, pp. 20-23.
- Islam MR, Shaikh MS, Siddique AB and Sumon MH (2014). Yield and nutrient uptake by wheat as influenced by integrated use of manures and fertilizers. Journal of the Bangladesh Agricultural University, 12(1): 73-78.
- Islam MS (1990). Soil characteristics and their effects on sulfur and zinc deficiency problems in Bangladesh. *In*: a workshop on sulfur and zinc management of soils and crops and balanced fertilization in Bangladesh, 10-11 April 1990, Dhaka, Bangladesh.

- Islam MS, Amin MS and Anwar MN (1994). Integrated soil fertility management in Bangladesh. *In*: Workshop on Integrated Nutrient Management for Sustainable Agriculture, 26 June 1994, SRDI, Dhaka, Bangladesh.
- Islam MS, Hasanuzzamanb M, Rokonuzzamanc M and Nahard K (2009). Effect of split application of nitrogen fertilizer on morpho-physiological parameters of rice genotypes. International Journal of Plant Production, 3(1): 51-61.
- Ismaeil FM, Abusuwar AO and El Naim AM (2012). Influence of chicken manure on growth and yield of forage sorghum (*Sorghum bicolor* L. Moench). International Journal of Agriculture and Forestry, 2(2): 56-60.
- ISTA (1985). International Seed Testing Association International rules for seed testing. Seed Science and Technology, 13(2): 299-513.
- Jackson ML (1962). Hydrogen activity determination for soil. Soil chemical analysis. Constable and Company Limited, London, pp. 38-56.
- Jackson ML (1973). Soil chemical analysis Prentice-Hall of India private limited M-97. New Delhi, India, pp. 498.
- Jahiruddin M and Satter MA (2010). Agricultural research priority: Vision 2030 and beyond. Sub-sector: land and soil resource management, Bangladesh. Final Report, pp. 1-47.
- Jan A, Daur I, Muhammad Z and Khan IA (2011). Effect of mung bean residue and nitrogen levels on barley. Pakistan Journal of Botany, 43(4): 1905-1908.
- Jat NK, Kumar A, Meena SR, Rana DS, Meena BP and Rana KS (2012). Influence of integrated nutrient management on the productivity, quality and soil health of maize (*Zea mays*) wheat (*Triticum aestivum*) cropping system. Indian Journal of Agronomy, 57(4): 327-332.
- Kabesh MO, El-Kramany MF, Sary GA, El-Naggar HM and Gehan SH (2009). Effects of sowing methods and some bio-organic fertilization treatments on yield and yield components of wheat. Research Journal of Agriculture and Biological Sciences, 5(1): 97-102.

- Kareem I, Jawando OB, Eifediyi EK, Bello WB and Oladosu Y (2017). Improvement of growth and yield of maize (*Zea mays* L.) by poultry manure, maize variety and plant population. Agronomical Research in Moldavia, 50(4): 51-64.
- Karim Z and Iqbal HA (2001). Impact of land degradation in Bangladesh. Changing scenario in agricultural land use. Bangladesh Agricultural Research Council, Farmgate, Dhaka-1215, Bangladesh.
- Khaliq T, Mahmood T and Masood A (2004). Effectiveness of farmyard manure, poultry manure and nitrogen for corn (*Zea mays*) productivity. International Journal of Agriculture and Biology, 6(2): 260-263.
- Khan MA, Shukla AK, Upadhyay NC and Singh OP (2005). Nutrient management strategies for maize-potato-wheat cropping system in typic ustochrept soils. Potato Journal, 32: 167-168.
- Khan TU, Jan MT, Khan A, Ahmad G, Ishaq M, Afridi K, Murad A, Qureshi MA, Ahmad I and Seed M (2018). Integrated management of fertilizer nitrogen and poultry manure enhances wheat production. Pakistan Journal of Agricultural Research, 31(3): 207-215.
- Khanam M, Rahman MH and Islam MR (2001). Effect of manure and fertilizers on the growth and yield of BRRI *Dhan* 30. Pakistan Journal of Biological Sciences, 4(2): 172-174.
- Kharub AS, Sharma RK, Mongia AD, Chhokar RS, Tripathi SC and Sharma VK (2004).
 Effect of rice (*Oryza sativa*) straw removal, burning and incorporation on soil properties and crop productivity under rice-wheat (*Triticum aestivum*) system.
 Indian Journal of Agricultural Science, 74(6): 295-299.
- Khatun, Khaleda MST, Hasan MK, Marina SR, Sabagh AE and Islam MS (2018). Response of growth and yield attributes of aromatic rice to cow dung and zinc fertilization. Advances in Plants and Agricultural Research, 8(6): 465-470.
- Kirchmann H and Gerzabek MH (1999). Relationship between soil organic matter and micropores in a long-term experiment at Ultuna, Sweden. Journal of Plant Nutrition and Soil Science, 162(5): 493-498.

- Kirchner MJ, Wollum II AG and King LD (1993). Soil microbial populations and activities in reduced chemical input agroecosystems soils. Soil Science Society of American Journal, 57(5): 1289-1295.
- Kowsar J, Boswal MV and Gunjan S (2015). Impact of bio-fertilizer and organic fertilizer on yield and yield parameters of bread wheat (*Triticum aestivum* L.). International Journal of Current Research, 7(2): 12388-12395.
- Krishna KR (2013). Agroecosystems: soils, climate, crops nutrients dynamics and productivity. Apple Academic Press Incorporated, Toronto, Canada, pp. 1-517.
- Krishnan P, Nagarajan S, Dadlani M and. Moharir AV (2003). Characterization of wheat (*Triticum aestivum* L.) and soybean (*Glycine max* L.) seeds under accelerated aging conditions by proton nuclear magnetic spectroscopy. Seed Science and Technology, 31(3): 541-550.
- Kumar K and Goh KM (2000). Crop residues and management practices: effects on soil quality, soil nitrogen dynamics, crop yields, and nitrogen recovery. Advances in Agronomy, 68: 197-319.
- Kumar S and Abraham T (2018). Productivity potential of wheat under certified organic production system. International Journal of Current Microbiology and Applied Sciences, 7(10): 281-288.
- Kuo S, Sainju UM and Jellum EJ (1997). Winter cover crop effects on soil organic carbon and carbohydrate in soil. Soil Science Society of American Journal, 61(1): 145-152.
- Ladha JK, Khind CS, Gupta RK, Meelu OP and Pasuquin E (2004). Long-term effects of organic inputs on yield and soil fertility in the rice-wheat rotation. Soil Science Society of America Journal, 68(3): 845-853.
- Lal R (2005). World crop residues production and implications of its use as a biofuel. Environment International, 31(4): 575-584.

- Lasaridi KE and Stentiford EI (1999). Composting of source separated MSW: an approach to respirometric techniques and biodegradation kinetics *In*: International Symposium on Composting of Organic Matter 549, 30 August 1999, Kassandra-Chalkidiki, Greece, pp. 81-88.
- Li D, Luo P, Han X and Yang J (2018). Influence of long term fertilization on soil physicochemical properties in a brown soil. IOP Conference Series: Earth and Environmental Science, 108: 1-6.
- Li JT, Zhong XL, Wang F and Zhao QG (2011). Effect of poultry litter and livestock manure on soil physical and biological indicators in a rice-wheat rotation system. Plant Soil Environment, 57(8): 351-356.
- Liebig MA, Varvel GE, Doran JW and Wienhold BJ (2002). Crop sequence and nitrogen fertilization effects on soil properties in the Western Corn Belt. Soil Science Society of American Journal, 66(2): 596-601.
- Liu JR and Shen RP (1992). The effects of fertilization on the properties and role of soil organic matter. *In*: Proceedings of the International Symposium on Paddy Soils, Nanjing, China, pp. 246-251.
- Ma BL, Dwyer LM and Gregorich EG (1999). Soil nitrogen amendment effects on nitrogen uptake and grain yield of maize. Agronomy Journal, 91(4): 650-656.
- Maguire JD (1962). Speed of germination-aid in selection and evaluation for seedling emergence and vigor. Crop Science, 2(2): 176-177.
- Mahato S, Bhuju S and Shrestha J (2018). Effect of *Trichoderma viride* as bio-fertilizer on growth and yield of wheat. Malaysian Journal of Sustainable Agriculture, 2(2): 1-5.
- Mahmood F, Khan I, Ashraf U, Shahzad T, Hussain S, Shahid M, Abid M and Ullah S (2017). Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. Journal of Soil Science and Plant Nutrition, 17(1): 22-32.
- Makinde EA (2007). Effects of an organo-mineral fertilizer application on the growth and yield of maize. Journal of Applied Sciences Research, 3(10): 1152-1155.

- Mandal KG, Misra AK, Hati KM, Bandyopadhyay KK, Ghosh PK and Mohanty M (2004).
 Rice residue-management options and effects on soil properties and crop productiviy.
 Journal of Food Agriculture and Environment, 2(1): 224-231.
- Mandal UK, Singh G, Victor US and Sharma KL (2003). Green manure: its effect on soil properties and crop growth under rice-wheat cropping system. European Journal of Agronomy, 19(2): 225-237.
- Marinari S, Masciandaro G, Ceccanti B and Grego S (2000). Influence of organic and mineral fertilizers on soil biological and physical characteristics. Bioresource Technology, 72(1): 9-17.
- Marshall TJ (1947). Mechanical composition of soils in relation to field description of texture. Council for Scientific and Industrial Research, Australia, Bulletin No. 224.
- Matsi T, Lithourgidis AS and Gagianas AA (2003). Effects of injected liquid cattle manure on growth and yield of winter wheat and soil characteristics. Agronomy Journal, 95(3): 592-596.
- Mbah CN and Onweremadu EU (2009). Effect of organic and mineral fertilizer inputs on soil and maize grain yield in an acid ultisol in Abakaliki-South Eastern Nigeria. American-Eurasian Journal of Agronomy, 2(1): 7-12.
- Meelu OP (1994). Recycling of crop residues and green manure for soil and crop productivity improvement in rice-wheat cropping system. Temperate rice: achievements and potential 2, pp. 605-613.
- Meena HN, Rana KS, Kumar A, Pooniya V, Shukla L and Biswas DR (2018). Productivity of wheat as influenced by crop residue and potassium management in maize (*Zea mays* L.)-wheat (*Triticum aestivum* L.)-mung bean cropping system. International Journal of Current Microbiology and Applied Sciences, 7(2): 1775-1785.
- Metho LA, Hammes PS and Beyers EA (1998). The effect of soil fertility on the contribution of main stem, tillers and kernel position to grain yield and grain protein content of wheat. South African Journal of Plant and Soil, 15(2): 53-60.

- Mia MMU and Karim Z (1995). Extension of Integrated Plant Nutrient System (IPNS) at farm level in Bangladesh: Progress and problems in the extension of integrated plant nutrient systems at farm level in Asia. FAO, RAPA Publication 12, pp. 29-42.
- Mia MMU, Habibullah AKM and Ali MF (1993). Depletion of organic matter in upland soils of Bangladesh. *In*: International Seminar on Managing Red and Lateric Soils for Sustainable Agriculture, 24-28 September 1993, Bangalore, India.
- Mihov M and Tringovska I (2010). Energy efficiency improvement of greenhouse tomato production by applying new biofertilizers. Bulgarian Journal of Agricultural Science, 16(4): 454-458.
- Mitchell CC and Tu S (2005). Long-term evaluation of poultry litter as a source of nitrogen for cotton and corn. Agronomy Journal, 97(2): 399-407.
- Muhammad FJ, Muhammad DA, Waqas L, Haseeb A and Wazir R (2018). Effect of poultry manure and phosphorous on phenology, yield and yield components of wheat. International Journal of Current Microbiology and Applied Sciences, 7(5): 3751-3760.
- Mukhtiar A, Waqar A, Khalil MK, Tariq M, Muhammad S, Hussain A and Kamal A (2018). Evaluating the potential organic manure for improving wheat yield and quality under agro-climatic conditions of Pakistan. Advances in Crop Science and Technology, 6(2): 1-4.
- Muller MM and Veronica S (1988). The fate of nitrogen (15N) released from different plant materials during decomposition under field conditions. Plant Soil, 105: 133-139.
- Munawar A, Blevins RL, Frye WW and Saul MR (1990). Tillage and cover crop management for soil water conservation. Agronomy Journal, 82(4): 773-777.
- Musa M, Bokhtiar SM and Gurung TR (2015). Status and future prospect of organic agriculture for safe food security in SAARC countries, pp. 1-19.
- Naeem M, Khan F and Ahmad W (2009). Effect of farmyard manure, mineral fertilizers and Mung bean residue on some microbiological properties of eroded soil in district Swat. Soil & Environment, 28(2): 162-168.

- Nahar MS, Rahman MA and Kibria MG (2012). Use of tricho-compost and tricho-leachate for management of soil borne pathogen production of healthy cabbage seedlings. Bangladesh Journal of Agricultural Research, 37(4): 653-664.
- Naseri R, Mirzaei A, Soleimani R and Nazarbeygi E (2010). Response of bread wheat to nitrogen application in calcareous soils of western Iran. American-Eurasian Journal Agricultural and Environmental Sciences, 9(1): 79-85.
- Nguyen ML, Haynes RJ and Goh KM (1995). Nutrient budgets and status in three pairs of conventional and alternative and mixed cropping farms in Canterbury, New Zealand. Agriculture, Ecosystems & Environment, 52(2&3): 149-162.
- Nimmo JR (2004). Porosity and pore size distribution. Encyclopedia of Soils in the Environment, 3(1): 295-303.
- Nyborg M, Solberg ED, Malhi SS and Izaurralde RC (1995). Fertilizer N, crop residue, and tillage alter soil C and N content in a decade. Advances in Soil Science: 93-99.
- OFRD (1998). Annual report: On Farm Research Division (Bangladesh Agricultural Research Institute). On-Farm soil fertility management. Progressive Agriculture, 11(1&2): 77-80.
- Ogunbanjo OR, Omueti JA, Jayeoba OJ, Omolaiye JA and Uzokwe NE (2010). Yield response of maize to animal manures and bulking agents in fortified composts. ASSET: An International Journal (Series A), 7(1): 87-98.
- Ogundare K, Agele S and Aiyelari P (2012). Organic amendment of an ultisol: effects on soil properties, growth, and yield of maize in Southern Guinea savanna zone of Nigeria. International Journal of Recycling of Organic Waste in Agriculture, 1(1): 1-7.
- Olsen SR and Sommers LE (1982). Phosphorus *In*: Methods of Soil Analysis Part 2 (2nd Editions). American Society of Agronomy, Madison Wisconsin, USA, pp. 403-427.
- Omara P, Aula L and Raun WR (2019). Nitrogen uptake efficiency and total soil nitrogen accumulation in long-term beef manure and inorganic fertilizer application. International Journal of Agronomy, 2019: 1-6. https://doi.org/10.1155/2019/9594369.

- Ouedraogo E, Mandob A and Zombrec NP (2001). Use of compost to improve soil properties and crop productivity under low input agricultural system in West Africa. Agriculture, Ecosystems & Environment, 84(3): 259-266.
- Page AL, Miller RH and Keeney DR (1982). Methods of Soil Analysis Part 2 (2nd Editions). American Society of Agronomy, Madison Wisconsin, USA.
- Page-Dumroese D, Harvey A, Jurgensen M and Graham R (1991). Organic matter function in the Western-Montane forest soil system. Symposium on: Proceedings-Management and productivity of Western-Montane forest soils, 10-12 April 1990. USDA, General Technical Report INT-280: 95-100.
- Park HM, Kim WC and Ahn SB (1995). The effect of long term application of compost and lime on rice yield and physic-chemical properties of paddy soil. Australian Journal of Soil Science Research, 36(2): 247-256.
- Parvez MS, Islam MR, Begum MS, Rahman MS and Abedin Miah MJ (2008). Integrated use of manure and fertilizers for maximizing the yield of BRRI *Dhan* 30. Journal of the Bangladesh Society for Agricultural Science and Technology, 5(1&2): 257-260.
- Patel NM, Sadaria SG, Kaneria BB and Khanpara VD (1995). Effect of nitrogen, potassium and zinc on growth and yield of wheat (*Triticum aestivum* L.) Indian Journal of Agronomy, 38(1): 113-115.
- Patra DD, Bhandari JC and Misra A (1992). Effect of plant residue on the size of microbial biomass and nitrogen mineralization in soil: incorporation of cowpea and wheat straw. Soil Science and Plant Nutrition, 38(1): 1-6.
- Pattanayak SK, Mishra KN, Jena MK and Nayak RK (2001). Evaluation of green manure crops fertilized with various phosphorus sources and their effect on subsequent rice crop. Journal of the Indian Society of Soil Science, 49(2): 285-291.
- Paul J, Choudhary AK, Suri VK, Sharma AK, Kumar V and Shobhna (2014). Bio resource nutrient recycling and its relationship with bio-fertility indicators of soil health and nutrient dynamics in rice–wheat cropping system. Communications in Soil Science and Plant Analysis, 45(7): 912-924.

- Paulsen HM (2005). Sulfur in organic farming. *In*: Aspects of sulfur nutrition of plants; evaluation of China's current, future and available resources to correct plant nutrient sulfur deficiencies–report of the first Sino-German Sulfur workshop, pp. 105.
- Pawar SK, Kumbhar GA and Sangale PN (2018). Effect of crop residues and green manure on soil properties and yield of cotton in salt affected soils of purna valley. Pharmacognosy and Phytochemistry, 7(3): 494-498.
- Peerzada OH, Mor VS, Abhinav D, Axay B, Dahiya OS, Pandey V, Anzer UI and Mohammad S (2016). Influence of integrated nutrient management on seed quality of fenugreek (*Rigonellafoenum graecum* L.). Environment and Ecology, 34(4): 2226-2230.
- Peltonen-sainio P, Kangas A, Salo Y and Jauhiainen L (2007). Grain numbers dominate grain weight in temperate cereal yield determination: evidence based on 30 years of multi-location trials. Field Crops Research, 100(2&3): 179-188.
- Phullan NK, Memon M, Shah JA, Memon MY, Sial TA, Talpur NA and Khushk GM (2017). Effect of organic manure and mineral fertilizers on wheat growth and soil properties. Basic and Applied Sciences, 13: 559-565.
- Polthanee A, Promkhambut A and Kaewrahan S (2011). Growth and yield of organic rice as affected by rice straw and organic fertilizer. International Journal of Environment and Rural Development, 2(1): 93-99.
- Polthanee A, Promkhambut A and Trelo-Ges V (2012). Effect of pre-rice mung bean and cattle manure application on growth and yield of organic rice. International Journal of Environment and Rural Development, 3(1): 1-9.
- Pradeep P (2018). Seed quality parameters (Germination percentage and seedling vigor index) of *Rabi* sorghum seeds influenced by rice weevil infestation. MedCrave Online Journal of Toxicology, 4(6): 391-396.
- Pradhan SB (1992). Status of fertilizer use in developing countries of Asia and the Pacific Region *In*: Procedure Regression FADINAP Seminar. Chang Mai, Thailand, pp. 37-47.

- Prasad BK, Singh DN and Singh SN (2003). Effect of long-term use of fertilizer, lime and manures on growth and yield of sweet corn. Journal of Indian Society Soil Science, 34: 271-274.
- Prasad NK and Kerketta R (1991). Soil fertility as influenced by different cropping sequences. Indian Agricultural Science, 61: 16-19.
- Prasad R and Power JF (1991). Crop residue management. Advances in Soil Science, 15: 205-239.
- Pratibha S, Amar NP, Mahesh KS and Swati D (2012). Field demonstration of *Trichoderma harzianum* as a plant growth promoter in wheat (*Triticum aestivum* L). Journal of Agricultural Sciences, 4(8): 65-73.
- Radford PJ (1967). Growth analysis formulae-their use and abuse. Crop Science, 7(3): 171-175.
- Rahman MA (2001). Integrated use of fertilizer and manure for crop production in wheatrice cropping pattern. PhD Thesis, Department of Soil Science, Bangladesh Agricultural University, Mymensingh, pp. 117-118.
- Rahman MA (2004). Effect of irrigation on nitrogen uptake, yield and yield attributes of wheat varieties. PhD Thesis, Crop Physiology Laboratory Department of Botany, Rajshahi University, Rajshahi.
- Rahman MA, Paul NK and Sarker MAR (2006). Study on growth and yield of wheat with irrigation and split application of nitrogen. Progressive Agriculture, 17(1): 1-7.
- Raissi A, Galavi M, Ramroudi M, Mousavi SR and Rasoulizadeh MN (2012). Effects of phosphate bio-fertilizer, organic manure and chemical fertilizers on yield, yield components and seed capabilities of isabgol (*Plantago ovata*). International Journal of Agriculture and Crop Sciences, 24: 1821-1826.
- Raj A, Jhariya MK and Toppo P (2014). Cow dung for ecofriendly and sustainable productive farming. International Journal of Scientific Research, 4(10): 201-202.
- Rajinder K, Gurkirpal, Bandari AI, Walia SS, Kumar R and Singh G (2000). Effect of manures and fertilizers on grain yield and changes in soil micronutrients under rice-wheat cropping system. Environment and Ecology, 18(1): 634-638.

- Rajput MKK, Ansari AH, Mehdi S and Hussai AM (1988). Effect of N and P fertilizers alone and in combination with organic matter on the growth and yield of Toria. Sarhad Journal of Agriculture, 4: 3-6.
- Ram M (2017). Effective use of cow dung manure for healthy plant growth. International Journal of Advanced Research and Development, 2(5): 218-221.
- Ramesh P, Panwar NR, Singh AB, Ramana S and Rao AS (2009). Impact of organicmanure combinations on the productivity and soil quality in different cropping systems in central India. Plant Nutrition and Soil Science, 172(4): 577-585.
- Rasoulzadeh A and Yaghoubi A (2010). Effect of cattle manure on soil physical properties on a sandy clay loam soil in North-West Iran. Journal of Food, Agriculture and Environment, 8(2): 976-979.
- Rasul GAM, Ahmed ST and Ahmed MQ (2015). Influence of different organic fertilizers on growth and yield of wheat. American-Eurasian Journal of Agriculture and Environmental Sciences, 15(6): 1123-1126.
- Rautaray SK, Ghosh BC and Mittra BN (2003). Efficacy of organic material on growth and yield of rice as influenced by time of application under integrated nutrient management. Oryza, 40(1&2): 18-21.
- Reddy DD, Rao AS and Rupa TR (2000). Effects of continuous use of cattle manure and fertilizer phosphorus on crop yields and soil organic phosphorus in a Vertisol. Bioresource Technology, 75(2): 113-118.
- Rees R and Castle K (2002). Nitrogen recovery in soils amended with organic manures combined with inorganic fertilizers. Agronomie, 22: 739-746.
- Rehman S, Khalil SK, Muhammad F, Rehman A, Khan AZ, Saljoki AR, Muhammad Z and Khalil IH (2010). Phenology, leaf area index and grain yield of rainfed wheat influenced by organic and inorganic fertilizer. Pakistan Journal of Botany, 42(5): 3671-3685.
- Rijmpa J and Jahiruddin M (2004). National strategy and plan for use of soil nutrient balance in Bangladesh: A Consultancy Report. SFFP, Khamarbari, Dhaka, Bangladesh.

- Rosales CM, Rivera FG, Hautea RA and Rosario ED (1998). Field evaluation of N_2 fixation by mung bean in the Philippines and residual effects on maize. *In*: Improving yield and nitrogen fixation of grain legumes in the tropics and subtropics of Asia, pp. 65-75.
- Ru SH, Zhang GY, Niu JF, Sun SY, Wang L and Geng N (2012). Effects of chicken manure application on crop yields and soil nutrients accumulation in wheat-corn rotation system. International Conference on Biomedical Engineering and Biotechnology, 28 May 2012, pp. 1392-1396.
- Sabaghina N, Janmohammadi M and Segherloo A (2014). Evaluation of some agromorphological traits diversity in Iranian bread wheat genotypes. UMCS-Biologia, 19(1): 79-90.
- Saddiq MH and Al-Ameer HA (2011). The effect of rice straw and poultry waste addition on the soil physical properties I- clay soil. Researches of the first international conference, University of Babylon, Iraq, pp. 107-114.
- Saini JP and Kumar R (2014) Long term effect of organic sources of nutrients on productivity and soil health in maize + soybean-wheat + gram cropping system.
 'Building Organic Bridges' at the organic world congress 13-15 October 2014, Istanbul, Turkey.
- Samra JS, Singh B and Kumar K (2003). Managing crop residues in the rice-wheat system of the Indo-Gangetic Plain. Improving the productivity and sustainability of rice-wheat systems: Issues and Impacts, 65: 173-195.
- Santillan YM, Moreno FP and Garcia FP (2014). Effect of the application of manure of cattle on the properties chemistry of soil in Tizayuca, Hidalgo, Mexico. International Journal of Applied Science and Technology, 4(3): 67-72.
- Sarkar MK, Acharya A and Sengupta PK (2002). Management of sheath blight of rice caused by *Rhizoctonia solani* by oilcake amendment of soil. Journal of Mycopathological Research, 40(2): 153-158.

- Sarwar G, Hussain N, Mujeeb F, Schmeisky H and Hassan G (2003). Biocompost application for the improvement of soil characteristics and dry matter yield of *Lolium perenne* (Grass). Asian Journal of Plant Science, 2(2): 237-241.
- Sarwar G, Hussain N, Schmeisky H, Muhammad S, Ibrahim M and Safdar E (2007). Use of compost an environment friendly technology for enhancing rice-wheat production in Pakistan. Pakistan Journal of Botany, 40(1): 1553-1558.
- Sarwar G, Hussain N, Schmeisky H, Suhammad S, Ibrahim M and Ahmad S (2008). Efficiency of various organic residues for enhancing rice-wheat production under normal soil condition. Pakistan Journal of Botany, 40(5): 2107-2113.
- Saunders DA (1991) Report of on farm survey: Jessore and Kushtia District: Farmers practices and problems and their Implications. *Monograph* No: 8, Wheat Research Center (BARI) Nasipur, Dinajpur, Bangladesh.
- Schaefer GL, Cosh MH and Jackson TJ (2007). The USDA natural resources conservation service Soil Climate Analysis Network (SCAN). Journal of Atmospheric and Oceanic Technology, 24(12): 2073-2077.
- Sen H and Jana PK (1998). Effect of soil moisture tensions and predicting winter crops on the yield, water use efficiency and moisture extraction pattern of succeeding pre-*Kharif* direct seed rice. Indian Agriculturist, 31: 199-205.
- Seufert V, Ramankutty N and Foley JA (2012). Comparing the yields of organic and conventional agriculture. Nature, 485(7397): 229-232.
- Sevaram, Sharma GC, Khybri ML and Gupta OP (1998). On sustainability in yield of maize wheat cropping system under integrated nutrient management. Journal of Andaman Science Association, 14: 21-24.
- Shah SA, Mohammad W, Haroon and Anwar A (2017). Residual effect of poultry manure and mineral N application on maize production under wheat-maize cropping system.
 Turkish Journal of Agriculture-Food Science and Technology, 5(9): 1061-1065.
- Shah SA, Shah SM, Mohammad W, Shafi M, Nawaz H, Shehzadi S and Amir M (2010). Effect of integrated use of organic and inorganic nitrogen sources on wheat yield. Sarhad Journal of Agriculture, 26(4): 559-563.

- Shahid M, Saleem MF, Khan HZ, Wahid MA and Sarwar M (2015). Improving wheat (*Triticum aestivum* L.) yield and quality by integration of urea with poultry manure. Soil & Environment, 34(2): 148-155.
- Shahrzad K, Mahmoud K, Hossein AK, Mehran H and Majid A (2014). Effects of incorporation of crops residue into soil on some chemical properties of soil and bioavailability of copper in soil. International Journal of Advanced Biological and Biomedical Research, 2(11): 2819-2824.
- Shanggan ZP, Shao A and Dychmans J (2000). Nitrogen nutrition and water stress effects on leaf photosynthetic gas exchange and water use efficiency in winter wheat. Environmental and Experimental Botany, 44(2): 141-149.
- Sharma N and Madan M (1988). Effects of various organic wastes alone and with earthworms on the total dry matter yield of wheat and maize. Biological Wastes, 25(1): 33-40.
- Sharma P, Patel AN, Saini MK and Deep S (2012). Field demonstration of *Trichoderma harzianum* as a plant growth promoter in wheat (*Triticum aestivum* L). Journal of Agricultural Science, 4(8): 65-73.
- Sharma SN and Prasad R (2008). Effect of crop residue management on the production and agronomic nitrogen efficiency in a rice-wheat cropping system. Plant Nutrition and Soil Science, 171(2): 295-302.
- Sharma SN, Prasad R and Singh RK (2000). Influence of summer legumes in rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system on soil fertility. Indian Journal of Agricultural Science, 70(6): 357-359.
- Sharma SN, Prasad R, Singh S and Singh P (2000). On-farm trials of the effect of introducing a summer green manure of mung bean on the productivity of a rice-wheat cropping system. The Journal of Agricultural Science, 134(2): 169-172.
- Sheela S and Khimiya S (2013). Vermicompost to save our agricultural land. International Science Congress Association, Research Journal of Agriculture and Forestry Sciences, 1(4): 18-20.

- Shelley IJ, Takahashi-Nosaka M, Kano-Nakata M, Haque MS and Inukai Y (2016). Rice cultivation in Bangladesh: present scenario, problems, and prospects. Journal of International Cooperation for Agricultural Development, 14(4): 20-29.
- Shen J, Yuan L, Zhang J, Li H, Bai Z, Chen X, Zhang W and Zhang F (2011). Phosphorus dynamics: from soil to plant. Plant Physiology, 156(3): 997-1005.
- Sheoran S, Raj D, Antil RS, Mor VS and Dahiya DS (2017). Productivity, seed quality and nutrient use efficiency of wheat (*Triticum aestivum*) under organic, inorganic and integrated nutrient management practices after twenty years of fertilization. Cereal Research Communications, 45(2): 315-325.
- Shewry PR and Hey SJ (2015). The contribution of wheat to human diet and health. Food and Energy Security, 4(3): 178-202.
- Shewry PR, Halford NG, Belton PS and Tatham AS (2002). The structure and properties of gluten: an elastic protein from wheat grain. Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences, 357(1418): 133-142.
- Siavoshi M, Nasiri A and Laware SL (2011). Effect of organic fertilizer on growth and yield components in rice (*Oryza sativa* L.). Journal of Agricultural Science, 3(3): 217-224.
- Sidhu BS, Beri V and Gosal SR (1995). Soil microbial health as affected by crop residue management. *In*: Proceedings of the national symposium on developments in soil science, Ludhiana, India. 2-5 November 1995. Indian Society of Soil Science, pp. 45-46.
- Silva SHSA, Chandrapala AG and Jayalath HAP (2005). Evaluation of different organic manures on soil properties, growth and yield of rice and maize under rice-maize crop rotation. Annals of the Srilanka, Peradeniya University, Department of Agriculture, Sri Lanka 17: 87-94.
- Simpson TW (1990). Agronomic use of poultry industry waste. Poultry Science, 70(5): 1126-1131.

- Sims TJ and Wolf DC (1992). Poultry waste management: Agricultural and Environmental issues. Advanced Agronomy, 52: 1-84.
- Singh AP, Sakal R and Singh BP (1983). Relative effectiveness of various types and methods of Zn application on the rice and maize crops grown in calcareous soil. Plant and Soil, 73(3): 315-322.
- Singh D, Sharma RP, Sankhyan NK and Meena SC (2017). Influence of long-term application of chemical fertilizers and soil amendments on physico-chemical soil quality indicators and crop yield under maize-wheat cropping system in an acid alfisol. Journal of Pharmacognosy and Phytochemistry, 6(3): 198-204.
- Singh KK, Srinivasarao C and Ali M (2008). Phosphorus and mung bean residue incorporation improve soil fertility and crop productivity in sorghum and mung bean-lentil cropping systems. Journal of Plant Nutrition, 31(3): 459-471.
- Singh KK, Srinivasarao CH, Swarnalakshmi K, Ganeshamurthy AN and Kumar N (2011). Influence of legume residues management and nitrogen doses on succeeding wheat yield and soil properties in Indo-Gangetic Plains. Food Legumes, 25(2): 116-120.
- Singh SK, Kumar D and Lal SS (2010). Integrated use of crop residues and fertilizers for sustainability of potato (*Solanum tuberosum*) based cropping systems in Bihar. Indian Journal of Agronomy, 55(3): 203-208.
- Singh V, Singh PN, Yadav RL, Awasthi SK, Joshi BB, Singh RK, Lal RJ and Duttamajumder SK (2010). Increasing the efficacy of *Trichoderma harzianum* for nutrient uptake and control of red rot in sugarcane. Journal of Horticulture and Forestry, 2(4): 66-71.
- Sistani KR, Sikora FJ and Rasnake M (2008). Poultry litter and tillage influences on corn production and soil nutrients in a Kentucky silt loam soil. Soil and Tillage Research, 98(2): 130-139.
- Slafer GA, Calderini DF and Miralles DJ (1996). Yield components and compensation in wheat: opportunities for further increasing yield potential. Increasing yield potential in wheat: Breaking the Barriers: 101-133.


- Sliesaravicius G and Altintas S (2006). Effects of *Trichoderma* spp. on the yield and grain quality of wheat plants grown in an unheated greenhouse. Australian Journal of Experimental Agriculture, 46: 131-136.
- SRDI (2017). Soil Resource Development Institute, Soil sample collection method, Soil building, Farmgate, Dhaka-1215, Bangladesh.
- Sultani MI, Gill MA, Anwar MM and Athar M (2007). Evaluation of soil physical properties as influenced by various green manure legumes and phosphorus fertilization under rain fed conditions. International Journal of Environmental Science and Technology, 4(1): 109-118.
- Sumner ME (2009). Soil Acidification. Encyclopedia of land use, land cover and soil sciences-VII. Encyclopedia of Life Support Systems, Oxford, United Kingdom, pp. 187-208.
- Suparno S, Hakim L and Aidawati N (2016). *Trichoderma* spp. as agent of biological control in local rice diseases in tidal swamp lands in South Kalimantan, Indonesia. Journal of Agriculture and Veterinary Science, 9(1): 1-6.
- Swarup A and Yaduvanshi NPS (2000). Effect of integrated nutrient management on soil properties and yield of rice in alkali soils. Journal of the Indian Society of Soil Science, 48(2): 279-282.
- Taheri RE, Ansari MH and Razavi NA (2017). Influence of cow manure and its vermicomposting on the improvement of grain yield quality of rice (*Oryza sativa* L.) in field conditions. Applied Ecology and Environmental Research, 16(1): 97-110.
- Tahir M, Ayub M, Javeed HMR, Naeem M, Rehman H, Waseem M and Ali M (2011).Effect of different organic matter on growth and yield of wheat (*Triticum aestivum* L.). Pakistan Journal of Life and Social Sciences, 9(1): 63-66.
- Talukdar P, Siddiqa MM, Masum MMI, Habibullah ABM and Bhuiyan AKM (2017). Effect of *Trichoderma* fortified compost on disease suppression, growth and yield of chickpea. International Journal of Environment, Agriculture and Biotechnology, 2(2): 831-839.

- Tapke R (2017). Zero budget natural reforming with the help of cow dung and cow urine. Proceedings of 109th: The IIER International Conference 10-11 July 2017, Pattaya, Thailand.
- Tarique MH and Paul NK (2005). Effects of different N levels on the yield contributing characters on the wheat. Progressive Agriculture, 16(2): 1-5.
- Tejada M, Hernandez MT and Garcia C (2009). Soil restoration using composted plant residues: Effects on soil properties. Soil and Tillage Research, 102(1): 109-117.
- Tewari L and Singh R (2005). Biological control of sheath blight of rice by *Trichoderma harzianum*. Indian Phytopathology, 58: 35-40.
- Tharmaraj K, Ganesh P, Kolanjinathan K, Kumar SR and Anandan A (2011). Influence of vermicompost and vermiwash on physic chemical properties of rice cultivated soil. Current Botany, 2(3): 18-21.
- Thiry DE, Sears RG, Shroyer JP and Paulsen GM (2002). Relationship between tillering and grain yield of Kansas wheat varieties. Kansas State University Agricultural Experimental Station and Cooperative Extension Service, pp. 1-5.
- Umanah EE, Ekpe EO, Ndon BA, Etim ME and Agbogu MS (2003). Effects of poultry manure on growth characteristics, yield and yield components of upland rice in south eastern Nigeria. Journal of Sustainable Agriculture and Environment, 5(1): 105-110.
- USDA (2011). United States Department of Agriculture, Natural Resources Conservation Service, Soil Quality Indicators, pp. 1-2.
- Usman MU, Ullah EH, Warriach EA, Farooq MU and Liaqat AM (2003). Effect of organic and inorganic manures on growth and yield of rice variety "Basmati-2000". International Journal of Agriculture and Biology, 5(4): 481-483.
- Uwah DF, Undie UL and John NM (2014). Comparative evaluation of animal manures on soil properties, growth and yield of sweet maize (*Zea mays* L. saccharata Strut.)Journal of Agriculture and Environmental Sciences, 3(2): 315-331.

- Vasilica ST, Virsta A, Dusa EM and Glavan AM (2009). Waste recycling and compost benefits. Notulae Botanicae Horti Agrobotanici Cluj-Napoca, 37(2): 9-13.
- Vengadaramana A and Jashothan PT (2012). Effect of organic fertilizers on the water holding capacity of soil in different terrains of Jaffna peninsula in Sri Lanka. Journal of Natural Product and Plant Resources, 2(4): 500-503.
- Verma TS and Bhagat RM (1992). Impact of rice straw management practices on yield, nitrogen uptake and soil properties in a wheat rice rotation in northern India. Fertilizer Research, 33(2): 97-106.
- Vinale F, Sivasithamparam K, Ghisalberti EL, Ruocco M, Wood S and Lorito M (2012). *Trichoderma* secondary metabolites that affect plant metabolism. Natural Production Communication, 7(11): 1545-1550.
- Wakene N, Heluf G and Friesen DK (2005). Integrated use of farmyard manure and NP fertilizers for maize on farmers' fields. Journal of Agriculture and Rural Development in Tropics and Subtropics, 106(2): 131-141.
- Walkley AC and Black TA (1935). Estimation of soil organic carbon by chromic acid titration method. Soil Science, 47: 29-38.
- Walter K, Don A, Tiemeyer B and Freibauer A (2016). Determining soil bulk density for carbon stock calculations: A systematic method comparison. Soil Science Society of America Journal, 80(3): 579-591.
- Walters DT, Aulakh MS and Doran JW (1992). Effects of soil aeration, legume residue and soil texture on transformations of macro and micronutrients in soils. Soil Science, 153(2): 100-107.
- Warren JG, Phillips SB, Mullins GL, Keahey D and Penn CJ (2006). Environmental and production consequences of using alum-amended poultry litter as a nutrient source for corn. Journal of Environmental Quality, 35(1): 172-182.
- Wilhelm WW, Johnson JM, Karlen DL and Lightle DT (2007). Corn stover to sustain soil organic carbon further constrains biomass supply. Agronomy Journal, 99(6): 1665-1667.

- Woignier T, Clostre F, Fernandes P, Rangon L, Soler A and Lesueur-Jannoyer M (2016).Compost addition reduces porosity and chlordecone transfer in soil microstructure.Environmental Science and Pollution Research, 23(1): 98-108.
- World Agricultural Outlook Board (2014). World Agricultural Supply and Demand Estimates-527. United States Department of Agriculture, pp. 1-38.
- Yaacob SZ, Abdullah N and Abdullah LC (2019). Potential of *Trichoderma harzianum* as cellulose biodegrader in biocomposting of paddy straw. Journal of Advanced Research in Materials Science, 39: 8-13.
- Yadvinder-Singh, Bijay-Singh, Gupta RK, Ladha JK, Bains JS and Singh J (2008). Evaluation of press mud cake as a source of nitrogen and phosphorus for the ricewheat cropping system in the Indo-Gangetic plains of India. Biology and Fertility of Soils, 44: 755-762.
- Yang Q, Luo W, Jiang Z, Li W and Yuan D (2016). Improve the prediction of soil bulk density by cokriging with predicted soil water content as auxiliary variable. Journal of Soils and Sediment, 16(1): 77-84.
- Yedidia I, Srivastva AK, Kapulnik Y and Chet I (2001). Effect of *Trichoderma harzianum* on microelement concentrations and increased growth of cucumber plants. Plant and Soil, 235(2): 235-242.
- Yildirim M, Yakut Z, Akinci C, Kurt F and Kizilgeci F (2016). Nitrogen rate and timing implementation on durum wheat in a bed planting system. Sains Malaysiana, 45(2): 221-228.
- Yuo T, Yamashita Y, Kanamori H, Matsumoto T, Lundqvist U, Sato K, Ichii M, Jobling SA and Taketa, S (2012). A short internodes (SHI) family transcription factor gene regulates awn elongation and pistil morphology in barley. Journal of Experimental Botany, 63(14): 5223-5232.
- Zaki SS (2016). Effects of compost and nitrogen fertilization on yield and nutrients uptake of rice crop under saline soil. Modern Chemistry and Applications, 4(2): 1-3.

- Zeleke TB, Grevers MCJ, Si BC, Mermut AR and Beyene S (2004). Effect of residue incorporation on physical properties of the surface soil in the South Central Rift Valley of Ethiopia. Soil and Tillage Research, 77(1): 35-46.
- Zhang HQ, Yu XY, Zhai BN, Jin ZY and Wang ZH (2016). Effect of manure under different nitrogen application rates on winter wheat production and soil fertility in dry land. IOP Conference Series: Earth Environmental Science, 39(1): 1-10.
- Zhao Y, Yan Z, Qin J and Xiao Z (2014). Effects of long-term cattle manure application on soil properties and soil heavy metals in corn seed production in Northwest China. Environmental Science and Pollution Research, 21(12): 7586-7595.
- Zhong H, Wang Q, Zhao X, Du Q, Zhao Y, Wang X, Jiang C, Zhao S, Cao M, Yu H and Wang D (2014). Effects of different nitrogen applications on soil physical, chemical properties and yield in maize (*Zea mays* L.). Agricultural Sciences, 5(14): 1440-1447.
- Zia MS, Nizami MI and Salim M (1994). Problems of soil degradation in Pakistan. The collection of land degradation data. Report of the expert consultation on the Asian network on problem soils. RAPA Publication 3, pp. 179-202.
- Zink RTA and Allen MF (1998). The effects of organic amendments on the restoration of a disturbed coastal sage scrub habitat. Restoration Ecology, 6(1): 52-58.



APPENDICES

Name of AEZ	Land type	Organic ma (Ave		Total depletion of organic
Name of ALL	Land type	1969-1970	1989-1990	matter (%)
	II. 11 1	17.8	12	22.59
1. Madhupur Tract	Highland	(13-24)	(6-17)	32.58
2 Derind Treest	Highland &	14.5	11.5	20.00
2. Barind Tract	MHL	(10.6-20)	(9-14)	20.69
3. Old Himalayan Piedmont	IT -1.1	13.2	12	0.0
Plain	Highland	(10-16.5)	(8-15)	9.0
4 Tiste Maanden Elee dalein	Highland &	15.5	12.3	20.6
4. Tista Meander Floodplain	MHL	(14.6-16)	(10-15)	20.6
	Highland &	20.4	13.2	25.2
5. Northern and Eastern Hills	MHL	(14.9-24.6)	(10-15)	35.3
6. Old Meghna Estuarine	YT' 1 1 1	21.6	11.7	45.0
Floodplain	Highland	(19.2-26.1)	(10-15)	45.8
7. High Ganges River	III ab land	12.1	9.8	10.0
Floodplain	Highland	(6.4-16.1)	(3-14)	19.0
	MIII 1 1	15.6	12.3	01.15
8. Old Brahmaputra Floodplain	MHL land	(10.9-21.6)	(9-15)	21.15

Table 1: Scenario of soil organic matter depletion during 20 years in different Agro-Ecological Zones of Bangladesh.

Source: Mia et al. (1993), MHL = Medium High Land.

								?
							Mg	Mg
						В	В	В
ents					Zn	Zn	Zn	Zn
Nutrients				S	S	S	S	S
Z			Κ	Κ	Κ	Κ	Κ	Κ
		Р	Р	Р	Р	Р	Р	Р
	Ν	Ν	Ν	Ν	Ν	Ν	Ν	Ν
Year	1951	1957	1960	1980	1982	1995	2000	2010

Table 2: Emergence of new nutrient deficiency with time in soil.

Source: Jahiruddin and Satter (2010).

Table 3: Nutrient elements content in different organic manures.

Organic matter	N (%)	P (%)	K (%)	S (%)	Zn (%)
Green manure	0.70	0.40	0.38	0.20	-
Rice straw	0.50	0.08	1.60	0.09	0.01
Cow dung	1.20	0.80	1.30	0.13	0.01
Compost	1.80	0.25	0.94	0.20	0.02
Vermicompost	1.75	1.00	0.75	0.03	0.02
Poultry manure	1.85	0.60	0.76	0.06	0.05

Analysed at Soil Resource Development Institute (SRDI) Laboratory, Shyampur, Rajshahi, Bangladesh.


Year	Month	*Te	mperatur	e (°C)	*]	Humidity	(%)	**Rainfall	*Sunshine
Teal	Monui	Max.	Min.	Average	6.0 am	6.0 pm	Average	(mm)	(hours)
	September	34.04	26.35	30.20	96.87	86.20	91.53	157.20	5.03
2016	October	33.07	23.92	28.50	96.97	83.48	90.23	94.60	7.73
2016	November	29.99	17.52	23.76	96.67	82.53	89.60	0.00	6.62
	December	26.15	13.65	19.90	97.87	83.45	90.66	0.00	4.49
	January	25.18	11.02	18.10	94.35	72.84	83.60	2.20	5.73
	February	28.74	13.50	21.12	95.25	60.93	78.09	0.00	6.34
	March	31.29	18.52	24.91	93.74	62.29	78.02	43.50	6.37
2017	April	35.06	23.24	29.15	92.47	63.57	78.02	102.00	6.14
2017	September	33.93	26.59	30.26	97.37	86.77	92.07	133.40	4.02
	October	31.81	24.38	28.09	97.55	87.68	92.61	237.50	5.98
	November	29.35	17.44	23.40	96.83	83.73	90.28	5.20	7.33
	December	26.18	14.64	20.41	96.70	84.58	90.64	19.60	6.06
	January	22.49	8.50	15.50	98.26	77.97	88.12	0.00	5.79
	February	28.90	14.20	21.55	96.10	65.40	80.75	12.60	5.50
	March	34.02	19.05	26.54	93.03	51.35	72.19	8.60	7.99
2010	April	34.13	22.21	28.17	93.16	66.16	79.66	142.50	7.68
2018	September	34.32	26.12	30.22	96.80	82.47	89.63	117.80	6.18
	October	32.11	21.84	26.98	95.35	83.00	89.18	85.80	7.23
	November	29.99	16.73	23.36	97.33	82.60	89.97	0.00	7.52
	December	25.27	11.64	18.45	94.94	79.61	87.27	17.20	6.41
	January	25.46	10.23	17.84	94.65	73.29	83.97	0.00	7.37
2010	February	27.62	13.17	20.40	95.36	65.29	80.32	47.20	7.63
2019	March	32.08	17.45	24.77	95.42	53.03	74.23	68.00	7.84
	April	34.52	22.95	28.74	94.27	64.87	79.59	113.90	7.16

Table 4: Monthly minimum, maximum and average air temperature (°C), rainfall (mm)and relative humidity (%) during the experimental period are given below:

*= Monthly average **= Monthly total.

Source: First class Meteorogical Observatory, Shyampur, Rajshahi, Bangladesh.

List of Publications and Abstract



EFFECTS OF PROBIOTIC AND ORGANIC FERTILIZERS AS SOIL AMENDMENTS ON THE GROWTH AND YIELD OF WHEAT

Md. Anwar Hossain^{1,2}, Md. Mahmodol Hasan³, Md. Belal Hossain³ and S.M. Shahinul Islam¹*

 ¹Plant Biotechnology and Genetic Engineering Laboratory, Institute of Biological Sciences,
 ³Plant Pathology Laboratory, Agronomy and Agricultural Extension Department, University of Rajshahi, Rajshahi - 6205 (Bangladesh)
 ²Directorate of Secondary and Higher Education, Dhaka - 1000 (Bangladesh)
 *e-mail: shahinul68@gmail.com

(Received 26 December, 2020; accepted 8 March, 2021)

ABSTRACT

Present study aimed to comprehensively assess the impact of organic amendments on crop health without compromising on growth and yield. To evaluate the effect of organic amendment a comprehensive study in wheat was conducted in a randomized complete block design with three replications for three consecutive wheat growing seasons (2017-2019). The experiment comprised of was nine soil amendment treatments wherein organic manures, probiotics and chemical fertilizer were used. The dry matter accumulation, leaf area index, and crop growth rate_were assessed at 25, 40, 55, 70 and 85 days after sowing (DAS). At 85 DAS, the maximum TDM accumulation (769.89 g m⁻²) was recorded by the chemical fertilizer which was statically similar (752.48 g m⁻²) with poultry manure + vermicompost + green manure (Sesbania rostrata). Similar trend was found in case of LAI (4.07 followed by 3.84) at 55 DAS. Likewise, the maximum CGR 20.30 and 19.07 g m⁻² d⁻¹ was noted at 55-70 DAS under the same treatment condition. The study revealed that the growth attributes significantly increased in chemical fertilizer than poultry manure + vermicompost + green manure treatment and untreated control. Application of probiotic and organic manures significantly influenced the growth attributes of wheat. The information generated is very useful for farmers and agriculturist for enhancing the growth and yield of wheat.

Keywords: Green and poultry manure, growth attributes, probiotics, vermicompost, yield, wheat

INTRODUCTION

Wheat (*Triticum aestivum* L.) has long been a highly popular cereal crop all over the world including Bangladesh. It is recognized as a pioneer source of protein for human consumption than other cereals. The major composition of wheat includes 66.0-71.6% carbohydrates, 13-16.7% protein, 2.5-3.1% fats and 2.5-3% crude fiber (Khan, 1984). Nearly 50% world grain production is occupied by wheat (Banglapedia, 2014; Haydar *et al.*, 2020). Bangladesh possesses 31st positions rank among wheat growing countries in the world. In Bangladesh, wheat crop occupies 4% of the total cropped area, 11% of area cropped in *rabi* and contributes 7% to the total output of food cereals (Anonymous, 2008).

Bangladesh is known as a rapidly increasing populated country and thus land under cultivation is decreased to provide their accommodation. To keep harmony with increased population intensive cultivation of wheat and other cereal crops like barley, maize, rye, etc. and its cultivated lands were affected by using many chemicals and fertilizers that directly affects to food security (Haque and Islam, 2018). The long-term application of excess inorganic fertilizer or poor fertilization practice results in soil acidification and nutrient depletion, besides deteriorating the soil and environmental health (Liebig et al., 2002). Further, synthetic fertilizers are chemical in natures that are harmful human, animals and environments. It has been reported that initial period of three years organic farming could give equal or more yields as compared to chemical use alone (Ramesh et al., 2005). To create a favorable crop growth environment and sustainable production, organic farming may be an alternative way without using chemical fertilizers in the soil (Kshirsagar, 2008) and it may contribute to increase 80% of the yield the developing countries (Badgley et al., 2007). Probitic i.e. Trichoderma is a common symbiont of plant root system (Vinale et al., 2012) which is used as biocontrol agent and plant growth enhancer (Talukder et al., 2017). Organic manures mainly poultry manure accelerates vegetative growth reported by Rasul et al. (2015) and Abubakar and Ali (2018). So, for achieving better crop yield and growth and retaining sound soil health, the use of high yielding varieties along with organic amendments are considered important. Hence, in view of above, a study was undertaken to assess the impact of probiotic and organic manure on wheat growth and yield.

MATERIALS AND METHODS

Preparation of soil and plant growth

The present research work was carried out in the experimental field and in the Laboratory of Plant Biotechnology & Genetic Engineering, Institute of Biological Sciences and Plant Pathology Lab., Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh during July 2016 to June 2019. The experimental field geographically was situated at $24^{\circ}17^{\circ}$ N latitude and $88^{\circ}28^{\circ}$ E longitude at an elevation of 20 m masl and belonged to the Agro-Ecological Zone-11 (AEZ-11). The soil of experimental field was poorly drained with moderately permeability, loam and slightly alkaline (pH, 8.1). The experimental soil had moisture content 19.4%, particle density 2.65 g cc⁻¹, bulk density 1.27 g cc⁻¹ and porosity 51.34%. Further, the soil had organic matter content of 1.2%, K 0.150 cmol (+) kg⁻¹, total N 0.07%, P 26.30 µg g⁻¹, S 12.50 µg g⁻¹, Zn 0.75 µg g⁻¹ and EC 145 µs cm⁻¹ with C: N of 10:1.

Experimental design and treatments

The experiment was laid out in a randomized complete block design with three replications. The treatments comprised of three wheat varieties viz., 'BARI Wheat-28', 'BARI Wheat-29' and 'BARI Wheat-30' [Bangladesh Agricultural Research Institute, Joydebpur, Gazipur] and nine soil amendments [control, rice straw + vermicompost + green manure, cow dung + vermicompost + green manure, compost + vermicompost + green manure, poultry manure + vermicompost + green manure, Trichoderma harzianum + vermicompost + green manure, mung bean straw + vermicompost + green manure, T. viride + vermicompost + green manure, and chemical fertilizer]. The unit plot size was 5.0 m^2 having a plot to plot 0.5 m, bed to bed 0.25 cm distance and 1 m from the surrounding boundary. The total unit plot was 81. Crop residues (rice straw and mung bean straw), cow dung, compost and poultry litter were applied at the rate 10 t ha⁻¹ 7 days prior to sowing [BARI, 2014]. Vermicompost was applied 5 t ha⁻¹ and *Trichoderma* spp. suspension $(1 \times 10^6 \text{ cfu g}^{-1} \otimes 5 \text{ kg ha}^1)$ before sowing. In case of chemical treatment, one third of urea (200 kg ha⁻¹), triple super phosphate 160 kg ha⁻¹, muriate of potash 45 kg ha⁻¹ and gypsum 115 kg ha⁻¹ were applied as basal dose (BARI, 2014). Rest urea was applied in two installments, one at 21 days after sowing (DAS) and another at 55 DAS. Crop residues were collected from the research field of IBSc, RU. Cow dung and compost were collected from a trained farmers pit under DAE Bagatipara, Natore. Poultry litter was collected from a poultry farm. Vermicompost was collected from Sumi Seed Vander, Rajshahi. Trichoderma spp. were cultured in Plant Pathology Laboratory, Department of Agronomy & Agriculture Extension, University of Rajshahi, Rajshahi, Bangladesh. The collected organic manures contained NPK as green manure (N 0.70%, P 0.40% and K 0.38%), rice straw (N 0.50%, P 0.08% and K 1.60%), cow dung (N 1.20%, P 0.80% and K 1.30%), compost (N 1.80%, P 0.25% and K 0.94%), vermicompost (N 1.75%, P 1.00% and K 0.75%) and poultry manure (N 1.85%, P 0.60% and K 0.76%) were used respectively.

Data collection on growth attributes

<u>Total dry matter (TDM)</u>: For TDM, five plants were collected at 25, 40, 55, 70 and 85 DAS from each plot and sundried before the collected samples were packed individually in a labeled brown paper bag and dried in an oven at 70-80°C for 72 h till constant weight was attained.

Leaf area index (LAI): For LAI determination, five plants were collected from each plot at 25, 40, 55, 70 and 85 DAS. At each time period, the collected plants were separated into leaf, stem and spike (when appeared) and leaf area was measured by disc method (Radford, 1967). For this, every leaf was segmented into three portion and middle portion length and breadth of each leaf were measured. Among the total leaves of a plant, each leaf was packed individually in a labeled brown paper bag and placed in an oven at 70-80°C for 72 h. The leaf area was measured and calculated by using the formula:

Area of leaf =
$$\frac{\text{Weight of leaves} \times \text{Area of segments}}{\text{Weight of segments}} \times \text{cm}^2$$

As growth and physiological parameters leaf area index (LAI) and crop growth rate (CGR) were calculated by using standard formulae (Radford, 1967).

Leaf area index (LAI) =
$$\frac{\text{Leaf area}}{\text{Ground area}}$$

<u>Crop growth rate (CGR)</u>: The crop growth rate was calculated from total dry matter in respect of sampling dates using the formulae (Radford, 1967):

Crop groth rate (CGR) =
$$\frac{(W2-W1)}{(t2-t1)}$$
 × g m⁻² d⁻¹

Where, W_2 and W_1 are the total dry weight plant at different DAS and t_2 and t_1 are the harvest time of total dry weight at the latter and former, respectively.

Statistical analysis

The data were compiled and tabulated for statistical analysis. The data was analyzed using analysis of variance and Duncan's multiple range tests through r-studio by the "Agricolae" package.

RESULTS AND DISCUSSION

The study revealed that the chemical fertilizer and combination of poultry manure + vermicompost + green manure (*Sesbania rostrata*) showed parallel results regarding total dry matter (TDM), leaf area index (LAI) and crop growth rate (CGR) with few exceptions.

Total dry matter

The study was planned to investigate the effects of different organic manures as soil amendment on TDM accumulation of wheat at 25, 40, 55, 70 and 85 DAS (Table 1). The soil amendments showed significant variations in TDM production. Three years pooled data revealed that TDM accumulation at 25 DAS was highest (30.7 g m⁻²) due to the application of chemical fertilizer followed by poultry manure + vermicompost + green manure (28.5 g m⁻²), while untreated control gave the lowest value (18.0 g m⁻²). The maximum TDM accumulation at 40, 55 and 70 DAS were recorded from chemical

fertilizer (102.4, 227.9 and 533.3 g m⁻²) and minimum in untreated control (51.8, 136.1 and 363.1 g m⁻²). At the peak (85 DAS) of TDM accumulation, the most statistical identity was found between chemical fertilizer (769.9 g m⁻²) and poultry manure + vermicompost + green manure (752.5 g m⁻²). Poultry manure-based treatment, in general followed a pattern similar to that of chemical fertilizer treatment for TDM accumulation. Generally, with the advancement of time the TDM accumulation of wheat positively increased. This increment of TDM might be for creating a favourable environment for plant growth activities employed by probiotic and organic amendments of soil (Ahmad *et al.*, 2008). Such increment of TDM might be found for the development of a large number of late tillers (Balyan, 1992; Rahman *et al.* 2006). The improvement of dry matter accumulation with poultry manure + vermicompost + green manure possibly due to high content of essential nutrients present in poultry manure, which accelerates vegetative growth (Muhammad *et al.* 2018). This finding is supported by Bakhtiar *et al.* (2017) who observed that N in poultry manure significantly improves leaf and stem diameter of wheat.

 Table 1: Effect of probiotic and organic fertilizers for soil amendment on total dry matter (g m⁻²) in wheat

	matter (g m) m wheat																			
	T	DM at					40 DA			DM at				DM at	70 DA	<u>s</u>	1	DM at	85 DA	S
Т		2017-	2018-	Mean	2016-	2017-	2018- 2019	Mean	2016-	2017- 2018	2018-	Mean	2016-	2017-	2018-	<u>S</u> Mean	2016-	2017-		Mean
		2018											2017	2010	2017		2017	2018		
	18.0	19.1					58.9						329.9							553.0
T ₀	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.6f	0				2.0h		3.1h	3.60		0	7.4h			18.6g				
		22.2				69.0							383.7			406.8				
T_1	±	±	±	± .	±	±	±	±	±	±	±	±	±	±	±	±	±	±	. ±	±
					0	-	0		0		0		10.0g	0					0	
T	21.5		24.6										412.7						680.8	
T_2	±	±	±	±	±	±	±	±	±	±	,±	±	±	±	±	±	±	±	±	±
	0.4e	0.4d					2.5e		4.0e											13.3cd
т	22.8	25.2			78.5		96.9	86.7	189.1				433.4							
T3	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
		0.4c					1.8d													14.9bc
T4	26.0			28.5									472.1							
14	±	±	±	±	±	±	±	±	±	± 4 Ohed	±	±	± 11.7b	±	±	±	±	±	±	±
	24.0	26.4		26.1									444.6							
T ₅	24.0 ±	20.4 ±	20.1 ±	20.1 ±	61.5 ±	89.0 ±	101.8 ±	90.8 ±	192.5 ±	205.5 ±	±	204.9 ±	444.0 ±	475.5 ±	498.5 ±	472.2 ±	000.0 ±	105.0 ±	132.5 ±	107.4 ±
15		± 0.5b		- 19 <u>77</u> ()	1000	1000	± 2.3c			± 3.8ghi						± 15.6c				
	21.3	22.7	24.3										398.5							
T ₆	±	+	±	±	±	+	±	+ 15.5	±	±	+	±	±	+17.1	+	+21.5	±	±	±	+
10	1000	0.4e							4.3f											15.8de
	23.1	25.7	27.2	25.4	80.8	85.9	96.3		185.1				438.6							
T ₇	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
~ /		0.5bc																		13.2bc
	27.9	29.4		30.7	92.4								504.0							
T ₈	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
	0.9a	0.8a	1.4a	2.1 a	2.5a	2.6a	3.3a	7.9a	5.5a	5.4ab	6.4a	7.9a	4.6a	13.8a	14.6a	16.6a	20.1a	21.0a	22.1a	15.6a
LS	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**
CV	1.9	2.2	2.6	6.2	2.2	2.4	2.5	6.5	2.0	2.3	2.3	6.6	1.9	2.2	2.2	6.5	1.8	2.2	2.1	6.5
(%)	1.9	2.2	2.0	0.2	2.2	2.4	2.5	0.5	2.0	2.5	2.5	0.0	1.9	2.2	2.2	0.5	1.0	2.2	2.1	0.5

Each value represents the average of three replicates. In column, the mean values superscripted with similar letter(s) are statistically identical as per Duncan's Multiple Range Test. T = Treatments, $T_0 = control$, $T_1 = rice$ straw + vermicompost + green manuring, $T_2 = cow dung + vermicompost + green manuring, <math>T_3 = compost + vermicompost + green manuring, <math>T_4 = poultry$ manure + vermicompost + green manuring, $T_5 = T$. *harzianum* + vermicompost + green manuring, $T_6 = mung$ bean straw + vermicompost + green manuring, $T_7 = T$. *viride* + vermicompost + green manuring, $T_8 = chemical fertilizer$, LS = Level of significance, **= 1% level of significance, CV = Coefficient of variation


Leaf area index (LAI)


Leaf area index expresses the efficacy of photosynthetic processes which ultimately indicates the good vegetative growth and yield. The data indicated that the positive residual effects of manures, especially poultry manure, on LAI were still evident for three growing seasons. It was observed that all the soil amendments provided significant results for LAI at every sampling date in each year of this study. The chemical fertilizer performed better performance for LAI followed by poultry manure + vermicompost + green manure in comparison to the untreated control (Table 2). The pooled data of three years study showed highest value (4.1) of LAI from chemical fertilizer and lowest LAI (2.4) was in untreated control at 55 DAS. The increment of LAI had an ascending tendency up to 55 DAS, whereas descending nature was exhibited in the next 70 and 85 DAS during the experiment. Throughout the study period chemical fertilizer and poultry manure + vermicompost + green manure exhibited almost similar results (2.5 and 2.4) at the last sampling date of LAI. Al-Rawashdeh and Abdel-Ghani (2008) supported the decline of LAI due to the losses of nitrogen (gaseous form) at the end of vegetative stage and leaf senescence. The possible reason of parallel performance of chemical fertilizer and poultry manure + vermicompost + green manure might be due to delivering more nutrients required for crop growth which were released from poultry manure combination. Continuous release of plant nutrients from organic manures which directed to plant growth and

Table 2: Effect of probiotic and organic fertilizers for soil amendment on leaf area index (LAI) in wheat

	in wheat																			
	L	AI at 2	25 DA	<u>.S</u>	Ī	AI at	40 DA	S	Ī	LAI at	55 DA	<u>S</u>	L	AI at	70 DA	S	Ī	AI at	85 DA	S
Т	2016- 2017	2017- 2018	2018- 2019	Mean	2016- 2017	2017- 2018	2018- 2019	Mean												
2 	0.2	0.3	0.2	0.2	1.2	1.4	1.3	1.3	2.1	2.5	2.5	2.4	2.0	2.2	2.3	2.1	1.4	1.5	1.6	1.5
T_0	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
	0.0j	0.0i	0.0i	0.0i	0.0f	0.0i	0.0i	0.1g	0.1i	0.1i	0.1i	0.1g	0.1e	0.1g	0.1i	0.1h	0.0f	0.0e	0.1i	0.1g
	0.2	0.3	0.3	0.3	1.3	1.6	1.6	1.5	2.3	2.7	2.8	2.6	2.1	2.4	2.6	2.3	1.5	1.8	1.8	1.7
T_1	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
	0.0h	0.0h	0.0h	0.0h	0.0e	0.1e	0.1h	0.1f	0.1h	0.1h	0.1h	0.1fg	0.1e	0.1g	0.1h	0.2g	0.0e	0.1d	0.1h	0.1f
	0.3	0.4	0.4	0.4	1.5	1.7	1.8	1.7	2.5	3.0	3.1	2.9	2.4	2.8	2.9	2.7	1.6	1.9	2.2	1.9
T_2	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±
	0.0f	0.0f	0.0f	0.0f	0.1d	0.1d	0.1f	0.1e	0.1f	0.1f	0.1f	0.2e	0.1d	0.1e	0.1f	0.2e	0.0d	0.1d	0.1f	0.2e
T	0.4	0.5	0.5	0.5	1.7	1.9	2.1	1.9	2.8	3.2	3.3	3.1	2.5	3.0	3.1	2.9	1.8	2.1	2.4	2.1
T_3	± 0.0e	± 0.0e	± 0.0e	± 0.0e	± 0.0c	± 0.1c	± 0.1d	± 0.1d	± 0.1e	± 0.1e	± 0.1e	± 0.2d	± 0.1c	± 0.1d	± 0.1e	± 0.2d	± 0.1c	± 0.1c	±0.1d	± 0.2d
	0.0e	0.0e	0.0e	0.0e	1.9	0.1c 2.3	2.4	2.2	3.4	4.0	4.2	3.8	0.1c 2.9	3.5	3.8	0.2d 3.4	0.1c 2.1	0.1c		2.4
T_4	0.0 ±	0.0 ±	±.0.7	0.0 ±	1.9 ±	2.5 ±	2.4 ±	2.2 ±	5.4 ±	4.0 ±	4.2 ±	5.0 ±	2.9 ±	5.5 ±	5.0 ±	5.4 ±	2.1 ±	2.5	2.8	2.4 ±
14	± 0.0b	0.0b	⊥ \0.0b	± 0.0b	±0.0b	± 0.0a	± 0.1b	± 0.2b	± 0.1b	± 0.1b	± 0.1b	0.2b	0.1b	0.1b	0.1b	0.3b	± 0.0ab	±0.1a	±0.1b	0.2b
	0.5	0.5	0.6	0.5	1.9	2.1	2.2	2.1	3.2	3.6	3.8	3.5	2.9	3.2	3.4	3.2	2.0			2.2
T_5	±	±	±	±	±	±	±	±	±	±	±	±	±,	±	±	±	±.0	2.2	2.5	+
- 5	0.0c	0.0c	0.0c	0.0c	0.0b	0.1b	0.1c	0.1c	0.1c	0.1c	0.1c	0.2c	0.1b	0.1c	0.1c	0.2c	0.0b	$\pm 0.1b$	±0.1c	0.1c
	0.3	0.3	0.4	0.3	1.5	1.7	1.7	1.6	2.4	2.8	2.9	2.7	2.2	2.6	2.7	2.5	1.6			1.8
T_6	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	1.8	2.1	±
	0.0g	0.0g	0.0g	0.0g	0.0d	0.1de	0.1g	0.1e	0.0g	0.1g	0.1g	0.2ef	0.0d	0.1f	0.1g	0.2f	0.0de	±0.1d	±0.1g	0.1e
	0.5	0.5	0.5	0.5	1.7	1.9	2.0	1.9	3.1	3.4	3.6	3.3	2.6	3.0	3.2	3.0	1.8	2.1	2.3	2.1
T_7	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±		$\pm 0.1e$	±
	0.0d	0.0d	0.0d	0.0d	0.0c	0.1c	0.1e	0.1d	0.1g	0.1d	0.1d	0.2c	0.1c	0.1d	0.1d	0.2d	0.0c	10.10	10.10	0.1d
	0.6	0.7	0.7	0.7	2.2	2.3	2.6	2.4	3.7	4.1	4.3	4.1	3.1	3.7	4.0	3.6	2.2	2.6	2.9	2.5
T_8	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±	±		$\pm 0.1a$	±
	0.0a	0.0a	0.0a	0.0a	0.1a	0.1a	0.1a	0.1a	0.1g	0.1a	0.1a	0.2a	0.1a	0.1a	0.1a	0.3a	0.1a	± 0.1a	± 0.1a	0.2a
LS	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**	**
CV (%)	3.7	4.1	2.4	4.1	6.4	6.4	2.4	4.3	2.2	1.6	1.4	7.0	6.3	5.8	1.6	4.1	6.6	6.4	2.1	4.3

Each value represents the average of three replicates. In column, the mean values superscripted with similar letter(s) are statistically identical as per Duncan's Multiple Range Test. T = Treatment, $T_0 = control$, $T_1 = rice$ straw + vermicompost + green manuring, $T_2 = cow dung + vermicompost + green manuring, <math>T_3 = compost + vermicompost + green manuring, T_4 = poultry manure + vermicompost + green manuring, <math>T_5 = T$. *harzianum* + vermicompost + green manuring, $T_6 = mung$ bean straw + vermicompost + green manuring, $T_7 = T$. *viride* + vermicompost + green manuring, $T_8 = chemical fertilizer$, LS = Level of significance, **= 1% level of significance, CV = Coefficient of variation.





production of more leaves resulting in increased leaf area (Sims and Wolf, 1992). Ogundare *et al.* (2012) and Kareem *et al.* (2017) confirmed the judgments through their research outcome.

Crop growth rate (CGR)

The data indicated that the soil amendments had significant effect on wheat CGR for all sampling dates. However, with few exceptions pooled data revealed that the chemical fertilizer had the highest activity in the enhancement of CGR than poultry manure + vermicompost + green manure where the lowest value was recorded in untreated control (Fig. 1). The maximum CGR rate (20.3 g m⁻² d⁻¹) was found in chemical fertilizer treatment while 2nd highest value (19.1 g m⁻² d⁻¹) was recorded from poultry manure-based combination. The minimum CGR value (15.1 g m⁻² d⁻¹) was recorded in untreated control at 55-70 DAS. In case of 70-85 DAS, poultry manure + vermicompost + green manure showed the higher value (16.7 g m⁻² d⁻¹) for CGR than chemical fertilizer treatment (15.8 g m⁻² d⁻¹). The crop growth rate was demonstrated slow increase at the initial stage of wheat and later

reached the peak at 55-70 DAS, and then declined in late growth stage. This observation may be linked to the delayed nutrient availability from poultry manure at early plant growth stage that produced less dry matter and then produced more dry matter in slow-fast-slow growth trend (Geng *et al.*, 2019). However, afterwards the decrease in CGR was noticed due to leaf senescence and tiller drying. These results are is an agreement with those of Boateng *et al.* (2006) and Ram (2017).

Conclusion: Chemical fertilizer gave effective results but its long-term use is not good for crop and soil health, therefore, organic soil amendment needs to be emphasized. Under this study potential and sustainable improvement of wheat growth with organic soil amendment was found. Inclusion of green manure, vermicompost and poultry manure may be a suitable alternative of chemical fertilizer for wheat growth. The capacity of surprising decomposition and nutrient release from those organic manures could be used to synchronize the nutrient demand of wheat. Data revealed that organic manures particularly poultry manure is a healthier source for improvement wheat growth as compared to other manures.

REFERENCES

- Abubakar, Z.A. and Ali, A.D. 2018. Screening effect of organic manure on the vegetative growth of maize (*Zea mays* L.). *Journal of Bioscience and Agriculture Research*, **16**(2): 1356-1364.
- Ahmad, R., Naveed, M., Aslam, M., Zahir, Z.A., Arshad, M. and Jilani, G. 2008. Economizing the use of nitrogen fertilizer in wheat production through enriched compost. *Renewable Agriculture and Food Systems*, 3: 243-349.
- Al-Rawashdeh, Y.A. and Abdel-Ghani, A.H. 2008. Effect of nitrogen application timing on dry matter and nitrogen assimilation and partitioning in six wheat cultivars under rain-fed conditions of Jordan. Archives of Agronomy and Soil Science, 54(2): 149-162.
- Anonymous 2008. *Statistical Year Book of Bangladesh-2006*. Bangladesh Bureau of Statistics (BBS), Planning Division, Ministry of Planning, Government of the People's Republic of Bangladesh.
- Badgley, C., Moghtader, J., Quintero, E., Zakem, E., Chappell, M.J., Aviles-Vazquez, K., Samulon, A. and Perfecto, I. 2007. Organic agriculture and the global food supply. *Renewable Agriculture and Food Systems*, 22(2): 86-108.
- Bakhtiar, M., Afridi, M.Z., Munsif, F., Arshad, I.U., Rehman, S., Khan, S., Adil, M., Khan, M.R., Khan, M.N. and Farooq, M. 2017. Dry matter partitioning and grain yield of wheat as affected by sources, methods and timing of nitrogen application. *Pure and Applied Biology*, 6(4): 1198-1215.
- Balyan, J.S. 1992. Effect of organic recycling on yield attributes, nitrogen concentration and uptake in wheat (*Triticum aestivum* L.). *Indian Journal of Agronomy*, **37**(4): 701-704.
- Banglapedia, M.S. 2014. National Encyclopedia of Bangladesh, Asiatic Society of Bangladesh.
- BARI. 2014. Krishi Projukti Hatboi (Handbook on Agro-Technology), (6th edn.), Gazipur, Bangladesh.
- Boateng, S.A., Zickerman, J. and Kernaharens, M. 2006. Poultry manure effect growth and yield of maize. West African Journal of Applied Ecology, 9: 1-11.
- Geng, Y., Cao, G., Wang, L. and Wang, S. 2019. Effects of equal chemical fertilizer substitutions with organic manure on yield, dry matter, and nitrogen uptake of spring maize and soil nitrogen distribution. *PloS One*, 14(7): 1-16.
- Haque, M. and Islam, S.M.S. 2018. Effect of sowing time on yield and yield contributing characters in comparison with local and European genotypes of barley (*Hordeum vulgare L.*). SKUAST Journal of Research, 20(1): 27-31.

- Haydar, F.M.A., Ahamed, M.S., Siddique, A.B., Uddin, G.M., Biswas, K.L. and Alam, M.F. 2020. Estimation of genetic variability, heritability and correlation for some quantitative traits in wheat (*Triticum aestivum L.*). *Journal of Bio-Science*, 28: 81-86.
- Kareem, I., Jawando, O.B., Eifediyi, E.K., Bello, W.B. and Oladosu, Y. 2017. Improvement of growth and yield of maize (*Zea mays* L.) by poultry manure, maize variety and plant population. *Cercetări Agronomice în Moldova*, 50(4): 51-64.
- Kshirsagar, K.G. 2008. Impact of organic farming on economics of sugarcane cultivation in Maharashtra. *Indian Journal of Agricultural Economics*, **63**(3): 396-405.
- Liebig, M.A., Varvel, G.E., Doran, J.W. and Wienhold, B.J. 2002. Crop sequence and nitrogen fertilization effects on soil properties in the Western Corn Belt. *Soil Science Society of American Journal*, **66**: 596-601.
- Muhammad, F.J., Muhammad, D.A., Waqas, L., Haseeb, A. and Wazir, R. 2018. Effect of poultry manure and phosphorus on phenology, yield and yield components of wheat. *International Journal of Current Microbiology and Applied Sciences*, 7(5): 3751-3760.
- Ogundare, K., Agele, S. and Aiyelari, P. 2012. Organic amendment of an ultisol: effects on soil properties, growth, and yield of maize in Southern Guinea savanna zone of Nigeria. *International Journal of Recycling of Organic Waste in Agriculture*, **1**(1): 11 doi:10.1186/2251-7715-1-11.
- Radford, P.J. 1967. Growth analysis formulae-their use and abuse 1. Crop Science, 7: 171-175.
- Rahman, M.A., Paul, N.K. and Sarker, M.A.R. 2006. Study on growth and yield of wheat with irrigation and split application of nitrogen. *Progressive Agriculture*, **17**(1): 1-7.
- Ram, A.A.M. 2017. Effective use of cow dung manure for healthy plant growth. International Journal of Advanced Research and Development, 2(5): 218-221.
- Ramesh, P., Singh, M. and Rao, A.S. 2005. Organic farming: Its relevance to the Indian context. *Current Science*, 88(4): 561-568.
- Rasul, G.A.M., Ahmed, S.T. and Ahmed, M.Q. 2015. Influence of different organic fertilizers on growth and yield of wheat. *American-Eurasian Journal of Agriculture and Environmental Sciences*, 15(6): 1123-1126.
- Sims, T.J. and Wolf, D.C. 1992. Poultry waste management: Agricultural and environmental issues. *Advanced Agronomy*, **52**: 1-84.
- Talukdar, P., Siddiqa, M.M., Masum. M.M.I., Habibullah, A.B.M. and Bhuiyan, A.K.M. 2017. Effect of *Trichoderma* fortified compost on disease suppression, growth and yield of chickpea. *International Journal of Environment, Agriculture and Biotechnology*, 2(2): 831-839.
- Vinale, F., Sivasithamparam, K., Ghisalberti, E.L., Ruocco, M., Wood, S. and Lorito, M. 2012. *Trichoderma* secondary metabolites that affect plant metabolism. *Natural Production Communication*, 7: 1545-1550.

SUSTAINABILITY OF WHEAT (*Triticum aestivum* L.) YIELD AND IMPROVEMENT OF SEED QUALITY THROUGH PROBIOTIC AND ORGANIC SOIL AMENDMENT

M.A. Hossain^{1, 2}, S.M.S. Islam¹ and M.M. Hasan³

¹Plant Biotech.& Genetic Eng. Lab., Inst. of Biol. Sci., University of Rajshahi ²Directorate of Secondary and Higher Education, Dhaka-1000 ³Plant Path. Lab., Agron. & Agril. Extn. Dept., University of Rajshahi Corresponding E-mail:mahmudul742001@yahoo.com

(Received: 27 November 2020, Accepted: 31 January 2021)

Keywords: Poultry manure, vermicompost, green manure, yield, seed quality, wheat

Abstract

In search of alternative counter to harmful effects of chemical fertilizers on soils and environment, probiotic and organic manures-based fertilizer management options need to be evaluated. The experiments were designed as randomized complete block design (RCBD) consisting of three wheat varieties and nine soil amendment treatments. The result revealed that organic amendments had prominent and variable effects on studied parameters and statistically at par with chemical fertilizer. Some yield-associated parameters like spike length, spikeletsspike⁻¹, fertile spikelets spike⁻¹, grains spike⁻¹, grains weight spike⁻¹ and 1000-grain weight were significantly influenced by organic amendments. Moreover, grain yield and straw yield were increased 73% and 27%, respectively under the treatment of poultry manure combination in comparison with control. In addition, the seed quality characters viz. germination, vigor index and total soluble protein content also exhibited significant improvement showing 23%, 44% and 17%, respectively by poultry manure + vermicompost + green manure. The above findings showed that to apply poultry manure + vermicompost + green manure as an effective soil amendment option and to obtain good yield and quality seed of wheat.

Introduction

In Bangladesh, wheat occupies 4% of the total cropped area and 11% of the cropped area in the *Rabi* season and contributes 7% to the total output of food cereals (BBS, 2008). In Bangladesh, the average yield of wheat is 3.16 t ha⁻¹ (BBS, 2017), which is below the achievable yield of 4.5 t ha⁻¹ (BARI, 2011). Alam *et al.* (2013) stated two reasons for the yield gap as (i) biotic factors including poor quality seeds and seedlings, insects, diseases, weeds and rodents; and (ii) abiotic factors including soil, nutrients and water. However, many reasons for this yield gap remain unexplained. Nutrients availability of soils is declining with time pace but which had been rich in the past. Among the different agricultural inputs, fertilizer is the most important one and nearly 50% of the modern agricultural production depends on this insert (Pradhan, 1992). The non-judicial and imbalanced use of chemical fertilizer generate hazards on soil, environment and human health which led the growers' considerable attention to turn organic manures application for sustainable production practices (Singh *et al.*, 2018). It is authentic that fairly good soil fertility and plant nutrients are important to farming, whether the practices are considered "conventional" or "sustainable" (Hue and Silva, 2000). Yadvinder-Singh *et al.* (2008) stated organic materials like crop residues, green manure, and animal manure show great influence on soil productivity. Therefore, probiotic and organic manures application should be evaluated on agricultural land reclamation and crop yield potentiality aspects in Bangladesh. In this view, the investigation was carried out to assess the effect of probiotic and organic manure amendment soil on the yield and seed quality of wheat.

Materials and Methods

The present piece of research work was conducted in the Institute of Biological Sciences (IBS) and Plant Pathology Laboratory, Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh during the period from July 2016 to June 2019. The experimental field is geographically situated at 24°17" N latitude and 88°28" E longitude at an elevation of 20 m above the sea level belonging to the Agro-ecological Zone-11(AEZ-11). The soil of the experimental field is characterized by poorly drained with moderate permeable, loamy and slightly alkaline (pH = 8.10) in nature. Before 2016, the field was occupied with barley production for two years. The experiment was laid out in randomized complete block design (RCBD) with three replications. Three wheat varieties viz. BARI Gom-28 (V_1), BARI Gom-29 (V_2) , BARI Gom-30 (V_3) and nine soil amendment treatments viz. control (T_0) , rice straw + vermicompost + green manure (T_1) , cow dung + vermicompost + green manure (T_2) , compost + vermicompost + green manure (T_3) , poultry manure + vermicompost + green manure (T_4) , Trichoderma harzianum + vermicompost + green manure (T_5) , mungbean residue + vermicompost + green manure (T_{7}), Trichoderma viride + vermicompost + green manure (T_{7}) and chemical fertilizer (T_8) were used . The unit plot was 5.0 m² having the plot to plot 0.5 m, bed to bed 0.25 cm distances and 1m from surrounding the boundary. Seeds of Dhaincha (Sesbania rostrata) were sown at the rate of 50 kg ha⁻¹to the respective plots and after 50 days of sowing young succulent green plants were incorporated into the soil. Crop residues (rice straw, mungbean residue), cow dung, compost and poultry litter were applied @ 10 t ha⁻¹ before 7 days of sowing. Vermicompost was applied @ 5 t ha⁻¹ and *Trichoderma* spp. suspension $(1 \ 10^6 \text{ cfu g}^{-1} @ 5 \text{ kg ha}^{-1})$ before sowing. In the case of chemical treatment, one-third of urea (200 kg ha⁻¹), TSP 160 kg ha⁻¹, MoP 45 kg ha⁻¹ and Gypsum 115 kg ha⁻¹ were used as basal dose (BARI, 2014). The rest of the urea was applied as two installments one at 21 DAS and another at 55 DAS.

Data recording

In the case of yield and yield attributes, ten spikes were selected randomly from each experimental plot for recording data on spike length, spikeletsspike⁻¹, fertile spikelets spike⁻¹, grains spike⁻¹ and grain weight spike⁻¹, while 1000-grain weight, grain yield and straw yield of crops were recorded which were collected from one square meter area at the center of each unit plot.

Harvest index (%)

The harvest index denotes the ratio of grain yield (economic yield) to biological yield and was calculated with the following formula:

Harvest index (%) = $\frac{\text{Grain yield}}{\text{Biological yield}} \times 100$

Here, Biological yield= Grain yield + straw yield.

Determination of seed germination

Four hundred seeds of each treatment were taken for germination test as recommended by ISTA (1985). Germination of seeds was collected regularly for up to 10 days. Percentage of germination was determined as follows:

Germination (%) = $\frac{\text{No. of germinated seedling}}{\text{No. of seed set for germination}} \times 100$

Vigor index

After setting seeds, the germination percentage was calculated from the final count. Vigor index was found out by using the following formula (Maguire, 1962):

 $Vigor index = \frac{No. of normal seedlings (First count)}{Days to first count} + - - - - - + + \frac{No. of normal seedlings (Final count)}{Days to final count}$

The total soluble protein content of seed

A technique formerly defined by Guy *et al.* (1992) was performed with some modification to estimate the total soluble protein of seed using a spectrophotometer.

Statistical analysis

The recorded data were compiled and tabulated for statistical analysis. The trial data for all parameters were analyzed statistically for analysis of variance (ANOVA) throughr-studio (<u>http://www.rstudio.com/</u>) of "Agricolae" package (<u>https://CRAN.R</u> <u>project.org/package=agricolae</u>) and Duncan's multiple range test (DMRT) was performed for comparing treatment means(Gomez and Gomez, 1984).

Results and Discussion

Effect of soil amendment on yield and yield attributes

Spike length

The spike length was affected significantly by various soil amendment treatments (Table 1). In the final year, it has been found that the treatment T_8 (chemical fertilizer) produced the maximum spike length (8.60 cm) which was followed by T_4 (poultry manure + vermicompost + green manure) (8.43 cm) while the smallest spike length recorded from T_0 (7.00 cm). It might be due to higher nutrient concentration and moisture retention of the soil amended with poultry manure combination treatment that led to the formation of longer spike length. This result was correlated with Ahmed *et al.* (2002). Nevertheless, green manure fortified poultry manure keeps the N balance which may enhanced the length of spike of wheat. The above statement was also supported by Nguyen *et al.* (1995).

Spikelets spike⁻¹ and fertile spikelets spike⁻¹

The spikelets spike⁻¹ is treated as an important element of wheat grain yield (Sabaghina *et al.*, 2014). From Table 1, it revealed that spikelets spike⁻¹ varied significantly from 13.80 to 17.97 where the maximum and the minimum value for this parameter were recorded from T_8 and T_0 .

Nitrogen-rich organic manures may have a positive effect on the growth and yield attributes of wheat where higher fertile spikelet spike⁻¹ from T_8 (17.24) and T_4 (17.00) but 13.22 was the lowest obtained from T_{0} . These findings were partially correlated with the observation of Siavoshi *et al.* (2011).

Grains spike⁻¹

Grains spike⁻¹ is significantly influenced by soil amendments. During the research period, a significantly higher number of grains spike⁻¹ was recorded from the treated plot in comparison with the control plot. The rank of treatments were $T8>T_4>T_5>T_7>T_3>T_6>T_2>T_1>T_0$ (Table 2). The maximum number of grains spike⁻¹ (45.73) was recorded from T_8 which was statistically similar to T_4 (44.80) and the least value from T_0 (33.67). Considering the organic soil amendment treatments, poultry manure (T_4) performed the better response; such result might be due greater amount of N within it, which produced more dry matter and partitioning it towards grains which enhanced the grains spike⁻¹. This result was confirmatory with the outcomes of Shah *et al.* (2010) and Mukhtiar *et al.* (2018).

Treatments	S	pike lengt (cm)	th	Spi	ikeletsspik (no.)	ke-1	Fertile	spikelets (no.)	spike ⁻¹
	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019
T ₀	7.31e ± 0.16	$\begin{array}{c} 7.12 f \\ \pm \ 0.10 \end{array}$	7.00e ± 0.05	$\begin{array}{c} 14.66 \text{d} \\ \pm \ 0.21 \end{array}$	$\begin{array}{c} 14.20d \\ \pm \ 0.20 \end{array}$	$\begin{array}{c} 13.80d\\ \pm\ 0.15\end{array}$	$\begin{array}{c} 14.00g\\ \pm\ 0.20\end{array}$	$\begin{array}{c} 13.78 \text{f} \\ \pm \ 0.17 \end{array}$	$\begin{array}{c} 13.22 \mathrm{f} \\ \pm \ 0.19 \end{array}$
T_1	7.45de ± 0.11	7.55e ± 0.11	$\begin{array}{c} 7.63d \\ \pm \ 0.08 \end{array}$	15.33c ± 0.14	15.50c ± 0.26	15.77c ± 0.17	$\begin{array}{c} 14.74 f \\ \pm \ 0.22 \end{array}$	14.86e ± 0.13	15.08e ± 0.22
T ₂	7.62cde ±0.12	7.73de ± 0.11	7.84d ± 0.08	15.60c ± 0.12	15.86c ± 0.20	16.10c ± 0.14	15.20de ± 0.12	$\begin{array}{c} 15.32d\\ \pm \ 0.10\end{array}$	15.51d ± 0.13
T ₃	7.77bcd ±0.09	7.99bcd ± 0.12	8.18c ± 0.10	$\begin{array}{c} 16.10b\\ \pm\ 0.14\end{array}$	$\begin{array}{c} 16.40b\\ \pm\ 0.23\end{array}$	16.63b ± 0.16	$\begin{array}{c} 15.50 \text{cd} \\ \pm \ 0.14 \end{array}$	15.72c ± 0.12	15.93c ± 0.08
T ₄	8.12ab ± 0.12	8.29ab ± 0.11	8.43ab ± 0.08	17.16a ± 0.13	17.59a ± 0.17	17.81a ± 0.16	16.60a ± 0.16	16.78a ± 0.09	17.00a ± 0.13
T_5	7.91bc ± 0.10	8.12abc ± 0.07	8.25bc ± 0.06	16.43b ± 0.14	16.68b ± 0.16	16.88b ± 0.21	15.90b ± 0.16	16.16b ± 0.15	16.40b ± 0.17
T ₆	7.54cde ± 0.10	7.66de ± 0.08	7.77d ± 0.08	15.51c ± 0.15	15.78c ± 0.20	16.01c ± 0.17	14.90ef ± 0.18	15.10de ± 0.17	15.41de ± 0.15
T ₇	7.79bcd ± 0.12	7.95cd ± 0.09	8.08c ± 0.06	16.26b ± 0.19	16.46b ± 0.11	16.70b ± 0.19	15.62bc ± 0.15	15.89bc ± 0.13	16.12bc ± 0.16
T ₈	8.27a ± 0.08	8.41a ± 0.09	8.60a ± 0.10	17.36a ± 0.17	17.68a ± 0.16	17.97a ± 0.17	16.89a ± 0.13	17.06a ± 0.10	17.24a ± 0.18
LS	**	**	**	**	**	**	**	**	**
CV (%)	4.58	4.03	2.93	2.76	3.23	3.23	2.42	2.00	2.61

Table 1. Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat

In the column, mean values bearing similar letter(s) or without letter are similar and those having dissimilar letters are differed significantly as per Duncan's Multiple Range Test. $T_0 = \text{control}$, $T_1 = \text{rice straw} + \text{vermicompost} + \text{green manure}$, $T_2 = \text{cow dung} + \text{vermicompost} + \text{green manure}$, $T_3 = \text{compost} + \text{vermicompost} + \text{green manure}$, $T_4 = \text{poultry manure} + \text{vermicompost} + \text{green manure}$, $T_5 = T$. harzianum + vermicompost + green manure, $T_6 = \text{mungbean residues} + \text{vermicompost} + \text{green manure}$, $T_7 = T$. viride+ vermicompost + green manure, $T_8 = \text{chemical fertilizer}$, LS = Level of significance, **= 1% level of significance, CV = Coefficient of variation

Grains weight spike⁻¹

As a yield component, grains weight spike⁻¹ of wheat is an important factor that contributes to the grain yield (Peltonen-Sainio *et al.*, 2007). Grain weight spike⁻¹ at harvest varied significantly due to different soil amendment treatments (Table 2). Among the different soil amendment treatments, T_4 (2.14 g) performed better than other treatments. The rest of the treatments were also showed as descending by T_5 (2.06 g), T_7 (2.01 g), T_3 (1.99 g), T_2 (1.93 g), T_6 (1.89 g), T_1 (1.86 g), while the lowest one in T_0 (1.60 g). The advancement of grain weight spike⁻¹ that provided with poultry manure treatment combination might be due to balanced nutrient stock during the grain filling stage and improvement of soil fertility. A similar result was also reported by Garg and Bahla (2008).

1000-grain weight

For cereal crops, 1000-grain weight is an important factor to determine the yield and yield attributes (Metho *et al.*, 1998). The data of 1000-grain weight of wheat indicated that chemical fertilizer (T_8) and poultry manure combination (T_4) significantly increased the weight (54.12 g and 53.90 g) while plants that did not receive any soil amendments (T_0) showed the lowest (48.78 g) might be due to availability of nutrients throughout the growing period and improvement of soil characters. This outcome was agreed with Ahmed *et al.* (2017).

	Grai	ins spike ⁻¹	(no.)	Grain	weight spi	ike ⁻¹ (q)	1000)-grainwei	ght (g)
Treatments	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018-
	2017	2018	2019	2017	2018	2019	2017	2018	2019
т	36.88f	35.04e	33.67i	1.66g	1.62g	1.60g	49.52f	48.61e	48.78i
T ₀	± 0.30	± 0.30	± 0.87	± 0.03	± 0.02	± 0.04	± 0.44	± 0.51	± 0.41
T ₁	36.98f	38.42d	39.53h	1.74f	1.80f	1.86f	50.03f	51.05d	51.30h
11	± 0.22	± 0.23	± 0.88	± 0.02	± 0.03	± 0.04	± 0.37	± 0.30	± 0.31
T ₂	38.11e	39.36d	40.67f	1.80ef	1.88de	1.93e	51.19de	51.84bcd	51.95f
12	± 0.21	± 0.26	± 0.96	± 0.03	± 0.03	± 0.05	± 0.33	± 0.59	± 0.33
т	38.76d	40.51c	41.58e	1.83de	1.91d	1.99d	51.77cd	52.34a-d	52.49e
T ₃	± 0.21	± 0.39	± 1.02	± 0.02	± 0.03	± 0.04	± 0.19	±0.34	± 0.34
т	42.11b	43.50a	44.80b	1.98b	2.10b	2.14b	52.94ab	53.51ab	53.90b
T_4	± 0.40	± 0.59	± 1.13	± 0.02	± 0.02	± 0.05	±0.27	± 0.47	± 0.39
т	40.17c	41.59b	43.14c	1.93bc	2.02c	2.06c	52.27bc	52.98abc	53.19c
T ₅	± 0.22	± 0.43	± 1.08	± 0.03	± 0.02	± 0.05	±0.21	± 0.90	± 0.38
т	37.20f	38.71d	40.23g	1.77ef	1.84ef	1.89f	50.48ef	51.39cd	51.60g
T ₆	± 0.20	± 0.27	± 0.97	± 0.03	± 0.03	± 0.05	± 0.40	± 0.31	± 0.34
т	39.73c	40.83bc	42.50d	1.89cd	1.99c	2.01d	51.97bcd	52.51a-d	52.64d
T ₇	± 0.25	± 0.42	± 1.07	± 0.02	± 0.02	± 0.05	±0.14	± 0.26	± 0.36
Τ.	42.74a	44.41a	45.73a	2.07a	2.15a	2.20a	53.28a	53.81a	54.12a
T ₈	± 0.37	± 0.53	± 1.04	± 0.03	± 0.03	± 0.05	± 0.30	± 0.29	± 0.40
LS	**	**	**	**	**	**	**	**	**
CV (%)	1.52	2.55	1.09	3.69	2.90	1.48	1.90	3.12	1.23

Table 2. Effect of organic and probiotic soil amendments on yield and yield contributing characters of wheat


In the column, mean values bearing similar letter(s) or without letter are similar and those having dissimilar letters are differed significantly as per Duncan's Multiple Range Test. $T_0 = \text{control}$, $T_1 = \text{rice straw} + \text{vermicompost} + \text{green manure}$, $T_2 = \text{cow dung} + \text{vermicompost} + \text{green manure}$, $T_3 = \text{compost} + \text{vermicompost} + \text{green manure}$, $T_4 = \text{poultry manure} + \text{vermicompost} + \text{green manure}$, $T_5 = T$. harzianum + vermicompost + green manure, $T_6 = \text{mungbean residues} + \text{vermicompost} + \text{green manure}$, $T_7 = T$. viride+ vermicompost + green manure, $T_8 = \text{chemical fertilizer}$, LS = Level of significance, **= 1% level of significance, CV = Coefficient of variation


Grain yield

Different yield component specifically spikesm⁻², number and weight of grains spike⁻¹ and 1000grain weight mathematically functioned on yield of cereal crop (Thiry *et al.*, 2002). Application of different soil amendments showed significant variation of grain yield of wheat (Figure 1, 2 and 3). In the last year, influences of different treatments on grain yield were found to be ranked in that following descending order T₈ (4.25 t ha⁻¹) > T₄ (4.09 t ha⁻¹) > T5 (3.85 t ha⁻¹) > T₇ (3.64 t ha⁻¹) > T₃ (3.60 t ha⁻¹) > T₂ (3.34 t ha⁻¹) > T₆ (3.27 t ha⁻¹) > T₁ (3.05 t ha⁻¹) > T₀ (2.36 t ha⁻¹). Better performance of T₄ might be enhanced soil fertility and improvement of physical conditions, reduction of nutrient losses which collectively contributed to the higher grain yield. This result was in confirmed to that with the observation of Zhang *et al.* (2016).

Straw yield

Soil amendments significantly influenced the straw yield (Figure 1, 2 and 3). Among the different treatments, T_8 (5.40 t ha⁻¹) performed superior. The rest of the treatments were also showed significant result ranked as descending by T_4 (5.31 t ha⁻¹), T_5 (5.14 t ha⁻¹), T_7 (5.05 t ha⁻¹), T_3 (4.93 t ha⁻¹), T_6 (4.83 t ha⁻¹), T_2 (4.73 t ha⁻¹) and T_1 (4.60 t ha⁻¹), while the lowest one in T_0 (4.18 t ha⁻¹). This performance of organic manures might be due to the capability of abundant nutrients supply and the creation of a favorable growth environment. A similar result was also noted by Auti *et al.* (1999) and Kumar and Abraham (2018).








Fig. 2. Effect of organic and probiotic soil amendments on grain yield and straw yield of wheat in 2017-2018.




Fig. 3. Effect of organic and probiotic soil amendments on grain yield and straw yield of wheat in 2018-2019.

Harvest index

Treatment T₈ showed a maximum harvest index (44.05%) followed by T₄ (43.48%) and the lowest one (36.13%) derived from T₀ during 2018-2019 (Figure 4). The boosting harvest index might be the continued supply of nutrients and higher dry matter distribution towards grain formation provided by the poultry manure treatment combination. This result was in agreement with the investigation of Kabesh *et al.* (2009) and Khan *et al.* (2018).




Fig. 4. Effect of organic and probiotic soil amendments on harvest index of wheat in 2016-2017, 2017-2018 and 2018-2019.

Effect of soil amendments on seed quality

After every harvesting, seeds were collected separately from each treatment and preserved. Major seed quality characters viz. germination, vigor index and total soluble protein content were determined. The introduction of designed soil amendments not only developed the growth and yield of wheat but also enriched parameters of seed quality.

Germination (%)

Soil amendments influenced the germination of seed. The treatment T_4 significantly enhanced (92.08%) the germination followed by T_5 (88.89%) and T_0 had a lower effect (74.78%) on germination in 2018-2019 (Table 3). A similar trend was also observed in the previous two cropping season. This might be due to an adequate supply of nutrients which helps to build up carbohydrate-like seed elements thus producing vigorous seeds. The outcome of the investigation was the conformity of previous results of Gowda *et al.* (2010) and Aslam *et al.* (2011).

Turn tar and	Ge	rmination	(%)	V	ligor inde	x	Soluble	protein co g ⁻¹ FW)	ntent(mg
Treatments -	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019	2016- 2017	2017- 2018	2018- 2019
T ₀	70.50h	72.53g	74.78g	24.70h	27.49f	28.84f	77.31e	77.47e	79.10e
10	± 0.92	± 0.93	± 0.85	± 0.42	± 0.40	± 0.27	± 0.28	± 0.58	± 0.42
T ₁	75.31g	77.54f	79.67f	28.20g	32.52e	34.39e	78.20e	79.68d	79.58e
11	± 1.08	± 1.02	± 1.18	± 0.55	± 0.54	± 0.36	±0.45	± 1.05	± 0.52
T_2	78.14f	80.88de	83.67d	31.69e	34.42d	35.93d	80.25d	80.97cd	82.98d
12	±0.82	± 0.98	± 0.96	± 0.85	± 0.96	± 0.38	±0.46	± 0.73	± 0.39
т	81.08d	83.86bc	86.17c	32.60d	36.84b	38.36b	80.20d	81.98c	82.24d
T ₃	± 0.85	± 0.92	±0.94	±0.43	± 0.49	± 0.34	± 1.02	± 1.44	± 1.28
т	88.27a	89.93a	92.08a	35.99a	39.48a	41.54a	88.17b	93.30a	92.56b
T_4	± 1.05	± 1.01	± 1.24	± 0.73	± 0.84	± 0.58	± 1.89	± 0.90	± 0.70
т	84.75b	86.52b	88.89b	34.57b	37.30b	39.26b	87.59b	88.30b	91.03c
T_5	± 1.06	± 1.15	±0.84	± 0.52	± 0.58	± 0.46	± 0.59	± 0.66	± 0.99
т	78.76e	78.40ef	81.83e	30.21f	34.68d	35.37de	81.51c	81.31c	82.62d
T ₆	± 0.89	± 1.46	± 1.42	±0.46	± 0.33	± 0.33	±0.72	± 0.63	±0.70
т	83.59c	83.33cd	86.28c	33.56c	35.78c	36.98c	87.88b	89.17b	91.99bc
T ₇	± 0.93	± 1.05	±0.96	±0.43	± 0.38	±0.42	±0.58	±0.83	±0.67
т	81.37d	84.57bc	86.76c	32.46d	34.03d	37.27c	93.32a	93.80a	96.43a
T ₈	± 0.86	± 1.05	± 01.16	± 0.50	± 0.34	±0.48	± 1.08	± 1.20	± 1.14
LS	**	**	**	**	**	**	**	**	**
CV (%)	0.82	3.34	1.79	0.97	3.16	2.91	1.58	1.61	1.42

Table 3. Effect of organic and probiotic soil amendments on seed quality of wheat

In the column, mean values bearing similar letter(s) or without letter are similar and those having dissimilar letters are differed significantly as per Duncan's Multiple Range Test. $T_0 = \text{control}$, $T_1 = \text{rice straw} + \text{vermicompost} + \text{green manure}$, $T_2 = \text{cow dung} + \text{vermicompost} + \text{green manure}$, $T_3 = \text{compost} + \text{vermicompost} + \text{green manure}$, $T_4 = \text{poultry manure} + \text{vermicompost} + \text{green manure}$, $T_5 = T$. *harzianum* + vermicompost + green manure, $T_6 = \text{mungbean residues} + \text{vermicompost} + \text{green manure}$, $T_7 = T$. *viride*+ vermicompost + green manure, $T_8 = \text{chemical fertilizer}$, LS = Level of significance, **= 1% level of significance, CV = Coefficient of variation

Vigor index

The vigor index is an imperative quality of seed that indicates the potentiality of constant emergence of seedlings (Pradeep, 2018). After three years of experimentation, considering the comparable trend of vigor index from respective treatments, poultry manure (41.54) and *Trichoderma harzianum* (39.26) in comparison with control treatment (28.84). The speed of germination i.e. vigor index might be significantly different for providing the substantial macro and micro nutrient in the grain filling stage and carbohydrate stored in seed enhances the quality. This result was in agreement with the reflection of Farhad *et al.* (2011).

Total soluble protein content

Among the major cereals, wheat is the prominent source of vegetal protein (13%). The results in general T_8 produced higher protein (96.43 mg g⁻¹ FW) followed by T_4 (92.56 mg g⁻¹ FW) and the least at T_0 (79.10 mg g⁻¹ FW). The increase of protein content might be due to the availability of high NPK that satisfy the plant requirement and accumulated N in leaf mobilized to seed components which leads the protein production. This result was in corroborates with the findings of Abedi *et al.* (2010) and Rasul *et al.* (2015). Considering the fact, the poultry manure combination treatment (T_4) was an environmentally friendly alternative to chemical fertilizer (T_8) for increasing soil fertility and crop productivity.

Conclusion

There is a potentiality for sustainable improvement of wheat growth, yield and seed quality with organic soil amendment through probiotic and organic manure. In that term, the inclusion of green manure, vermicompost and poultry manure might be a worthy alternative. Organic manures particularly poultry manure was a healthier source for improvement of growth, yield parameters and seed quality of wheat as compared to other manures. So, the poultry manures could be suggested for soil amendment and increase wheat production in the studied region.

References

- Abedi, T., A. Alemzadeh and S.A. Kazemeini. 2010. Effect of organic and inorganic fertilizers on grain yield and protein banding pattern of wheat. Aust. J. Crop Sci. 4:384-389.
- Ahmed, B.E.A.M., A.A. Ishag, M.K. Hassan and M.O. Ahmed. 2017. Response of two wheat cultivars (*Triticum aestivum* L.) to amended nitrogen fertilizer on yield and grain quality in Halfa Elgadidah area. MOJ. Biol. Med. 1(5): 133-136.
- Ahmed, S.I., M.K. Abbasi and G. Rasool. 2002. Integrated plant nutrition system (IPNS) in wheat under rain fed conditions of Rawalakot Azad Jammu and Kashmir [Pakistan]. Pak. J. Soil Sci. 21(1&2): 79-86.
- Alam, M.M., M.R. Karim and J.K. Ladha. 2013. Integrating best management practices for rice with farmers' crop management techniques: A potential option for minimizing rice yield gap. Field Crops Res. 144: 62-68.
- Aslam, M., M.A. Khan, I.U. Awan, E.A. Khan, A.A. Khan and G. Jilani. 2011. Effect of single and combined use of various organic amendments on wheat grown over green manured soil: growth and yield attributes. Pak. J. Nutr. 7: 640-646.
- Auti, A.K., S.C. Wadile and V.S. Pawar. 1999. Yield, quality and nutrient removal of wheat (*Triticum aestivum* L.) as influenced by levels and sources of fertilizer. Indian J. Agron.44: 119-120.
- BARI (Bangladesh Agricultural Research Institute). 2014. Krishi Projukti Hatboi (Handbook on Agrotechnology), 6th edition, Gazipur, Bangladesh.

- BARI (Bangladesh Agricultural Research Institute).2011. Krishi Projukti Hatboi (Handbook on Agrotechnology), 5th edition, Gazipur, Bangladesh.
- BBS (Bangladesh Bureau of Statistics). 2008. Statistical Yearbook of Bangladesh. Statistics and Informatics Div., Ministry of Planning. Govt. People's Repub., Bangladesh, Dhaka.
- BBS (Bangladesh Bureau of Statistics).2017.Statistical Yearbook of Bangladesh. Statistics and Informatics Div., Ministry of Planning. Govt. People's Repub., Bangladesh, Dhaka.
- Farhad. W, M.F. Saleem, M.A. Cheema, H.Z. Khan and H.M. Hammad. 2011. Influence of poultry manure on the yield and quality of spring maize. Crop Environ. 2: 6-10.
- Garg, S. and G.S. Bahla. 2008. Phosphorus availability to maize as influenced by organic manures and fertilizer P associated phosphatase activity in soils. Bioresour. Technol. 99: 5773-5777.
- Gomez. K.A. and A.A. Gomez.1984. Statistical Procedures for Agricultural Research (2nd edition). International Rice Research Institute. John Wiley and Sons, a Wiley-Inter Science Publication.
- Gowda, C., P.N.K. Biradar, B.N. Patil, J.S. Awakknavar and R. Hunje. 2010. Effect of organic manures on growth, seed yield and seed quality of wheat. Karnataka J. Agric. Sci. 21: 366-368.
- Guy, C., D. Haskell, L. Neven, P. Klein and C. Smelser. 1992. Hydration-state-responsive proteins link cold and drought stress in spinach. Planta. 188: 265-270.
- Hajieghrari, B. and M. Mohammadi. 2016. Growth-promoting activity of indigenous *Trichoderma* isolates on wheat seed germination, seedling growth and yield. Aust. J. Crop Sci. 10: 1339-1347.
- Hue, N.V. and J.A. Silva. 2000. Organic Soil Amendments for Sustainable Agriculture: Organic Sources of Nitrogen, Phosphorus and Potassium: Plant nutrient management in Hawaii's soils, approaches for tropical and subtropical agriculture. College of Tropical Agriculture and Human Resources, University of Hawaii, Manoa. 133-144.
- ISTA (International Seed Testing Association). 1985. International rules for seed testing. Seed Sci. Technol. 13: 299-513.
- Kabesh, M.O., M.F. El-Kramany, G.A. Sary, H.M. El-Naggar and S.H. Gehan. 2009. Effects of sowing methods and some bio-organic fertilization treatments on yield and yield components of wheat. Res. J. Agric. Biol. Sci. 5: 97-102.
- Khan, T.U., M.T. Jan, A. Khan, G. Ahmad, M. Ishaq, K. Afridi, A. Murad, M.A. Qureshi, I. Ahmad and M. Seed. 2018. Integrated management of fertilizer nitrogen and poultry manure enhances wheat production. Pak. J. Agric. Res. 31: 207-215.
- Kumar, S. and T. Abraham. 2018. Productivity potential of wheat under certified organic production system. Int. J. Curr. Microbiol. Appl. Sci. 7: 281-288.
- Maguire, J.D. 1962. Speed of germination-aid in selection and evaluation for seedling emergence and vigor. Crop Sci. 2: 176-177.
- Metho, L.A., P.S. Hammes and E.A. Beyers. 1998. The effect of soil fertility on the contribution of main stem, tillers and kernel position to grain yield and grain protein content of wheat. S. Afr. J. Plant Soil 15: 53-60.
- Mukhtiar, A., A. Waqar, M.K. Khalil, M. Tariq, S. Muhammad, A. Hussain and A. Kamal. 2018. Evaluating the potential organic manure for improving wheat yield and quality under agroclimatic conditions of Pakistan. Adv. Crop Sci. Tech. 6: 1-4.
- Nguyen, M.L., R.J. Haynes and K.M. Goh. 1995. Nutrient budgets and status in three pairs of conventional and alternative and mixed cropping farms in Canterbury, New Zealand. Agric. Ecosyst. Environ. 52: 149-162.

- Peltonen-sainio, P., A. Kangas, Y. Salo and L. Jauhiainen. 2007. Grain numbers dominate grain weight in temperate cereal yield determination: evidence based on 30 years of multi-location trials. Field Crops Res. 100: 179-188.
- Pradeep, P. 2018. Seed quality parameters (Germination percentage and seedling vigor index) of rabi sorghum seeds influenced by rice weevil infestation. MOJ. Toxicol. 4: 391–396.
- Pradhan, S.B. 1992. Status of fertilizer use in developing countries of Asia and the Pacific Region In: Proc. Reg. FADINAP Sem. Chiang Mai, Thailand: 37-47.
- Rasul, G.A.M., S.T. Ahmed and M.Q. Ahmed. 2015. Influence of different organic fertilizers on growth and yield of wheat. American-Eurasian J. Agric. Environ. Sci. 15: 1123-1126.
- Sabaghina, N., M. Janmohammadi and A.E. Segherloo. 2014. Evaluation of some agromorphological traits diversity in Iranian bread wheat genotypes. Annales UMCS, Biologia 69: 79-92.
- Shah, S.A., S.M. Shah, W. Mohammad, M. Shafi, H. Nawaz, S. Shehzadi and M. Amir. 2010. Effect of integrated use of organic and inorganic nitrogen sources on wheat yield. Sarhad J. Agric. 26: 559-563.
- Shewry, P.R., N.G. Halford, P.S. Belton and A.S. Tatham. 2002. The structure and properties of gluten: An elastic protein from wheat grain. Philosophical Transactions of the Royal Society of London Series B: Biol. Sci. 357: 133-142.
- Siavoshi, M.,A. Nasiri and S.L. Laware. 2011. Effect of organic fertilizer on growth and yield components in rice (*Oryza sativa* L.). J. Agric. Sci. 3:217.224.
- Singh, A., M.S. Darhvankar and G. Singh. 2018. Impact of organic and inorganic amendments on yield and growth of wheat (*Triticum aestivum* L.). Int. J. Curr. Microbiol. Appl. Sci. 7: 789-794.
- Thiry, D.E., R.G. Sears, J.P. Shroyer and G.M. Paulsen. 2002. Relationship between tillering and grain yield of Kansas wheat varieties. Kansas State University Agricultural Experimental Station and Cooperative Extension Service: 1-5.
- Yadvinder-Singh, S. Bijay, R.K. Gupta, J. Ladha, J.S. Bains and J. Singh. 2008. Evaluation of press mud cake as a source of nitrogen and phosphorus for the rice-wheat cropping system in the Indo-Gangetic plains of India. Biol. Fertil. Soils. 45: 289-296.
- Zeidan, M.S. and M.F.E. Kramany. 2001. Effect of organic manure and slow-release N Fertilizer on the productivity of wheat (*Triticum aestivum* L.) in sandy soil. Acta Agronomica Hungarica. 49: 379-385.
- Zhang, H.Q., X.Y. Yu, B.N. Zhai, Z.Y. Jin and Z.H. Wang. 2016. Effect of manure under different nitrogen application rates on winter wheat production and soil fertility in dry land. IOP Conf. Ser.: Earth Environ. Sci. 39(1):1-10.



J. Bio-Sci. 29(2): 61-70, 2021 (December) http://www.banglajol.info/index.php/JBS/index DOI: https://doi.org/10.3329/jbs.v29i2.54955

EFFECTS OF PROBIOTIC AND ORGANIC MANURES FOR SOIL AMENDMENTS ON SEEDLING SURVIVABILITY AND LEAF CHLOROPHYLL FOR CROP IMPROVEMENT IN WHEAT (*TRITICUM AESTIVUM* L.)

MA Hossain^{1, 2}, MM Hasan³ and SMS Islam^{1*}

¹Plant Biotechnology and Genetic Engineering Laboratory, Institute of Biological Sciences, University of Rajshahi, Rajshahi-6205, Bangladesh ²Directorate of Secondary and Higher Education, Dhaka-1000, Bangladesh

³Agronomy and Agricultural Extension Department, University of Rajshahi, Rajshahi-6205, Bangladesh

Abstract

Under this study a total of nine treatments were considered for soil amendments where organic manures, chemical fertilizer and probiotics were used to evaluate the efficiency on growth and yield of wheat. On the basis of seedling disease incidence and chlorophyll presence parameters data were recorded on seedling infection at 14 days, seedling blight at 21 days, and chlorophyll content at 70 days after sowing. In this study the same experiments were conducted up to three years (2016-2019) and data were compiled and analyzed. It was observed that the disease prevalence attributes significantly decreased in T₅ (*Trichoderma harzianum* + vermicompost + green manuring) than other treatment combinations and control (T₀). Noteworthy results were found on seedling health, application of probiotic and organic manures which significantly influenced the less infection of wheat. Besides this, chlorophyll content of wheat leaf and plant number with effective tiller enhanced largely under T₈ (chemical fertilizer) than T₄ (poultry manure + vermicompost + green manuring) and control (T₀). The main advantage of this research is to evaluate efficiency for growing healthy wheat plant which reflects on plant growth and yield and ultimately helps for food security.

Key words: Blight, Chlorophyll, Green and poultry manure, Infection, Vermicompost, Wheat

Introduction

Wheat (*Triticum aestivum* L.) is a staple food of billions of people all over the world and third most agricultural commodity next to rice and maize. The contribution of wheat is being increased significantly in different countries as Nigeria (1% - 6.64%), India (11.85% - 20.41%) and China (12.20% - 17.83%) regarding total kcal (Shewry and Hey 2015). In Bangladesh, wheat is one of the important grain crops regarding economic and consumption purposes. Wheat is used as industrial and commercial products for human consumption having high protein with others nutrients that is also a cheaper source of feed for livestock (Hammad et al. 2011). Land engaged for cultivation decreased with time to meet up the accommodation demand for geometric growth of population. To face the land limitation problem and provide food for increased population, sustainable use of land is burning issue for agricultural development (Haque and Islam 2018, Alam et al. 2020). The substantial pressure on land may create various constraints due to imbalanced and inadequate

ISSN 1023-8654

^{*}Author for correspondence: shahinul68@gmail.com

application of fertilizer (Kumar and Abraham 2018). Chemical fertilizer may increase the wheat yield but prolonged application resulted human health hazard and imbalanced natural resources (Hossain et al. 2021a). So, need to link soil management and cropping design that are more environmentally sustainable (Karnel et al. 1994). Though transition period of three years produce lower performance but higher soil fertility found over time in conventional to organic farming (Tester 1990). So, for achieving higher yield and retain soil health sound organic amendments are felt to be important. A good soil should have organic matter content of more than 2.5%. Most of the soils in Bangladesh have low to very low organic matter content, generally less than 1.5% (BARC 2012). Soil productivity is greatly influenced by the application organic manures like crop residues, green manure and animal manure in the soil-plant system (Yadvinder-Singh et al. 2008, Hossain et al. 2021a). Probiotics are live microorganisms which confer a beneficial health benefit on the host. Trichoderma is considered as a probiotic having the characteristic of fungi and considered as a potential and promising bio-control agent and growth promoter for many crops. Bio-fertilizer is regarded as an eco-friendly substitute of chemical fertilizer trends to enhance soil fertility and increase crop productivity and yield without creating harm to the environment (Hermosa et al. 2012, Hossain et al. 2021b). It was established that Trichoderma is a biological degrader and promotes plant disease defense, increasing the immunity of the plant. Hence, Trichoderma have tremendous opportunities for disease management of soil borne pathogens but also have the capacity to improve plant growth parameters and soil health. It is authentic that fairly good soil fertility and plant nutrients are important to farming. Hence to improve soil fertility and increase plant nutrients availability, efforts need to be made to increase soil organic matter content. So, in view of the above realities an initiative was undertaken to focus the probiotic and organic manure performance on wheat seedling and plant growth resulted on yield, which may help to increase the productivity of wheat.

Materials and Methods

Soil and field preparation

The present research work has been carried out in the Experimental Field of the Institute of Biological Sciences (IBSc) and laboratory works has been carried out at Plant Biotechnology and Genetic Engineering Lab., IBSc and Plant Pathology Lab., Department of Agronomy and Agricultural Extension, University of Rajshahi, Bangladesh during the period from July 2016 to June 2019. The experimental field geographically situated at 24°17` N latitude and 88°28` E longitude at an elevation of 20 m above the sea level belonging to the Agro-Ecological Zone-11 (AEZ-11). The soil of the experimental field has characteristics like poorly drained with moderately permeability, loam and slightly alkaline (pH = 8.10) in nature. The physical properties of the experimental soil had moisture content 19.40%, particle density 2.65 g/cc, bulk density 1.27 g/cc and porosity 51.34%. Furthermore, chemical properties like the organic matter content 1.20%, K 0.150 cmol (+)/kg, total N 0.07%, P 26.30 μ g/g, S 12.50 μ g/g, Zn 0.75 μ g/g, EC 145 μ s/cm and C:N 10:1 were recorded.

Experimental design and treatments

The experiment was laid out in randomized complete block design (RCBD) with three replications. Three wheat varieties *viz*. BARI wheat-28 (V₁), BARI wheat-29 (V₂), BARI wheat-30 (V₃) and nine treatments (T) including control (T_0), rice straw + vermicompost + green manuring (T_1), cow dung + vermicompost + green manuring (T_2), compost + vermicompost + green manuring (T_3), poultry manure + vermicompost + green

manuring (T₄), *Trichoderma harzianum* + vermicompost + green manuring (T₅), mung bean residue + vermicompost + green manuring (T₆), *Trichoderma viride* + vermicompost + green manuring (T₇) and chemical fertilizer (T₈) were used in this study. The unit plot was 5.0 m² having a plot to plot 0.5 m, bed to bed 0.25 cm distance and 1 m from surrounding the boundary. The total unit plot was 81. Seeds of *Dhaincha* (*Sesbania rostrata*) were sown at the rate of 50 kg/ha (BARI 2014) and after 50 days young succulent green manure plants incorporated into the soil to the respective plots (Dubey et al. 2015). Crop residues (rice straw and mung bean), cow dung, compost and poultry litter were applied at the rate 10 t/ha before 7 days of sowing. Vermicompost was applied 5 t/ha and *Trichoderma* spp. suspension (1 × 10⁶ cfu/g at the rate 5 kg/ha) before sowing. In case of chemical treatment, one third urea (200 kg/ha), TSP 160 kg/ha, MP 45 kg/ha and gypsum 115 kg/ha was used as basal dose (BARI 2014). Rest of the urea was applied following two installments, one at 21 days after sowing (DAS) and rest one at 55 DAS.

Data collection on seedling health, leaf chlorophyll content and plant characters

Seedling infection

The seedlings which were found yellow and rotted at the base are considered as infected. From germination up to 14 days regular observation was made and infected seedlings were counted in each plot. Infection was estimated by the following formula:

% Seedling infection =
$$\frac{\text{Number of infected seedlings}}{\text{Total number of seedlings}} \times 100$$

Seedling blight

Seedlings which were dead and became straw in color defined as seedling blight. Seedling infection is treated as the first step of seedling blight. Thus, blighted seedlings were counted at 21 DAS in every plot. It was estimated as by the following formula:

% Seedling blight = $\frac{\text{Number of blighted seedlings}}{\text{Total number of seedlings}} \times 100$

Leaf chlorophyll content

The atLEAF handled chlorophyll meter was used to assess leaf greenness. The meter determines light transmittance through the leaf at 660 and 940 nm wavelengths. The readings were obtained by inserting the middle portion of the topmost fully expanded leaf in the slit of the meter. At least three readings from leaves of randomly selected plants in each plot were recorded and mean value was determined. Abnormally looking or insect attacked plants were not selected for measurement (Ali et al. 2020).

Total plant/m²

Before harvesting 1 m² area was selected with quadrate and the number of total plants was counted. This operation was done in each plot preferably choosing the five rows in the middle of the plot.

Effective tiller/plant

The numbers of tillers bearing panicles were counted at harvest with the help of quadrate from 1 m × 1 m area, which expressed as effective tillers/plant.

Statistical analysis

The recorded data were compiled and tabulated for statistical analysis. The data were analyzed statistically for analysis of variance (ANOVA) and Duncan's Multiple Range Test (DMRT) through r-studio by the "Agricolae" package.

Results

Seedling infection

Regarding various treatments effect, the lower rate of seedling infection (0.32%) was found after application *Trichoderma harzianum* (T₅) in comparison with the maximum infection from T₀ (2.32%) in 2018-2019. All the treatments in the experimental period presented significant differences among each other. But T₁ (rice straw + vermicompost + green manuring) and T₈ (chemical fertilizer) was statistically identical. Reduction of seedling infection at 14 DAS under different soil amendments showed more or less the following order as T₅>T₇>T₄>T₃>T₂>T₆>T₁>T₈>T₀ during the whole research period shown in Table 1.

Seedling blight

Under this study, recorded results showed that there was a significant variation among the treatments. The highest inhibition effect against seedling blight (0.87%) was observed by treatment T_5 and the lowest (4.74%) from control (T_0) during 2018-2019 (Fig.1 a-b). Next to treatment T_5 , the treatment T_7 also showed remarkable effect in reducing seedling blight (1.07%) which was statistically identical (1.09%) with T_4 (Table 1).




Fig. 1(a-b): Seedling status and plant growth by adding probiotic and organic manures for soil amendments, a) seedling infection and, b) seedling blight.

Leaf chlorophyll content

A significant difference was observed from various soil amendments for the chlorophyll content of wheat leaf at 70 DAS. The highest value of chlorophyll content (51.48 SPAD) was recorded from chemical fertilizer (T₈) and which was followed by T₄ (48.37 SPAD) and T₅ (48.08). In this regard, the smallest value (40.33 SPAD) was noticed from T₀ during the season 2018-2019. Due to soil amendment with chemical fertilizer (T₈)

provided more chlorophyll content and the trend was maintained from beginning to the end of the study (Table 1).

	Seedling infection (%)			Seedling blight (%)			Leaf chlorophyll content (SPAD)		
Treat.	14 DAS			21 DAS			70 DAS		
	2016-	2017-	2018-	2016-	2017-	2018-	2016-	2017-	2018 -
	2017	2018	2019	2017	2018	2019	2017	2018	2019
T ₀	2.14	1.99	2.32	3.52	3.30	4.74	38.26	38.82	40.33
	± 0.12a	± 0.09a	± 0.13a	± 0.16a	± 0.16a	± 0.20a	± 0.85h	± 0.90g	± 0.79d
T ₁	1.49	1.47	1.46	2.53	2.41	2.48	40.18	41.62	42.77
	± 0.03b	± 0.05b	± 0.09b	± 0.08b	± 0.08b	± 0.19c	± 0.90g	± 0.86f	± 0.87c
T ₂	1.34	1.24	1.20	2.17	2.04	2.11	41.39	43.56	42.96
	± 0.04d	± 0.05c	± 0.04c	± 0.06c	± 0.04d	± 0.08d	± 0.87f	± 1.05de	± 0.76c
T ₃	0.93	0.87	0.92	1.77	1.71	1.61	42.45	43.85	44.09
	± 0.04e	± 0.03d	± 0.05d	± 0.05d	± 0.05e	± 0.04e	± 0.85d	± 1.04d	± 0.85c
T ₄	0.59	0.58	0.55	1.24	1.19	1.09	44.59	46.63	48.37
	± 0.02f	± 0.02e	± 0.01e	± 0.05e	± 0.06f	± 0.05f	± 0.81b	± 0.89b	± 0.80b
T ₅	0.35	0.34	0.32	0.93	0.87	0.87	43.03	44.73	48.08
	± 0.02g	± 0.02g	± 0.02g	± 0.06f	± 0.04g	± 0.05g	± 0.85c	± 0.91c	± 0.61b
T ₆	1.43	1.44	1.42	2.44	2.33	2.16	41.08	41.80	43.06
	± 0.02c	± 0.06b	± 0.08b	± 0.07b	± 0.08bc	± 0.11d	± 0.89f	± 0.92f	± 0.63c
T 7	0.57	0.46	0.45	1.29	1.11	1.07	41.78	43.45	45.11
	± 0.02f	± 0.02f	± 0.03f	± 0.04e	± 0.03f	± 0.06f	± 0.92e	± 0.95e	± 0.71c
T ₈	1.48	1.47	1.42	2.44	2.26	2.57	47.14	48.92	51.48
	± 0.10c	± 0.08b	± 0.04b	± 0.09b	± 0.04c	± 0.11b	± 0.97a	± 1.08a	± 0.70a
LS	**	**	**	**	**	**	**	**	**
CV (%)	5.42	6.89	4.38	5.59	6.16	4.30	0.80	0.79	5.09

Table 1. Effect of organic and probiotic soil amendments on seedling infection and seedling blight of wheat

Each value represents the average of three replicates. In the column, mean values bearing similar letter(s) or without letter are identical and those having dissimilar letters are differed significantly as per Duncan's Multiple Range Test. T_0 = control, T_1 = rice straw + vermicompost + green manuring, T_2 = cow dung + vermicompost + green manuring, T_3 = compost + vermicompost + green manuring, T_4 = poultry manure + vermicompost + green manuring, T_5 = *T. harzianum* + vermicompost + green manuring, T_6 = mung bean residues + vermicompost + green manuring, T_7 = *T. viride* + vermicompost + green manuring, T_8 = chemical fertilizer, SPAD = The soil plant analysis development. LS = level of significance, ** = 1% level of significance, CV = co-efficient of variation.

Total plant/m²

Total plant/m² at harvest varied significantly due to different soil amendments all over the study period. The highest frequency of total plant/m² was found in T₈ (199.11%) and the lowest result was recorded in T₀ (158.67%) in 2016-2017 (Fig. 2). The next highest value was recorded from T₄ (191.56%) and T₅ (185.00%) had the statistical likeness. In the year 2017-2018, total plant/m² exhibited the matching trend due to various soil amendments as reported in 2016-2017. Where the highest (207.78%) plant/m² produced by T₈ and had similar identity with the second most from T₄ (202.89%). The lowest total plant/m² (163.89%) was noticed in T₀ coupled with T₁ (170.22%). Among the different soil amendments T₈ (216.11%) performed superior results besides T₄ (209.89%) to other treatments in 2018-2019. Rest of the treatments were showed also significant difference ranked as descending by T₅ (199.78%), T₇ (190.22%), T₃ (187.22%), T₂ (179.00%), T₆ (177.67%) and T₁ (173.44%), whereas the lowest one in T₀ (159.00%).

Effective tiller/plant

Under this study, recorded results on effective tiller/plant showed significant difference by the influence of treatments. Out of different soil amendments, chemical fertilizer (T₈) and poultry manure (T₄) performed as superior to the rest of the treatments mainly unamend (T₀) all over the research period (Fig. 3). On the basis of the above statement, T₈ was produced the maximum number of effective tiller/plant (3.49%) followed by T₄ (3.40%), whereas the minimum value (2.07%) was given by T₀ in 2016-2017. During 2017-2018, treatments were exhibited approximately equal performance with slight increase of effective tiller/plant. However, 3.64% and 2.10% were the greater and less value of effective tiller/plant obtained by treatment T₈ and T₀. In the last year 2018-2019, the maximum value on this parameter was found from T₈ (3.89%) and T₄ (3.69%) and then gone downward as followed T₅ (3.36%) >T₇ (3.22) >T₃ (3.09) >T₂ (2.91%) >T₆ (2.84%) >T₁ (2.73%) over T₀ (2.16%).




Fig. 2: Effect of organic and probiotic soil amendments on total plant/m² of wheat.




Fig. 3: Effect of organic and probiotic soil amendments on effective tiller/plant of wheat.

Discussion

Seedling infection

A significant variation of seedling infection was noticed all over the research period. The trend of infection was found to decline with the progress of the study due to application of probiotic and organic manure. The highest rate of seedling infection was found in T₀ (2.32%), while the minimum rate of infection was recorded from T₅ (0.32%) in 2018-2019. It might be due to the definite role of probiotic fortified-organic manures that exerted antagonistic effect against the fungi causing seedling infection. Furthermore, the addition of organic matter may create a hospitable environment for growth and development of fungal antagonists resulting in enhanced biocontrol activity (Sarkar et al. 2002). This finding was in harmony with the reports Tewari and Rajbir (2005).

Seedling blight

Seedling infection later on converted into seedling blight. Like seedling infection, this parameter was also varied significantly under various treatments all over the study period. The control activity against seedling blight was highly observed by T_5 (0.87%) and the lowest exhibited from check T_0 (4.74%) during 2018-2019. Such result may be obtained for the efficacy of the prohibitory and biocontrol action of the treatment. This efficacy of *Trichoderma* fortified-organic manures was evaluated to reduce the pre-emergence and post-emergence seedling mortality, diseases of stem and root of chickpea by Talukder et al. (2017). Similar control activity of *Trichoderma* spp. against plant pathogen and seedling mortality was also reported by Bhuiyan et al. (2007) and Ha 2010.

Leaf chlorophyll content

Chlorophyll content of leaf from diverse treatments was provided significant dissimilarities from beginning to end of the study. On the focus of organic amendments, treatment T_4 (48.37 SPAD) next to T_8 (51.48 SPAD) had the bigger accumulation of chlorophyll and the smaller was noticed from T_0 (40.33 SPAD). It may probably relate with the decomposition rate of organic manures that happened in case of T_4 treatment. This result was supported by Khan et al. (2005) with the statement that N supply in organic treatment is generally restricted and slows N mineralization as compared to crop demand of N. Chlorophyll content is closely related to the presence of N and poultry manure (T_4) had the higher ability of supplying N. This result is an agreement with the findings of Sims and Wolf (1992), Khanam et al. (2001), Ru et al. (2012) and Krishna (2013).

Total plant/m²

The parameter plant/m² is one of the major vegetative growth indicators of wheat. The best values of plant/m² were obtained from T₈ (216.11%) and then T₄ (202.81%) where the lowest at T₀ (158.66%). The ability of such enhancement plant/m² due to application of poultry manure might contain high amounts of nutrients and organic matter that improved the soil structure and environment which aids density of plant. Result in the study was in the same manner previously notified by Hassan (2002), Dauda et al. (2008) and Ismaeil et al. (2012).

Effective tiller/plant

Tillering is one of the important growth stages of wheat which provides the necessary stalks for satisfactory production. Though tillers are lateral branches which emerge from the main stem, but not all produce an ear. Generally, the first two-three tillers are noted as productive tillers of wheat and production is related to environment and nutrition supply. In this investigation, a greater number of productive tiller/plant was demonstrated by T_8 (3.89%) and T_4 (3.69%). This advancement of effective tiller might be due to optimum moisture and available nutrients provided by poultry manure which may have created the favorable condition for wheat growth. This was supported by Simpson (1990), Eck and Stewart (1995), Mitchell and Tu (2005) and Sistani et al. (2008). The result of this investigation was related to the observations of Belefant-Miller (2007), Abbasi and Khaliq (2016).

Conclusion

Quick response of chemical fertilizer offered an instant results nevertheless it has harmful effect not only crops as well as soil health. Therefore, organic soil amendment has given emphasis in this study. Seedling survivability was found greater by the application of probitic combination, whereas potential and sustainable improvement of wheat growth with organic soil amendment was found. Regarding this findings, *Trichoderma harzianum*, green manure and vermicompost may be useful for against seedling mortality and in addition of green manure, vermicompost and poultry manure may be a worthy substitute for wheat growth. Data of this study revealed that the probiotic particularly *Trichoderma harzianum* produce healthier seedling and poultry manure exhibited the capacity of supply nutrient demand and chlorophyll synthesis for improvement in growth of wheat as compared to other manures.

References

- Abbasi MK and Khaliq A (2016). Nitrogen mineralization of a loam soil supplemented with organic-inorganic amendments under laboratory incubation. Frontiers in Plant Science, 7: 1-9.
- Alam MS, Haydar FMA and Hoque A (2020). Effects of planting method on yield and yield attributing characters in chickpea (*Cicer arietinum* L.). Journal of Bio-Science, 28(20): 105-110.
- Ali AM, Ibrahim SM and Bijay-Singh (2020). Wheat grain yield and nitrogen uptake prediction using atLeaf and GreenSeeker portable optical sensors at jointing growth stage. Information Processing in Agriculture, 7(3): 375-383.
- BARC (2012). Fertilizer Recommendation Guide. Farmgate, Dhaka-1215, Bangladesh.
- BARI (2014). Krishi Projukti Hatboi (Handbook on Agro-Technology), 6th Edition, Gazipur-1701, Bangladesh.
- Belefant-Miller H (2007). Poultry litter induces tillering in rice. Journal of Sustainable Agriculture, 31(1): 151-160.
- Bhuiyan MK, Ahmed I and Begum F (2007). Integrated management of seedling mortality of cauliflower (*Brassica oleraceae* var. capitata) caused by *Sclerotium rolfsii*. Journal of Agriculture, Science and Technology, 8(1&2): 79-86.
- Dauda SN, Ajayi FA and Ndor E (2008). Growth and yield of water melon (*Citrullus lanatus*) as affected by poultry manure application. Journal of Agriculture and Social Science, 4(3): 121-124.
- Dubey L, Dubey M and Jain P (2015). Role of green manure in organic farming. Plant archives, 15(1): 23-26.
- Eck HV and Stewart BA (1995). Manure in soil amendments and environmental quality. Lewis publication, Boca Raton, Florida: 169-98.
- Ha TN (2010). Using *Trichoderma* species for biological control of plant pathogens in Viet Nam. International Society for Southeast Asian Agricultural Sciences, 16(1): 17-21.
- Hammad HM, Khaliq A, Ashfaq A, Aslam M, Malik AH, Farhad W and Laghari KQ (2011). Influence of different organic manures on wheat productivity. International Journal of Agriculture and Biology, 13(1): 137-140.
- Haque M and Islam SMS (2018). Effect of sowing time on yield and yield contributing characters in comparison with local and European genotypes of barley (*Hordeum Vulgare* L.). SKUAST Journal of Research, 20(1): 27-31.
- Hassan EA (2002). Effect of chicken manure and season on the performance and HCN Content of two forage sorghum cultivar. PhD Thesis, Faculty of Agriculture, University of Khartoum, Sudan.
- Hermosa R, Viterbo A, Chet I and Monte E (2012). Plant-beneficial effects of *Trichoderma* and of its genes. Microbiology, 158: 17-25.
- Hossain MA, Hasan MM and Islam SMS (2021b) Sustainability of wheat (*Triticum aestivum* L.) yield and improvement of seed quality through probiotic and organic soil amendment. Bangladesh Agronomy Journal 389. Bangladesh Agronomy Journal, 24(1).
- Hossain MA, Hasan MM, Hossain MB and Islam SMS (2021a). Effects of probiotic and organic fertilizers as soil amendments on growth and yield of wheat plants. Applied Biological Research, 23(2): 157-164. DOI: 10.5958/0974-4517.2021.00021.5
- Ismaeil FM, Abusuwar AO and El Naim AM (2012). Influence of chicken manure on growth and yield of forage sorghum (Sorghum bicolor L. Moench). International Journal of Agriculture and Forestry, 2(2): 56-60.
- Karlen DL, Varvel GE, Bullock DG and Cruse RM (1994). Crop rotations for the 21st Century. Advances in Agronomy, 53: 01-45.
- Khan MA, Shukla AK, Upadhyay NC and Singh OP (2005). Nutrient management strategies for maize-potato-wheat cropping system in typic ustochrept soils. Potato Journal, 32: 167-168.

- Khanam M, Rahman MH and Islam MR (2001). Effect of manure and fertilizers on the growth and yield of BRRI Dhan 30. Pakistan Journal of Biological Sciences, 4: 172-174.
- Krishna KR (2013). Agroecosystems: soils, climate, crops nutrients dynamics and productivity. Apple Academic Press Incorporated, Toronto, Canada: 1-517.
- Kumar S and Abraham T (2018). Productivity potential of wheat under certified organic production system. International Journal of Current Microbiology and Applied Sciences, 7(10): 281-288.
- Mitchell CC and Tu S (2005). Long-term evaluation of poultry litter as a source of nitrogen for cotton and corn. Agronomy Journal, 97: 399-407.
- Ru SH, Zhang GY, Niu JF, Sun SY, Wang L and Geng N (2012). Effects of chicken manure application on crop yields and soil nutrients accumulation in wheat-corn rotation system. International Conference on Biomedical Engineering and Biotechnology, 28 May 2012, pp. 1392-1396.
- Sarkar MK, Acharya A and Sengupta PK (2002). Management of sheath blight of rice caused by *Rhizoctonia solani* by oilcake amendment of soil. Journal of Mycopathological Research, 40(2): 153-158.
- Shewry PR and Hey SJ (2015). The contribution of wheat to human diet and health. Food and Energy Security, 4(3): 178-202.
- Simpson TW (1990). Agronomic use of poultry industry waste. Poultry Science, 70: 1126-1131.
- Sims TJ and Wolf DC (1992). Poultry waste management: Agricultural and environmental issues. Advanced Agronomy, 52: 1-84.
- Sistani KR, Sikora FJ and Rasnake M (2008). Poultry litter and tillage influences on corn production and soil nutrients in a Kentucky silt loam soil. Soil and Tillage Research, 98(2): 130-139.
- Talukdar P, Siddiqa MM, Masum MMI, Habibullah ABM and Bhuiyan AKM (2017). Effect of *Trichoderma* fortified compost on disease suppression, growth and yield of chickpea. International Journal of Environment, Agriculture and Biotechnology, 2(2): 831-839.
- Tester CF (1990). Organic amendment effects on physical and chemical properties of a sandy soil. Soil Science Society of America Journal, 54(3): 827-831.
- Tewari L and Rajbir S (2005). Biological control of sheath blight of rice by *Trichoderma harzianum*. Indian Phytopathology, 58: 35-40.
- Yadvinder-Singh, Bijay-Singh, Gupta RK, Ladha JK, Bains JS and Singh J (2008). Evaluation of press mud cake as a source of nitrogen and phosphorus for the rice-wheat cropping system in the Indo-Gangetic plains of India. Biology and Fertility of Soils, 45: 289-296.

(Manuscript received on 21 February 2021; revised on 02 June 2021)

Modification of soil properties through probiotic and organic manures to enhance soil fertility and productivity

Md. Anwar Hossain^{1, 2}, S.M Shahinul Islam¹ and Md. Mahmodol Hasan^{3*} ¹Plant Biotechnology and Genetic Engineering Laboratory, Institute of Biological Sciences, University of Rajshahi, Rajshahi-6205, Bangladesh ²Directorate of Secondary and Higher Education, Dhaka-1000, Bangladesh ³Plant pathology Laboratory, Agronomy and Agricultural Extension Department, University of Rajshahi, Rajshahi-6205, Bangladesh

*Corresponding Authors e-mail:

anwar11nubah55@gmail.com/mahmudul742001@yahoo.com/ shahinul68@gmail.com

Abstract

Presently shifting is accepted from chemical management to organic agriculture for reducing environmental hazard and to ensure soil health for productivity. So, to attain the knowledge on probiotic and organic manures influencing the physical and chemical properties of soil, the investigation was carried out during the period from July 2016 to June 2019. There were nine treatments included in the research for soil amendments. The experiment was followed as Randomized Complete Block Design (RCBD) having three replications. At this point, along with probiotic and organic amendments, chemical fertilizer and control were used for comparative study. The foremost findings were as (1) consistent addition of probiotic and organic manures increased soil physical properties like moisture content, porosity and silt particle and decreased the bulk density, particle density and sand particles. (2) Increased some chemical properties like organic matter content and nutrients availability but decreased pH and C: N ratio. Moreover, most of the physical properties greatly modified ($p \le 0.05$) by compost + vermicompost + green manure (T₃) while maximum parameters of chemical properties changed ($p \le 0.05$) with poultry manure + vermicompost + green manure (T₄). Hence based on soil modification requirements, farmers can be used above mentioned treatment combinations.

Key words: Physical property, chemical property, soil amendment, probiotic and organic manure.

CURRICULUM VITAE

01. Personal detail:

Present address:				
Assistant Professor (Agriculture)				
Officer on Special Duty (OSD)				
Directorate of Secondary and Higher Education				
Dhaka, Bangladesh				
Permanent address:				
Vill: Sailkona Bazar P.O: Nazirpur				
P.S: Bagatipara Dist: Natore-6410				
Bangladesh				

02. Educational qualification:

Name of Examination	Course Duration	Division /Class	Marks Obtained	Passing Year	Board/University	
Secondary School Certificate	10	First	75.60	1989	Rajshahi	
Higher Secondary School Certificate	02	First	74.10	1991	Rajshahi	
Bachelor of Science in Agriculture	04	First	62.20	1995	Bangladesh Agricultural University, Mymensingh	
MS in Agronomy	1.5	First	70.00	2001	Bangladesh Agricultural University, Mymensingh	
Masters in Education	02	CGPA-2.80	70.00	2011	Bangladesh Open University	

03. List of publications:

- 01. **MA Hossain**, MSU Bhuyian and SMA Hossain (2006). Influence of soil moisture management practice and form of urea fertilizer on the growth and yield of maize. Bangladesh Journal of Crop Science. 17(1): 243-249.
- 02. HM Zakir, M Moshfiqur Rahman, A Rahman, I Ahmed and **MA Hossain** (2012). Heavy metals and major ionic pollution assessment in waters of Midstream of the River Karatoa in Bangladesh. Journal of Environmental Science and Natural Resources, 5(2): 149-160.
- 03. Hossain MA, Hasan MM, Hossain MB and Islam SMS (2021). Effects of probiotic and organic fertilizers as soil amendments on growth and yield of wheat plants. Applied Biological Research, 23(2): 157-164. DOI: 10.5958/0974-4517.2021.00021.5
- 04. **Hossain MA**, Hasan MM and Islam SMS (2021) Sustainability of wheat (*Triticum aestivum* L.) yield and improvement of seed quality through probiotic and organic soil amendment. Accepted, BAJ-389. Bangladesh Agronomy Journal, 24(1): 1-11.
- 05. Hossain MA, Hasan MM and Islam SMS (2021). Effects of probiotic and organic manures for soil amendments on seedling survivability and leaf chlorophyll for crop improvement in wheat (*Triticum aestivum* L.). Accepted, Journal of Bio-Science. 29(2): 61-70.
- 06. **Hossain MA**, Hasan MM and Islam SMS. Modification of soil properties through probiotic and organic manures to enhance soil fertility and productivity (In preparation).