

University of Rajshahi

Rajshahi-6205

Bangladesh.

RUCL Institutional Repository

<http://rulrepository.ru.ac.bd>

Department of Mathematics

MPhil Thesis

2015

Separation Axioms on L-Topological Spaces


Islam, Rafiqul

University of Rajshahi

<http://rulrepository.ru.ac.bd/handle/123456789/373>

Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

SEPARATION AXIOMS ON L-TOPOLOGICAL SPACES




**A THESIS
SUBMITTED TO THE UNIVERSITY OF RAJSHAHI FOR
THE DEGREE OF MASTER OF PHILOSOPHY
IN MATHEMATICS**

**BY
RAFIQUL ISLAM**


**In the
Department of Mathematics
Faculty of Science
University of Rajshahi
Rajshahi-6205, Rajshahi
Bangladesh**

December, 2015



Dr. Md. Sahadat Hossain

Associate Professor
Department of Mathematics



University of Rajshahi
Rajshahi – 6205, Bangladesh.
Phone: (Dept.) 0721/711108
(Cell): (01914) 254710
Fax: 0088-0721-750064
E-mail: sahadat@ru.ac.bd

No

Dated

Declaration

Certified that the thesis entitled “SEPARATION AXIOMS ON L -TOPOLOGICAL SPACES” submitted by Rafiqul Islam in fulfillment of the requirements for the degree of Master of Philosophy in Mathematics, Faculty of Science, University of Rajshahi, Rajshahi, Bangladesh has been completed under my supervision. I believe that this research work is an original one and that it has not been submitted elsewhere for any degree.

I wish him a bright future and every success in life.

Supervisor

(Dr. Md. Sahadat Hossain)

Associate Professor
Department of Mathematics
University of Rajshahi
Rajshahi, Bangladesh

STATEMENT OF ORIGINALITY

I declare that the content in my M.Phil. thesis entitled “SEPARATION AXIOMS ON L-TOPOLOGICAL SPACES” is original and accurate to the best of my knowledge. I also declare that the materials contained in my research work have not been previously published or written by any person for any degree or diploma.

(Rafiqul Islam)

Author

ACKNOWLEDGEMENTS

I wish to express first my sincere admiration, appreciation and profound gratitude to my most respected supervisor as well as my most honorable teacher Dr. MD. Sahadat Hossain, Associate Professor, Department of Mathematics, University of Rajshahi for his sincere guidance, valuable suggestions and logical criticisms throughout my research work and also for his rigorous proof-reading of the manuscript during the preparation of my thesis. It is a great pleasure to acknowledge most humbly my indebtedness to him.

I would like to express my gratitude to my honorable teachers Professor Dewan Muslim Ali, Professor Subrata Majumdar, Professor Abdulla Ansary, Professor Abdul Latif, Professor M. Asaduzzaman and Dr. M. Altab Hossain for their valuable advice, inspiration and encouragement during my research work. I am also thankful to my other friends for their inspiration and indirect co-operation during my research work.

I am deeply thankful to my wife Tahsin Islam for her constant inspiration and encouragement. My sense of fathomless gratitude to my parents and other members of the family for their unmeasured sacrifices, continuous inspiration and support, that have led me to all the success.

Finally, I would like to express special thanks to the authority of the University of Rajshahi and in particular to Professor Wazed Ali Pramanik, Chairman, Department of Mathematics for extending all facilities and co-operation during the research work of my M. Phil. program.

University of Rajshahi
Rajshahi, Bangladesh

(Rafiqul Islam)
Author

Contents

	Page No
Introduction	1-4
 Chapter 1: Preliminaries	 5-24
1.1 Introduction	5-6
1.2 Fuzzy set	6-9
1.3 Fuzzy topology	9-10
1.4 L-Fuzzy set	11-16
1.5 Mapping in L-fuzzy set	16-17
1.6 L-Topology	18-21
1.7 Continuous map, Open map and closed map	21-22
1.8 “Good extension” and Product in L-topology	22-24
 Chapter 2: On T_0 Space in L-Topological Spaces	 25-34
2.1 Introduction	25
2.2 T_0 -property in L-Topological Spaces	25-28

2.3	“Good extension”, Hereditary, Productive and Projective Properties in L-Topology	28-32
2.4	Mapping in L-topological spaces	32-34
Chapter 3:	On T_1 Space in L-Topological Spaces	35-44
3.1	Introduction	35
3.2	T_1 -property in L-Topological Spaces	35-38
3.3	“Good extension”, Hereditary, Productive and Projective Properties in L-Topology	38-42
3.4	Mapping in L-topological spaces	42-44
Chapter 4:	On T_2 Space in L-Topological Spaces	45-55
4.1	Introduction	45
4.2	T_2 -property in L-Topological Spaces	45-48
4.3	“Good extension”, Hereditary, Productive and Projective Properties in L-Topology	48-52
4.4	Mapping in L-topological spaces	52-55
Chapter 5:	On R_0 Space in L-Topological Spaces	56-64

5.1	Introduction	56
5.2	R_0 -property in L-Topological Spaces	56-57
5.3	“Good extension”, Hereditary, Productive and Projective Properties in L-Topology	57-61
5.4	Mapping in L-topological spaces	61-64
Chapter 6:	On R_1 Space in L-Topological Spaces	65-76
6.1	Introduction	65
6.2	R_1 -property in L-Topological Spaces	65-68
6.3	“Good extension”, Hereditary, Productive and Projective Properties in L-Topology	69-73
6.4	Mapping in L-topological spaces	73-76
	References	77-88

Introduction

The notion of a fuzzy set, as proposed by L.A. Zadeh[90] in 1965 to provide a foundation for the evolution of many areas of knowledge. After then in quick succession, L-fuzzy sets were introduced by Goguen[24] in 1967. As a result, this provides a natural frame work for generalizing many algebraic and topological concepts in various directions such as L-fuzzy sets, fuzzy logics, fuzzy control, fuzzy groups, fuzzy rings, fuzzy vector spaces, fuzzy topology, fuzzy bitopology, L-topology etc. Many other branches of mathematics have been developed all over the world during the last five decades. Chang [12] first introduced and studied the concept of a fuzzy topological space by using the fuzzy set in 1968. Hutton[31-34], Reilly[33-34], Wong [83-84], Lowen[47-48], Srivastava[73-79], Dude[16-17], Cutler[14], Ying[88], Ali[2-9], Hossain[26-30], Pu and Liu[53-54], etc., discussed and developed various aspects of fuzzy topological spaces. Ying [88] introduced fuzzifying topology and developed this in a new direction with the semantic methods of continuous valued logic. With the help of fuzzifying topology, Sinha[69-70] introduced and studied T_0 , T_1 , T_2 -(Hausdorff), T_3 -(regular), T_4 -(normal), separation axioms. Mashhour et al. [51-52] introduced and studied the concepts of the family of fuzzifying semi-open sets, fuzzifying neighbourhood structure of a point and fuzzifying semi-closure of a fuzzy set. Also in fuzzifying

topology they introduced and studied semi- T_0 -, semi- R_0 -, semi- T_1 -, semi- R_1 -, semi- T_2 (semi Hausdorff)-, semi- T_3 (semi regular)-, semi- T_4 (semi normal)-, separation axioms. Ali and Hossain [28] developed the R_0 and R_1 separation axioms and studied their relations with the T_1 and T_2 -separation axioms respectively. In 1993, Warner and Mclean [80] introduced on compact Hausdorff L-fuzzy spaces. Later Jin-xuan, Ren Bai-lin[35] introduced and studies a set of new separation axioms in L-fuzzy topological spaces. After wards, Kudri[41-42], Li[43-44], Song[71], Xu[86], ZHAO Bin[93], introduced the L-fuzzy topological spaces and studied the strong Hausdorff Separation property in L-fuzzy topological spaces.

In this present thesis, we are going to introduce some new definitions of separation axioms in L-topological spaces using the ideas of Jin-xuan and Ren Bai-lin[35]. Some of their equivalent formulations along with various new characterizations and results concerning the existing ones are presented here. Our criterion for definitions has been preserving as much as possible the relation between the corresponding separation properties for L-topological spaces. Moreover, it will be seen that the definitions of these axioms are ‘good extensions’ in the sense of Lowen [47-48].

We aim to develop theories of L- T_0 , L- T_1 , L- T_2 (Hausdorff), L- R_0 and L- R_1 -separation axioms analogous to its counterpart in ordinary topology. The

materials of this thesis have been divided into six chapters, a brief scenario of which we present as follows.

The first one is to incorporate some of the basic definitions and results of fuzzy set, fuzzy topology, fuzzy mapping, L-topology and its mapping. These results are ready references for the work in the subsequent chapter. Results are stated without proof and can be seen in the papers referred to.

Our work starts from chapter two. In this chapter, we have introduced and studied T_0 properties in L-topological spaces. Here we add eight more definitions to this list and we have established relationship among them. All these eight definitions are ‘good extensions’ of the corresponding concept T_0 in a topological space. We prove that all these definitions satisfy property of hereditary, productive and projective. Also we have studied some other properties of these concepts.

In chapter three, we have developed and studied T_1 properties in L-topological spaces. Here, we include eight more definitions to this chapter and we have established relations among them. All these eight notions are ‘good extensions’ of the corresponding concept of T_1 in a topological space. We prove that all these notions are hereditary, productive and projective. We have discussed some other properties of these concepts.

We have introduced and studied T_2 (Hausdorff) properties in L-topological spaces, in chapter four. We have developed here seven more definitions and we

established relations among them. All these properties are ‘good extensions’ of the corresponding concept T_2 (Hausdorff) in a topological space. We have observed that all the definitions satisfy property of hereditary, productive and projective.

Chapter five is based on the R_0 properties in L-topological spaces. Here we have obtained seven more definitions. We see that all these properties are ‘good extensions’ of the corresponding notions in topological spaces. We have discussed that all the properties are hereditary, productive and projective. We have also studied several other properties of these concepts.

The R_1 properties in L-topological spaces are to be studied in chapter six. We have given here seven more definitions and we have established relations among them. All these are ‘good extensions’ of their corresponding concept in a topological space. Some other pleasant properties of these concepts have been studied here.

CHAPTER-1

Preliminaries

1.1 Introduction:

We have discussed the fuzzy sets, fuzzy topological spaces and L-topological spaces in this chapter. We have incorporated some of the basic definitions and results of the fuzzy sets, Grade of membership, L-fuzzy sets, complement of L-fuzzy sets, some laws of L-fuzzy sets, different mapping on L-fuzzy sets, Fuzzy topological spaces, L-topological spaces, L-T₀, L-T₁, L-T₂-spaces, fuzzy product topological spaces and L-product topological spaces which are to be used as ready references for understanding the subsequent chapters. Most of the results are quoted from various research papers. We make use the following general notations in this thesis paper.

Λ : Index set.

L : Complete distributive lattice with 0 and 1.

$I = [0, 1]$: Closed unit interval.

$I_1 = [0, 1)$: Right open unit interval.

$I_0 = (0, 1]$: Left open unit interval.

$\mu, \alpha, \beta, \gamma, \dots$: L-Fuzzy sets.

(X, t) : Fuzzy topological space.

(X, τ) : L-topological space.

(X, T) : General topological space.

$\prod_{i \in \Lambda} X_i$: Usual product of X_i .

$(X, \tau_1 \times \tau_2)$: Product of L-topologies τ_1 and τ_2 on the set X .

$I_\alpha(\tau) = \{u^{-1}(\alpha, 1] : u \in \tau^*\}$, $\alpha \in I_1$: General topology on X .

1.2 Fuzzy Set

1.2.1 Definition [90]: Let X be a non-empty set and $I = [0, 1]$. A fuzzy set in X is a function $u: X \rightarrow I$ which assign to each element $x \in X$, a degree of membership, $u(x) \in I$. Thus a usual subset of X , is a special type of a fuzzy set in which the range of the function is $[0, 1]$.

1.2.2 Definition: Let X be a nonempty set and A be a subset of X . The function $1_A : X \rightarrow \{0, 1\}$ defined by

$$1_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{if } x \notin A \end{cases} \quad \text{is called the characteristic function of } A. \text{ We also}$$

write 1_x for the characteristic function of X .

1.2.3 Definition: The characteristic functions of subsets of a set X are referred to as the crisp sets in X .

1.2.4 Example: Suppose X is real number R and the fuzzy set of real numbers much greater than 5 in X that could be defined by the continuous function

$$U : X \rightarrow [0,1] \text{ such that } u(x) = \begin{cases} 0 & \text{if } x \leq 5 \\ \frac{x-5}{50} & \text{if } 5 < x < 55 \\ 1 & \text{if } x \geq 55 \end{cases}.$$

1.2.5 Definition [47]: A fuzzy subset is empty if and only if grade of membership is identically zero in X . It is denoted by 0.

1.2.6 Definition [47]: A fuzzy subset is whole if and only if its grade of membership is identically one in X . It is denoted by 1.

1.2.7 Definition [82]: Let u and v be two fuzzy subsets of a set X . Then u is said to be subset of v , i.e., $u \subseteq v$ if and only if $u(x) \leq v(x)$ for every $x \in X$.

1.2.8 Definition [82]: Let u and v be two fuzzy subsets of a set X . Then u is said to be equal to v , i.e., $u = v$ if and only if $u(x) = v(x)$ for every $x \in X$.

1.2.9 Definition [82]: Let u and v be two fuzzy subsets of a set X . Then u is said to be the complement of v , i.e., $u = v^c$ if and only if $u(x) = 1 - v(x)$, for every $x \in X$. Obviously $(v^c)^c = v$.

1.2.10 Definition [11]: Let u and v be two fuzzy subsets of a set X . Then the union w of u and v , i.e., $w = u \cup v$ if and only if $w(x) = (u \cup v)(x) =$

$\max \{ u(x), v(x) \}$, for every $x \in X$. The union w is a fuzzy subset of X .

In general, if Λ be an index set and $A = \{u_i : i \in \Lambda\}$ be a family of fuzzy sets of X then the union $\cup u_i$ is defined by

$$(\cup u_i)(x) = \sup \{u_i(x) : i \in \Lambda\}, x \in X.$$

1.2.11 Definition [11]: Let u and v be two fuzzy subsets of a set X . Then the intersection m of u and v , i.e., $m = u \cap v$ is a fuzzy subset of X if and only if $m(x) = (u \cap v)(x) = \min \{u(x), v(x)\}, \forall x \in X$, and $(\cap u_i)(x) = \inf \{u_i(x) : i \in \Lambda\}, x \in X$, where $\{u_i, i \in \Lambda\}$.

1.2.12 Definition: Let u and v be two fuzzy subsets of a set X . Then the difference of u and v is defined by $u - v = u \cap v^c$.

1.2.13 Definition: If $\alpha \in I$ and $u \in I^X$ define by $u(x) = \alpha$, for all $x \in X$, we refer to u as a constant fuzzy set and denote it by α itself. In particular, we have the constant fuzzy sets 0 and 1.

1.2.14 Example: Let $X = \{x, y, z\}$ and $u, v \in I^X$ are defined by $u(x) = .6$, $u(y) = .7$, $u(z) = .5$ and $v(x) = .7$, $v(y) = .5$, $v(z) = .4$. Then $(u \cup v)(x) = \max \{u(x), v(x)\} = .7$, $(u \cup v)(y) = \max \{u(y), v(y)\} = .7$, $(u \cup v)(z) = \max \{u(z), v(z)\} = .5$, $(u \cap v)(x) = \min \{u(x), v(x)\} = .6$, $(u \cap v)(y) = \min \{u(y), v(y)\} = .5$, $(u \cap v)(z) = \min \{u(z), v(z)\} = .4$, $u^c(x) = 1 - u(x) = .4$, $u^c(y) = 1 - u(y) = .3$, $u^c(z) = 1 - u(z) = .5$.

1.2.15 Laws of the algebra of fuzzy sets:

As in ordinary set theory, idempotent laws, associative law, commutative law, distributive laws, identity law, demorgan's laws hold in the case of fuzzy sets also. But the complement laws are not necessarily true. For example, if

$X = \{a, b, c\}$ and u is a fuzzy subset of X where is defined by

$$u = \{ (a, .2), (b, .7), (c, 1) \},$$

$$\text{then } u^c = \{ (a, .8), (b, .3), (c, 0) \}$$

$$\text{so } u \cup u^c = \{ (a, .8), (b, .7), (c, 1) \} \neq 1,$$

$$u \cap u^c = \{ (a, .2), (b, .3), (c, 0) \} \neq 0.$$

Also in ordinary set theory $U \cap V = \phi$ if and only if $U \subset V^c$. But in fuzzy subsets reverse is not necessary true. For example if

$$v = \{ (a, .6), (b, .2), (c, 0) \} \text{ then } u \subset v^c,$$

$$u \cap v = \{ (a, .2), (b, .2), (c, 0) \} \neq 0.$$

1.3 Fuzzy Topology

1.3.1 Definition [12]: Let $I = [0,1]$, X be a non-empty set and I^X be the collection of all mappings from X into I , i. e. the class of all fuzzy sets in X . A fuzzy topology on X is defined as a family t of members of I^X , satisfying the following conditions:

$$(i) \quad 1, 0 \in t$$

- (ii) if $u_i \in t$ for each $i \in \Lambda$ then $\bigcup_{i \in \Lambda} u_i \in t$
- (iii) if u_1, u_2 then $u_1 \cap u_2 \in t$.

The pair (X, t) is called a fuzzy topological space (fts, in short) and the members of t are called t -open (or simply open) fuzzy sets. A fuzzy set v is called a t -closed (or simply closed) fuzzy set if $1 - v \in t$.

1.3.2 Example : Let $X = \{ a, b, c, d \}$, $t = \{ 0, 1, u, v \}$,

where $1 = \{ (a, 1), (b, 1), (c, 1), (d, 1) \}$

$$0 = \{ (a, 0), (b, 0), (c, 0), (d, 0) \}$$

$$u = \{ (a, .2), (b, .5), (c, .7), (d, .9) \}$$

$$v = \{ (a, .3), (b, .5), (c, .8), (d, .95) \}$$

Then (X, t) is a fuzzy topological space.

1.3.3 Definition: A fuzzy topological space (X, t) is said to be fuzzy regular if and only if for each $x \in X$ and closed fuzzy set u with $u(x) = 0$, there exists open fuzzy sets $v, w \in t$ such that $v(x) = 1$, $u \subseteq w$ and $v \subseteq 1 - w$.

1.3.4 Definition: A fuzzy topological space (X, t) is said to be fuzzy normal if and only if for each close fuzzy set m and open fuzzy set u with $m \subseteq u$, there exists a fuzzy set v such that $m \subseteq v^\circ \subseteq \bar{v} \subseteq u$.

1.4 L-Fuzzy Set

1.4.1 Definition [24]: Let X be a non-empty set and L be a complete distributive lattice with 0 and 1. An L -fuzzy set in X is a function $\alpha: X \rightarrow L$ which assign to each element $x \in X$, a degree of membership, $\alpha(x) \in L$.

1.4.2 Definition [24]: Let α be an L -fuzzy set in X . Then $1 - \alpha = \alpha'$ is called the complement of α in X .

1.4.3 Definition [24]: An L -fuzzy subset is empty if and only if grade of membership is identically zero in X . It is denoted by 0^* .

1.4.4 Definition [24]: An L -fuzzy subset is whole if and only if its grade of membership is identically one in X . It is denoted by 1^* .

1.4.5 Definition [24]: Let α and β be two L -fuzzy subsets of a set X . Then α is said to be subset of β , i.e., $\alpha \subseteq \beta$ if and only if $\alpha(x) \leq \beta(x)$ for every $x \in X$.

1.4.6 Definition [24]: Let α and β be two L -fuzzy subsets of a set X . Then α is said to be equal to β , i.e., $\alpha = \beta$ if and only if $\alpha(x) = \beta(x)$ for every $x \in X$.

1.4.7 Definition [24]: Let α and β be two L -fuzzy subsets of a set X . Then α is said to be the complement of β , i.e., $\alpha = \beta^c$ if and only if $\alpha(x) = 1 - \beta(x)$, for every $x \in X$. Obviously $(\beta^c)^c = \beta$.

1.4.8 Definition [24]: Let α and β be two L-fuzzy subsets of a set X . Then the union γ of α and β , i.e., $\gamma = \alpha \cup \beta$ if and only if $\gamma(x) = (\alpha \cup \beta)(x) = \max \{ \alpha(x), \beta(x) \}$, for every $x \in X$. The union γ is an L-fuzzy subset of X .

In general, if Λ be an index set and $A = \{ \alpha_i : i \in \Lambda \}$ be a family of L-fuzzy sets of X then the union $\cup \alpha_i$ is defined by $(\cup \alpha_i)(x) = \sup \{ \alpha_i(x) : i \in \Lambda \}, x \in X$.

1.4.9 Definition [24]: Let α and β be two L-fuzzy subsets of a set X . Then the intersection m of α and β , i.e., $m = \alpha \cap \beta$ is an L-fuzzy subset of X if and only if $m(x) = (\alpha \cap \beta)(x) = \min \{ \alpha(x), \beta(x) \}, \forall x \in X$, and $(\cap \alpha_i)(x) = \inf \{ \alpha_i(x) : i \in \Lambda \}, x \in X$, where $\{ \alpha_i, i \in \Lambda \}$.

1.4.10 Definition [24]: Let α and β be two L-fuzzy subsets of a set X . Then the difference of α and β is defined by $\alpha - \beta = \alpha \cap \beta^c$.

1.4.11 Example: Let $X = \{x, y, z\}$, $L = \{0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1\}$ and $\alpha, \beta \in L^X$ are defined by $\alpha(x) = .6, \alpha(y) = .7, \alpha(z) = .5$ and $\beta(x) = .7, \beta(y) = .5, \beta(z) = .4$. Then $(\alpha \cup \beta)(x) = \max \{ \alpha(x), \beta(x) \} = .7$, $(\alpha \cup \beta)(y) = \max \{ \alpha(y), \beta(y) \} = .7$, $(\alpha \cup \beta)(z) = \max \{ \alpha(z), \beta(z) \} = .5$, $(\alpha \cap \beta)(x) = \min \{ \alpha(x), \beta(x) \} = .6$, $(\alpha \cap \beta)(y) = \min \{ \alpha(y), \beta(y) \} = .5$, $(\alpha \cap \beta)(z) = \min \{ \alpha(z), \beta(z) \} = .4$, $\alpha^c(x) = 1 - \alpha(x) = .4$, $\alpha^c(y) = 1 - \alpha(y) = .3$, $\alpha^c(z) = 1 - \alpha(z) = .5$.

1.4.12 Definition [45]: Two L-fuzzy sets α and β in X are said to be intersected if and only if there exist a point $x \in X$ such that $(\alpha \cap \beta)(x) \neq 0$. In this case we say that α and β intersect at x .

1.4.13 Definition [46]: Let X be a non empty set and μ be an L-fuzzy set in X . A α -cut of μ is defined by $\alpha_\mu = \{x: \mu(x) \geq \alpha, \forall \alpha \in L\}$.

1.4.14 Definition [46]: Let X be a non empty set and μ be an L-fuzzy set in X . A strong α -cut of μ is defined by $\alpha_\mu^+ = \{x: \mu(x) > \alpha, \forall \alpha \in L\}$. We see that α -cut and strong α -cut are crisp subsets of X . The 1-cut of μ is called the core of μ .

1.4.15 Definition [46]: Let X be a non empty set and μ be an L-fuzzy set in X . The support of μ in X is the crisp subset of X that contains all the elements of X that have none zero membership grads in μ , i.e., $\text{supp}\mu = \{x: \mu(x) > 0\}$.

1.4.16 Definition [46]: The height $h(\mu)$ of an L-fuzzy set μ is the largest membership grade obtained by any element in that set, i.e., $h(\mu) = \sup_{x \in X} \mu(x)$.

1.4.17 Definition [46]: For a finite L-fuzzy set, the cardinality $|\alpha|$ defined as $|\alpha| = \sum_{x \in X} \alpha(x)$. $\|\alpha\| = \frac{|\alpha|}{|x|}$ is called the relative cardinality of α .

1.4.18 Definition [46]: An L-fuzzy set μ is called normal when $h(\mu) = 1$; it is called subnormal when $h(\mu) < 1$. The height of μ may also be viewed as the supremum of α for which $\alpha_\mu \neq \emptyset$.

1.4.19 Definition [90]: If $r \in L$ and α is an L-fuzzy set in X defined by $\alpha(x) = r, \forall x \in X$ then we refer to α as a constant L-fuzzy set and denoted it by r itself.

In particular, we have the constant L-fuzzy sets 0 and 1.

1.4.20 Definition [46]: An L-fuzzy point p in X is a special L-fuzzy set with membership function $p(x) = r$ if $x = x_0$

$$= 0 \text{ otherwise } x \neq x_0 \text{ where } r \in L .$$

1.4.21 Definition [46]: An L-fuzzy point p is said to belong to an L-fuzzy set α in X ($p \in \alpha$) if and only if $p(x) < \alpha(x)$ and $p(y) \leq \alpha(y)$

$$(y \neq x) \text{ i.e. } x_r \in \alpha \Rightarrow r < \alpha(x) .$$

1.4.22 Definition [90]: An L-fuzzy singleton in X is an L-fuzzy set in X which is zero everywhere except at one point say x , where it takes a value say r with $0 < r \leq 1$ and $r \in L$. We denote it by x_r and $x_r \in \alpha$ iff $r \leq \alpha(x)$.

1.4.23 Definition [46]: An L-fuzzy singleton x_r is said to be quasi-coincident (q-coincident, in short) with an L-fuzzy set α in X , denoted by $x_r q \alpha$ iff $r + \alpha(x) > 1$. Similarly, an L-fuzzy set α in X is said to be q-coincident with an L-fuzzy set β in X , denoted by $\alpha q \beta$ if and only if $\alpha(x) + \beta(x) > 1$ for some $x \in X$. Therefore $\alpha \bar{q} \beta$ iff $\alpha(x) + \beta(x) \leq 1$ for all $x \in X$, where $\alpha \bar{q} \beta$ denotes that an L-fuzzy set α in X is not q-coincident with an L-fuzzy set β in X .

1.4.24 Laws of the algebra of L-fuzzy sets:

As in ordinary set theory, idempotent laws, associative laws, commutative laws, distributive laws, identity laws, Demorgan's laws hold in the case of L-fuzzy sets also. But the complement laws are not necessarily true. For example, if $X = \{a, b, c\}$, $L = \{0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1\}$ and α is an L-fuzzy subset of X where is defined by

$$\alpha = \{ (a, .2), (b, .7), (c, 1) \},$$

$$\text{then } \alpha^c = \{ (a, .8), (b, .3), (c, 0) \}$$

$$\text{so } \alpha \cup \alpha^c = \{ (a, .8), (b, .7), (c, 1) \} \neq 1,$$

$$\alpha \cap \alpha^c = \{ (a, .2), (b, .3), (c, 0) \} \neq 0.$$

Also in ordinary set theory $U \cap V = \emptyset$ if and only if $U \subset V^c$. But in L-fuzzy subsets reverse is not necessary true. For example

if $\beta = \{ (a, .6), (b, .2), (c, 0) \}$ then $\alpha \subset \beta^c$,

$\alpha \cap \beta = \{ (a, .2), (b, .2), (c, 0) \} \neq 0$.

1.5 Mapping in L-fuzzy set

1.5.1 Definition [23]: Let f be a mapping from a set X into a set Y and α is an L-fuzzy subset on X . Then f and α induced an L-fuzzy subset $\beta = f(\alpha)$ of Y whose membership function is defined by

$$\beta(y) = f(\alpha)(y) = \begin{cases} \sup\{\alpha(x)\} & \text{if } f^{-1}[\{y\}] \neq \emptyset, x \in X \\ 0, & \text{otherwise} \end{cases}$$

1.5.2 Definition [11]: Let f be a mapping from a set X into Y and β is an L-fuzzy subset of Y . Then the inverse of β written as $\alpha = f^{-1}(\beta)$ is an L-fuzzy subset of X and is defined by $\alpha(x) = (f^{-1}(\beta))(x) = \beta(f(x))$, for $x \in X$.

1.5.3Example: Suppose that $X = \{x, y, z, w\}$ and $Y = \{a, b, c\}$. Define $f: X \longrightarrow Y$ by $f(x) = b, f(y) = c, f(z) = a, f(w) = a$. Let $\alpha \in L^X$ be given by $\alpha(x) = .2, \alpha(y) = .3, \alpha(z) = .5$ and $\alpha(w) = .4$. Then $(f(\alpha))(a) = \sup\{\alpha(z), \alpha(w)\} = .5$. Similarly, $(f(\alpha))(b) = \sup\alpha(x) = .2$, and $(f(\alpha))(c) = \sup\alpha(y) = .3$.

On the other hand, if β is an L-fuzzy set in Y given by $\beta(a) = .6, \beta(b) = .8, \beta(c) = .7$. Then $(f^{-1}(\beta))(x) = \beta(f(x)) = \beta(b) = .8, (f^{-1}(\beta))(y) =$

$$\beta(f(y)) = \beta(c) = .7, (f^{-1}(\beta))(z) = \beta(f(z)) = \beta(a) = .6, (f^{-1}(\beta))(w) = \beta(f(w)) = \beta(a) = .6.$$

We now mention some properties of L-fuzzy subsets induced by mappings.

1.5.4 Definition [11]: Let f be a mapping from X into Y , α be an L-fuzzy subset of X and β be an L-fuzzy subset of Y . Then the following properties are true.

- (a) $f^{-1}(\beta^c) = (f^{-1}(\beta))^c$ for any L-fuzzy subset β of Y .
- (b) $f(\alpha^c) = (f(\alpha))^c$ for any L-fuzzy subset α of X .
- (c) $\beta_1 \subset \beta_2 \Rightarrow f^{-1}(\beta_1) \subset f^{-1}(\beta_2)$, where β_1 and β_2 are two L-fuzzy subsets of Y .
- (d) $\alpha_1 \subset \alpha_2 \Rightarrow f(\alpha_1) \subset f(\alpha_2)$, where α_1 and α_2 are two L-fuzzy subsets of X .
- (e) $\beta \supset f(f^{-1}(\beta))$, for any L-fuzzy subset β of Y .
- (f) $\alpha \subset f^{-1}(f(\alpha))$, for any L-fuzzy subset α of X .
- (g) Let f be a function from X into Y and g be a function from Y into Z .

Then $(g \circ f)^{-1}(\gamma) = f^{-1}(g^{-1}(\gamma))$, for any L-fuzzy subset γ in Z ,

where $(g \circ f)$ is the composition of g and f .

1.6 L-topology

1.6.1 Definition [46]: Let X be a non-empty set and L be a complete distributive lattice with 0 and 1. Again suppose that τ be the sub collection of all mappings from X to L i.e. $\tau \subseteq L^X$. Then τ is called L-topology on X if it satisfies the following conditions:

- (i) $0^*, 1^* \in \tau$
- (ii) If $u_1, u_2 \in \tau$ then $u_1 \cap u_2 \in \tau$
- (iii) If $u_i \in \tau$ for each $i \in \Lambda$ then $\bigcup_{i \in \Lambda} u_i \in \tau$.

Then the pair (X, τ) is called the L-topological space (lts, in short) and the members of τ are called open L-fuzzy sets. An L-fuzzy set v is called a closed L-fuzzy set if $1 - v \in \tau$.

1.6.2 Example: Let $X = \{a, b, c\}, \tau = \{0^*, u, v, 1^*\}$ and

$L = \{0, 0.05, 0.1, 0.15, \dots \dots \dots 0.95, 1\}$. Where $0^* = \{(a, 0), (b, 0), (c, 0)\}, 1^* =$

$\{(a, 1), (b, 1), (c, 1)\}, u = \{(a, 0.1), (b, 0.3), (c, 0.5)\}$ and

$v = \{(a, 0.2), (b, 0.4), (c, 0.6)\}$. Then τ is an L topology on X and the pair (X, τ) is called L-topological space.

1.6.3 Definition [46]: Let λ be an L-fuzzy set in lts (X, τ) . Then the closure of λ is denoted by $\bar{\lambda}$ or $cl\lambda$ and defined as $\bar{\lambda} = \bigcap \{\mu: \lambda \subseteq \mu, \mu \in \tau^c\}$.

The interior of λ written λ^0 or $\text{int}\lambda$ is defined by $\lambda^0 = \cup \{\mu : \mu \subseteq \lambda, \mu \in \tau\}$.

1.6.4 Definition [46]: If (X, τ) is an lts and $A \subseteq X$ then $\tau_A = \{u|A : u \in \tau\}$

is called the sub space L-topology on A and (A, τ_A) is referred to as an

L-sub space of (X, τ) .

1.6.5 Definition [46]: Let (X, τ) be an L-topological space. A subfamily B of τ is a base for τ if and only if each member of τ can be express as the union of some members of B .

1.6.6 Definition [46]: Let (X, τ) be an L-topological space. A subfamily S of τ is a sub-base for τ if and only if the family of finite intersection of members of S forms a base for τ .

1.6.7 Definition [46]: Let x_r be an L-fuzzy point in an lts (X, τ) . An

L-fuzzy set α in X is called a neighborhood (in short, nhd) of x_r if and only

if there exists an open L-fuzzy set β in X such that $x_r \in \beta \subseteq \alpha$.

1.6.8 Definition [46]: An L-fuzzy set u in an L-topological space (X, τ) is called a neighborhood of an L-fuzzy point x_r if and only if there exist an

L-fuzzy set $u_1 \in \tau$ such that $x_r \in u_1 \subseteq u$. A neighborhood u is called an open neighborhood if u is open. The family consisting of all the neighborhoods of x_r is called the system of x_r .

1.6.9 Definition [46]: An L-topological space (X, τ) is said to be $L-T_0$ if and only if

- a) for all distinct elements $x, y \in X$, there exists $u \in \tau$ such that $u(x) = 1, u(y) = 0$ or $u(x) = 0, u(y) = 1$.
- b) for all distinct elements $x, y \in X$, there exists $u \in \tau$ such that $u(x) < u(y)$ or $u(y) < u(x)$.
- c) for all distinct elements $x, y \in X$, $\bar{x}_1 \cap \bar{y}_1 < 1$.

1.6.10 Definition [46]: An L-topological space (X, τ) is said to be T_1 if and only if

- (a) for all distinct elements $x, y \in X$, there exist $u, v \in \tau$ such that $u(x) = 1, u(y) = 0$ and $v(x) = 0, v(y) = 1$;
- (b) for all distinct elements $x, y \in X$, there exist $u, v \in \tau$ such that $u(x) > 0, u(y) = 0$ and $v(x) = 0, v(y) > 0$;
- (c) for all distinct elements $x, y \in X$, there exist $u, v \in \tau$ such that $u(x) > u(y)$ and $v(y) > v(x)$.

1.6.11 Definition [46]: An L-topological space (X, τ) is said to be L-fuzzy Hausdorff or $L-T_2$ if and only if

- (a) for all distinct elements $x, y \in X$, there exist $u, v \in \tau$ such that $u(x) = 1 = u(y)$ and $u \cap v = 0$;

- (b) for all pair of distinct L-fuzzy points $x_r, y_s \in S(X)$, there exist $u, v \in \tau$ such that $x_r \in u, y_s \in v$ and $u \cap v = 0$;
- (c) for all distinct elements $x, y \in X$, there exist $u, v \in \tau$ such that $u(x) > 0$, $v(y) > 0$ and $u \cap v = 0$.

1.7 Continuous map Open map and closed map

1.7.1 Definition [46]: Let f be a real-valued function on an L-topological space. If $\{x: f(x) > \alpha\}$ is open for every real α , then f is called lower-semi continuous function (lsc, in short).

1.7.2 Definition [46]: Let (X, τ) and (Y, s) be two L-topological spaces and f be a mapping from (X, τ) into (Y, s) i . e. $f: (X, \tau) \rightarrow (Y, s)$. Then f is called-

- (i) L-Continuous iff for each open L-fuzzy set $u \in s \Rightarrow f^{-1}(u) \in \tau$
- (ii) L-Open iff $f(\mu) \in s$ for each open L-fuzzy set $\mu \in \tau$.
- (iii) L-Closed iff $f(\lambda)$ is s-closed for each $\lambda \in \tau^c$ ie. λ is a closed L-fuzzy set in X .
- (iv) L-Homeomorphism iff f is bijective and both f and f^{-1} is L-continuous.

1.7.3 Proposition[46]: Let $f : (X, \tau) \longrightarrow (Y, s)$ be an L-fuzzy continuous function, then the following properties hold :

- (i) For every s – closed v , $f^{-1}(v)$ is τ – closed.
- (ii) For each L-fuzzy point p in X and each neighborhood u of $f(p)$, then there exist a neighborhood v of p such that $f(v) \subseteq u$.
- (iii) For any L-fuzzy set u in X , $f(u) \subseteq (f(u))^0$.
- (iv) For any L-fuzzy set v in Y , $(f^{-1}(v))^0 \subseteq f^{-1}(v^0)$.

1.7.4 Proposition[46]: Let $f : (X, \tau) \longrightarrow (Y, s)$ be a L-fuzzy open function , then the following properties hold:

- (i) $f(u^0) \subseteq (f(u))^0$, for each L-fuzzy set u in X .
- (ii) $(f^{-1}(v))^0 \subseteq f^{-1}(v^0)$, for each L-fuzzy set v in Y .

1.7.5 Proposition[46]: Let $f : (X, t) \longrightarrow (Y, s)$ be a function. Then f is closed if and only if $f(u) \subseteq f(u)$ for each fuzzy set u in X .

1.8 “Good extension” and Product in L-topology

1.8.1 Definition [46]: Let X be a none empty set and T be a topology on X .

Let $\tau = \omega(T)$ be the set of all lower semi continuous (lsc) functions from

(X, T) to L (with usual topology). Thus $\omega(T) = \{u \in L^X : u^{-1}(\alpha, 1] \in T\}$ for each $\alpha \in L$. It can be shown that $\omega(T)$ is an L-topology on X .

Let “P” be the property of a topological space (X, T) and LFP be its L-topological analogue. Then LFP is called a “good extension” of P “if the statement (X, T) has P iff $(X, \omega(T))$ has LFP” holds good for every topological space (X, T) .

1.8.2 Definition [91]: Let $\{(X_i, \tau_i) : i \in \Lambda\}$ be a family of L-topological space. Then the space $(\prod X_i, \prod \tau_i)$ is called the product Lts of the family $\{(X_i, \tau_i) : i \in \Lambda\}$ where $\prod \tau_i$ denote the usual product L-topologies of the families $\{\tau_i : i \in \Lambda\}$ of L-topologies on X .

1.8.3 Definition [91] : If u_1 and u_2 are two L-fuzzy subsets of X and Y respectively then the Cartesian product $u_1 \times u_2$ of two L-fuzzy subsets u_1 and u_2 is a fuzzy subsets of $X \times Y$ defined by $(u_1 \times u_2)(x, y) = \min\{u_1(x), u_2(y)\}$, for each pair $(x, y) \in X \times Y$.

1.8.4 Definition [46] : Let $\{X_i, i \in \Lambda\}$, be any class of sets and let X denoted the Cartesian product of these sets, i.e., $X = \prod_{i \in \Lambda} X_i$. Note that X consists of all points $p = \langle a_i, i \in \Lambda \rangle$, where $a_i \in X_i$. Recall that, for each $j_0 \in \Lambda$, we define the projection π_{j_0} from the product set X to

the coordinate space X_{j_0} . ie $\pi_{j_0} : X \longrightarrow X_{j_0}$ by $\pi_{j_0} (\langle a_i : i \in \Lambda \rangle) = a_{j_0}$

.These projections are used to define the product L-topology.

1.8.5 Definition [46]: If (X_1 , τ_1) and (X_2 , τ_2) be two L- topological spaces and $X = X_1 \times X_2$ be the usual product and τ be the coarsest L-topology on X , then each projection $\pi_i : X \longrightarrow X_i$, $i = 1 , 2$., is L-fuzzy continuous. The pair (X, τ) is called the product space of the L-topological spaces (X_1, τ_1) and (X_2, τ_2) .

Chapter-2

On T_0 Space in L-Topological Spaces

2.1 Introduction

Fuzzy T_0 spaces have been defined and studied by Hutton and Reilly[33,34], Pu and Liu [53,54]. After then, in quick succession, a large number of seemingly different definitions of fuzzy T_0 spaces were developed and studied by several workers, e.g. Ali [2], Hossain [30], Srivastava [77,78] and Choubey[13] etc. In this chapter we define possible eight definitions of T_0 space in L-topological spaces. We established all these definitions satisfied “good extension” property; also we show that these spaces possess many nice properties and that they are hereditary, productive and projective.

2.2 T_0 -property in L-Topological Spaces

We now give the following definitions of T_0 -property in L-topological spaces.

2.2.1 Definition: An lts (X, τ) is called-

- (a) $L - T_0(i)$ if $\forall x, y \in X, x \neq y$ then $\exists u \in \tau$ such that $u(x) \neq u(y)$.
- (b) $L - T_0(ii)$ if $\forall x, y \in X, x \neq y$ then $\exists u \in \tau$ such that $u(x) = 1, u(y) = 0$ or $u(x) = 0, u(y) = 1$.


- (c) $L - T_0(iii)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u \in \tau$ such that $x_r \in u, y_s \notin u$ or $x_r \notin u, y_s \in u$.
- (d) $L - T_0(iv)$ if for all pairs of distinct L-fuzzy singletons $x_r, y_s \in S(X)$ with $x_r \bar{q} y_s$ then $\exists u \in \tau$ such that $x_r \subseteq u, y_s \bar{q} u$ or $y_s \subseteq u, x_r \bar{q} u$.
- (e) $L - T_0(v)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u \in \tau$ such that $x_r \in u, u \bar{q} y_s$ or $y_s \in u, u \bar{q} x_r$.
- (f) $L - T_0(vi)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u \in \tau$ such that $x_r \in u, y_s \cap u = 0$ or $y_s \in u, x_r \cap u = 0$.
- (g) $L - T_0(vii)$ if $\forall x, y \in X, x \neq y$ then $\exists u \in \tau$ such that $u(x) > 0, u(y) = 0$ or $u(x) = 0, u(y) > 0$.
- (h) $L - T_0(viii)$ if $\forall x, y \in X, x \neq y$ then $\exists u \in \tau$ such that $u(x) > u(y)$ or $u(y) > u(x)$.

A complete comparison of the definitions $L - T_0(ii), L - T_0(iii),$

$L - T_0(iv), L - T_0(v), L - T_0(vi), L - T_0(vii)$ and $L - T_0(viii)$ with

$L - T_0(i)$ are given below:

2.2.2Theorem: Let (X, τ) be an lts. Then we have the following implications:



The reverse implications are not true in general.

Proof: $L - T_0(ii) \Rightarrow L - T_0(i)$, $L - T_0(iii) \Rightarrow L - T_0(i)$ and

$L - T_0(iv) \Rightarrow L - T_0(i)$ can be proved . Now $L - T_0(v) \Rightarrow L - T_0(i)$ since $L - T_0(v) \Leftrightarrow L - T_0(iii)$.

$L - T_0(vi) \Rightarrow L - T_0(i)$, since $L - T_0(vi) \Rightarrow L - T_0(v)$. $L - T_0(vii)$ and $L - T_0(viii) \Rightarrow L - T_0(i)$ since $L - T_0(viii) \Rightarrow L - T_0(vii)$ and

$L - T_0(vii) \Rightarrow L - T_0(ii)$.

None of the reverse implications are true, it can be seen through the following counter example: Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u\}$ where $u(x) = 0.5$, $u(y) = 0.3$ and $L = \{0, 0.05, 0.1, 0.15, \dots, 0.95, 1\}$.

Proof: $L - T_0(i) \not\Rightarrow L - T_0(ii)$: Here the lts (X, τ) is clearly $L - T_0(i)$ but it is not $L - T_0(ii)$. Since there is no non empty L-fuzzy set in τ which takes zero value at x or y .

$L - T_0(i) \not\Rightarrow L - T_0(iii)$: For if we take the distinct L-fuzzy points $x_{3/4}, y_{4/5}$, there does not exist $u \in \tau$ such that $x_{3/4} \in u, y_{4/5} \notin u$ or $x_{3/4} \notin u, y_{4/5} \in u$.

$L - T_0(i) \not\Rightarrow L - T_0(iv)$: As for the distinct L-fuzzy singletons x_1, y_1 in τ there does not exist $u \in \tau$ such that $x_1 \subseteq u, y_1 \bar{q}u$ or $y_1 \subseteq u, x_1 \bar{q}u$.

$L - T_0(i) \not\Rightarrow L - T_0(v)$: This follows automatically from the fact that

$L - T_0(v) \Leftrightarrow L - T_0(iii)$ and it has already been shown that $L - T_0(i) \not\Rightarrow L - T_0(iii)$.

$L - T_0(i) \not\Rightarrow L - T_0(vi)$: Since for any two distinct L-fuzzy points $x_{3/4}, y_{4/5}$ in X , there does not exist $u \in \tau$ which is disjoint with $x_{3/4}$ or $y_{4/5}$.

$L - T_0(i) \not\Rightarrow L - T_0(vii)$ and $L - T_0(i) \not\Rightarrow L - T_0(viii)$: It is obvious because $L - T_0(vii) \Rightarrow L - T_0(ii)$ and $L - T_0(viii) \Rightarrow L - T_0(ii)$ and it has already been shown that $L - T_0(i) \not\Rightarrow L - T_0(ii)$:

2.3 “Good extension”, Hereditary, Productive and Projective Properties in L-Topology

Now all the definitions $L - T_0(i), L - T_0(ii), L - T_0(iii), L - T_0(iv),$

$L - T_0(v), L - T_0(vi), L - T_0(vii), L - T_0(viii)$ are ‘good extensions’ of T_0 - property, is shown below:

2.3.1Theorem: Let (X, T) be a topological space. Then (X, T) is T_0 iff $(X, \omega(T))$ is $L - T_0(i)$.

Proof: Let (X, T) be T_0 space. Choose $x, y \in X$ with $x \neq y$. Then $\exists U \in T$ such that $x \in U, y \notin U$ or $y \in U, x \notin U$. Now consider the characteristics function 1_U . Then $1_U \in \omega(T)$ such that $1_U(x) = 1, 1_U(y) = 0$ and so that $1_U(x) \neq 1_U(y)$. Thus $(X, \omega(T))$ is $L - T_0(i)$.

Conversely, let $(X, \omega(T))$ be $L - T_0(i)$. To show that (X, T) is T_0 . Choose $x, y \in X$ with $x \neq y$. Then $\exists u \in \omega(T)$ such that $u(x) \neq u(y)$. Let $u(x) < u(y)$. Choose r such that $u(x) < r < u(y)$ and consider $u^{-1}(r, 1]$. Then $u^{-1}(r, 1] \in T$ with $x \notin u^{-1}(r, 1]$ and $y \in u^{-1}(r, 1]$. Hence (X, T) is T_0 . Similarly we can easily show that each of $L - T_0(ii), L - T_0(iii), L - T_0(iv), L - T_0(v), L - T_0(vi), L - T_0(vii), L - T_0(viii)$ are also holds 'good extension' property.

2.3.2Theorem: Let (X, τ) be an lts, $A \subseteq X$ and $\tau_A = \{u|_A : u \in \tau\}$, then

- (a) (X, τ) is $L - T_0(i) \Rightarrow (A, \tau_A)$ is $L - T_0(i)$.
- (b) (X, τ) is $L - T_0(ii) \Rightarrow (A, \tau_A)$ is $L - T_0(ii)$.
- (c) (X, τ) is $L - T_0(iii) \Rightarrow (A, \tau_A)$ is $L - T_0(iii)$.
- (d) (X, τ) is $L - T_0(iv) \Rightarrow (A, \tau_A)$ is $L - T_0(iv)$.
- (e) (X, τ) is $L - T_0(v) \Rightarrow (A, \tau_A)$ is $L - T_0(v)$.
- (f) (X, τ) is $L - T_0(vi) \Rightarrow (A, \tau_A)$ is $L - T_0(vi)$.
- (g) (X, τ) is $L - T_0(vii) \Rightarrow (A, \tau_A)$ is $L - T_0(vii)$.

(h) (X, τ) is $L - T_0(viii) \Rightarrow (A, \tau_A)$ is $L - T_0(viii)$.

Proof: We prove only (b). Suppose (X, τ) is L -topological space and $L - T_0(ii)$.

We shall prove that (A, τ_A) is $L - T_0(ii)$. Let $x, y \in A$ with $x \neq y$, then $x, y \in X$ with $x \neq y$ as $A \subseteq X$. Since (X, τ) is $L - T_0(ii)$, $\exists u \in \tau$ such that $u(x) = 1, u(y) = 0$ or $u(x) = 0, u(y) = 1$. For $A \subseteq X$ we find $u|_A \in \tau_A$ such that $u|_A(x) = 1, u|_A(y) = 0$ or $u|_A(x) = 0, u|_A(y) = 1$. Hence it is clear that the subspace (A, τ_A) is $L - T_0(ii)$.

Similarly, (a), (c), (d), (e), (f), (g), (h) can be easily proved.

2.3.3 Theorem: Given $\{(X_i, \tau_i): i \in \Lambda\}$ be a family of L -topological space. Then the product of L -topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_0(j)$ iff each coordinate space (X_i, τ_i) is $L - T_0(j)$ where $j = i, ii, iii, iv, v, vi, vii, viii$.

Proof: Let each coordinate space $\{(X_i, \tau_i): i \in \Lambda\}$ be $L - T_0(ii)$. We show that the product space is $L - T_0(ii)$. Suppose $x, y \in X$ with $x \neq y$, again suppose that $x = \Pi x_i, y = \Pi y_i$ then $x_j \neq y_j$ for some $j \in \Lambda$. Now consider $x_j, y_j \in X_j$. Since (X_j, τ_j) is $L - T_0(ii)$, $\exists u_j \in \tau_j$ such that $u_j(x_j) = 1, u_j(y_j) = 0$ or $u_j(x_j) = 0, u_j(y_j) = 1$. Suppose $u_j(x_j) = 1, u_j(y_j) = 0$. Now take $u = \Pi u'_j$ where $u'_j = u_j$ and $u_i = 1$ for $i \neq j$. Then u is such that $u(x) = 1, u(y) = 0$. Hence the product L -topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_0(ii)$.

Conversely, let the product L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_0(ii)$. Take any coordinate space (X_j, τ_j) , choose $x_j, y_j \in X_j, x_j \neq y_j$. Now construct $x, y \in X$ such that $x = \Pi x'_i, y = \Pi y'_i$ where $x'_i = y'_i$ for $i \neq j$ and $x'_j = x_j, y'_j = y_j$. Then $x \neq y$ and hence $\exists u \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0$ or $u(x) = 0, u(y) = 1$. Suppose $u(x) = 1, u(y) = 0$. Now u must be the union of basic open L-fuzzy set say $u = \bigcup_{k \in K} b_k$. Thus $\bigcup b_k(x) = 1$ and $\bigcup b_k(y) = 0$ which implies that there exist at least one k such that $b_k(x) = 1, b_k(y) = 0$. Now let $b_k = \Pi v_i$ where $v_i = 1$ except for finitely many i 's. So $\Pi v_i(x) = 1, \Pi v_i(y) = 0, i.e. \inf v_i(x'_i) = 1$ and $\inf v_i(y'_i) = 0$, which implies that $v_j(x_j) = 1, v_j(y_j) = 0$. Since $x'_i = y'_i$ for $i \neq j$, thus (X_i, τ_i) is $L - T_0(ii)$.

Moreover one can easily verify that

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(i) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(i)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(iii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(iii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(iv) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(iv)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(v) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(v)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(vi) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(vi)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(vii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(vii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_0(viii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_0(viii).$$

Hence, we see that $L - T_0(i), L - T_0(ii), L - T_0(iii), L - T_0(iv),$

$L - T_0(v), L - T_0(vi), L - T_0(vii)$ and $L - T_0(viii)$ Property is productive and projective.

2.4 Mapping in L-topological spaces

We show that $L - T_0(j)$ property is preserved under one-one, onto and continuous mapping for $j = i, ii, iii, iv, v, vi, vii, viii$.

2.4.1Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, onto and L-open map, then-

- (a) (X, τ) is $L - T_0(i) \Rightarrow (Y, s)$ is $L - T_0(i)$.
- (b) (X, τ) is $L - T_0(ii) \Rightarrow (Y, s)$ is $L - T_0(ii)$.
- (c) (X, τ) is $L - T_0(iii) \Rightarrow (Y, s)$ is $L - T_0(iii)$.
- (d) (X, τ) is $L - T_0(iv) \Rightarrow (Y, s)$ is $L - T_0(iv)$.
- (e) (X, τ) is $L - T_0(v) \Rightarrow (Y, s)$ is $L - T_0(v)$.
- (f) (X, τ) is $L - T_0(vi) \Rightarrow (Y, s)$ is $L - T_0(vi)$.
- (g) (X, τ) is $L - T_0(vii) \Rightarrow (Y, s)$ is $L - T_0(vii)$.
- (h) (X, τ) is $L - T_0(viii) \Rightarrow (Y, s)$ is $L - T_0(viii)$.

Proof: Suppose (X, τ) is $L - T_0(ii)$. We shall prove that (Y, s) is

$L - T_0(ii)$. Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto, $\exists x_1, x_2 \in X$ such that $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one. Again

since (X, τ) is $L - T_0(ii)$, $\exists u \in \tau$ such that $u(x_1) = 1, u(x_2) = 0$ or $u(x_1) = 0, u(x_2) = 1$.

Now

$$f(u)(y_1) = \{supu(x_1): f(x_1) = y_1\} = 1$$

$$f(u)(y_2) = \{supu(x_2): f(x_2) = y_2\} = 0$$

or

$$f(u)(y_1) = \{supu(x_1): f(x_1) = y_1\} = 0$$

$$f(u)(y_2) = \{supu(x_2): f(x_2) = y_2\} = 1.$$

Since f is L -open, $f(u) \in s$. Now it is clear that $\exists f(u) \in s$ such that $f(u)(y_1) = 1, f(u)(y_2) = 0$ or $f(u)(y_1) = 0, f(u)(y_2) = 1$. Hence it is clear that the L -topological space (Y, s) is $L - T_0(ii)$. Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

2.4.2 Theorem: Let (X, τ) and (Y, s) be two L -topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be L -continuous and one-one map, then-

- (a) (Y, s) is $L - T_0(i) \Rightarrow (X, \tau)$ is $L - T_0(i)$.
- (b) (Y, s) is $L - T_0(ii) \Rightarrow (X, \tau)$ is $L - T_0(ii)$.
- (c) (Y, s) is $L - T_0(iii) \Rightarrow (X, \tau)$ is $L - T_0(iii)$.
- (d) (Y, s) is $L - T_0(iv) \Rightarrow (X, \tau)$ is $L - T_0(iv)$.
- (e) (Y, s) is $L - T_0(v) \Rightarrow (X, \tau)$ is $L - T_0(v)$.
- (f) (Y, s) is $L - T_0(vi) \Rightarrow (X, \tau)$ is $L - T_0(vi)$.

(g) (Y, s) is $L - T_0(vii) \Rightarrow (X, \tau)$ is $L - T_0(vii)$.

(h) (Y, s) is $L - T_0(viii) \Rightarrow (X, \tau)$ is $L - T_0(viii)$.

Proof: Suppose (Y, s) is $L - T_0(ii)$. We shall prove that (X, τ) is $L - T_0(ii)$.

Let $x_1, x_2 \in X$ with $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$ as f is one-one. Since (Y, s) is $L - T_0(ii)$, then $\exists u \in s$ such that $u(f(x_1)) = 1, u(f(x_2)) = 0$ or $u(f(x_1)) = 0, u(f(x_2)) = 1$. Suppose $u(f(x_1)) = 1, u(f(x_2)) = 0$. This implies that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0$ and $f^{-1}(u) \in \tau$ as f is

L -continuous and $u \in s$. Now it is clear that $f^{-1}(u) \in \tau$ such that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0$. Hence the L -topological space (X, τ) is $L - T_0(ii)$.

Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

Chapter-3

On T_1 Space in L-Topological Spaces

3.1 Introduction

The concept of fuzzy T_1 space was introduced Ali [2, 3], Hossain[26], Srivastava and Lal[79], Sinha[69,70], Malghan[49,50] and other mathematician have contributed to the development of the theory. In this chapter, we discuss possible eight definitions of T_1 space in L-topological spaces; all these notions satisfy “good extension” property. We show that these notions possess many nice properties which are hereditary, productive and projective.

3.2 T_1 -property in L-Topological Spaces

Here, we define the following definitions of T_1 -property in L-topological spaces.

3.2.1 Definition: An lts (X, τ) is called-

- (a) $L - T_1(i)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) \neq u(y)$ and $v(x) \neq v(y)$.


- (b) $L - T_1(ii)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) = 1, u(y) = 0$ and $v(x) = 0, v(y) = 1$.
- (c) $L - T_1(iii)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u, y_s \notin u$ and $x_r \notin v, y_s \in v$.
- (d) $L - T_1(iv)$ if for all pairs of distinct L-fuzzy singletons $x_r, y_s \in S(X)$ with $x_r \bar{q} y_s$ then $\exists u, v \in \tau$ such that $x_r \subseteq u, y_s \bar{q} u$ and $y_s \subseteq v, x_r \bar{q} v$.
- (e) $L - T_1(v)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u, u \bar{q} y_s$ and $y_s \in v, v \bar{q} x_r$.
- (f) $L - T_1(vi)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u, y_s \cap u = 0$ and $y_s \in v, x_r \cap v = 0$.
- (g) $L - T_1(vii)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) > 0, u(y) = 0$ and $v(x) = 0, v(y) > 0$.
- (h) $L - T_1(viii)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) > u(y)$ and $v(y) > v(x)$.

Here, we established a comparison of the definitions $L - T_1(ii)$,

$L - T_1(iii), L - T_1(iv), L - T_1(v), L - T_1(vi), L - T_1(vii), L - T_1(viii)$ with

$L - T_1(i)$ is given below:

3.2.2Theorem: Let (X, τ) be an lts. Then we have the following implications:



The reverse implications are not true in general.

Proof: $L - T_1(ii) \Rightarrow L - T_1(i)$, $L - T_1(iii) \Rightarrow L - T_1(i)$ and $L - T_1(iv) \Rightarrow$

$L - T_1(i)$ can be proved easily. Now $L - T_1(v) \Rightarrow L - T_1(i)$ since

$$L - T_1(v) \Leftrightarrow L - T_1(iii).$$

$$L - T_1(vi) \Rightarrow L - T_1(i), \quad \text{since} \quad L - T_1(vi) \Rightarrow L - T_1(v). \quad L - T_1(vii) \text{ and}$$

$$L - T_1(viii) \Rightarrow L - T_1(i) \text{ since } L - T_1(viii) \Rightarrow L - T_1(vii) \text{ and } L - T_1(vii) \Rightarrow$$

$$L - T_1(ii).$$

None of the reverse implications are true; it can be seen through the following example:

Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u, v\}$

where $u(x) = 0.5$, $u(y) = 0.6$ and $v(x) = 0.7$, $v(y) = 0.4$ and

$$L = \{0, 0.05, 0.1, 0.15, \dots \dots 0.95, 1\}.$$

Proof: $L - T_1(i) \not\Rightarrow L - T_1(ii)$: Here the lts (X, τ) is clearly $L - T_1(i)$ but it is not $L - T_1(ii)$. Since there is no non empty L-fuzzy set in τ which takes zero value at x or y .

$L - T_1(i) \not\Rightarrow L - T_1(iii)$: For if we take the distinct L-fuzzy points $x_{3/5}, y_{1/2} \in S(X)$, then there does not exist $u, v \in \tau$ such that $x_{3/5} \in u, y_{1/2} \notin u$ and $x_{3/5} \notin v, y_{1/2} \in v$.

$L - T_1(i) \not\Rightarrow L - T_1(iv)$: As for the distinct L-fuzzy singletons x_1, y_1 in τ there does not exist $u, v \in \tau$ such that $x_1 \subseteq u, y_1 \bar{q} u$ and $y_1 \subseteq v, x_1 \bar{q} v$.

$L - T_1(i) \not\Rightarrow L - T_1(v)$: This follows automatically from the fact that

$L - T_1(v) \Leftrightarrow L - T_1(iii)$ and it has already been shown that $L - T_1(i) \not\Rightarrow L - T_1(iii)$.

$L - T_1(i) \not\Rightarrow L - T_1(vi)$: Since for any two distinct L-fuzzy points $x_{3/5}, y_{1/2}$ in $S(X)$, then there does not exist $u, v \in \tau$ which is disjoint with $x_{3/5}$ and $y_{1/2}$.

$L - T_1(i) \not\Rightarrow L - T_1(vii)$ and $L - T_1(i) \not\Rightarrow L - T_1(viii)$: It is obvious because $L - T_1(vii) \Rightarrow L - T_1(ii)$ and $L - T_1(viii) \Rightarrow L - T_1(ii)$ and it has already been shown that $L - T_1(i) \not\Rightarrow L - T_1(ii)$.

3.3 “Good extension”, Hereditary, Productive and Projective Properties in L-Topology

Here, we show that all the definitions $L - T_1(i), L - T_1(ii), L - T_1(iii),$

$L - T_1(iv), L - T_1(v), L - T_1(vi), L - T_1(vii)$ and $L - T_1(viii)$ are ‘good extensions’ of T_1 - property, is shown below:

3.3.1Theorem: Let (X, T) be a topological space. Then (X, T) is T_1 iff $(X, \omega(T))$ is $L - T_1(i)$.

Proof: Let (X, T) be T_1 . Choose $x, y \in X$ with $x \neq y$. Then $\exists U, V \in T$ such that $x \in U, y \notin U$ and $y \in V, x \notin V$. Now consider the lower semi continuous functions $1_U, 1_V$. Then $1_U, 1_V \in \omega(T)$ with $1_U(x) = 1, 1_U(y) = 0$ and $1_V(x) = 0, 1_V(y) = 1$ and so that $1_U(x) \neq 1_U(y)$ and $1_V(x) \neq 1_V(y)$. Thus $(X, \omega(T))$ is $L - T_1(i)$.

Conversely, let $(X, \omega(T))$ be $L - T_1(i)$. To show that (X, T) is T_1 . Choose $x, y \in X$ with $x \neq y$. Then $\exists u, v \in \omega(T)$ such that $u(x) \neq u(y)$ and $v(x) \neq v(y)$. Let $u(x) < u(y)$ and $v(y) < v(x)$. Choose r and s such that $u(x) < r < u(y)$ and $v(y) < s < v(x)$ and consider $u^{-1}(r, 1]$ and $v^{-1}(s, 1]$. Then $u^{-1}(r, 1], v^{-1}(s, 1] \in T$ and is $x \notin u^{-1}(r, 1], y \in u^{-1}(r, 1]$ and $x \in v^{-1}(s, 1], y \notin v^{-1}(s, 1]$. Hence (X, T) is T_1 .

Similarly we can show that $L - T_1(ii), L - T_1(iii), L - T_1(iv), L - T_1(v),$

$L - T_1(vi), L - T_1(vii), L - T_1(viii)$ are also hold ‘good extension’ property.

3.3.2Theorem: Let (X, τ) be an lts, $A \subseteq X$ and $\tau_A = \{u|_A : u \in \tau\}$, then

- (a) (X, τ) is $L - T_1(i) \Rightarrow (A, \tau_A)$ is $L - T_1(i)$.
- (b) (X, τ) is $L - T_1(ii) \Rightarrow (A, \tau_A)$ is $L - T_1(ii)$.
- (c) (X, τ) is $L - T_1(iii) \Rightarrow (A, \tau_A)$ is $L - T_1(iii)$.
- (d) (X, τ) is $L - T_1(iv) \Rightarrow (A, \tau_A)$ is $L - T_1(iv)$.

(e) (X, τ) is $L - T_1(v) \Rightarrow (A, \tau_A)$ is $L - T_1(v)$.

(f) (X, τ) is $L - T_1(vi) \Rightarrow (A, \tau_A)$ is $L - T_1(vi)$.

(g) (X, τ) is $L - T_1(vii) \Rightarrow (A, \tau_A)$ is $L - T_1(vii)$.

(h) (X, τ) is $L - T_1(viii) \Rightarrow (A, \tau_A)$ is $L - T_1(viii)$.

Proof: We prove only (b). Suppose (X, τ) is L -topological space and

$L - T_1(ii)$. We shall prove (A, τ_A) is $L - T_1(ii)$. Let $x, y \in A$ with $x \neq y$, then $x, y \in X$ with $x \neq y$ as $A \subseteq X$. Since (X, τ) is $L - T_1(ii)$, $\exists u, v \in \tau$ such that $u(x) = 1, u(y) = 0$ and $v(x) = 0, v(y) = 1$. For $A \subseteq X$ we find $u|_A, v|_A \in \tau_A$ and $u|_A(x) = 1, u|_A(y) = 0$ and $v|_A(x) = 0, v|_A(y) = 1$ as $x, y \in A$. Hence it is clear that the subspace (A, τ_A) is $L - T_1(ii)$.

Similarly, (a), (c), (d), (e), (f), (g), (h) can be easily proved.

3.3.3 Theorem: Given $\{(X_i, \tau_i): i \in \Lambda\}$ be a family of L -topological space. Then the product of L -topological space $(\prod X_i, \prod \tau_i)$ is $L - T_1(j)$ iff each coordinate space (X_i, τ_i) is $L - T_1(j)$ where $j = i, ii, iii, iv, v, vi, vii, viii$.

Proof: Let each coordinate space $\{(X_i, \tau_i): i \in \Lambda\}$ be $L - T_1(ii)$. Then we show that the product space is $L - T_1(ii)$. Suppose $x, y \in X$ with $x \neq y$, again suppose $x = \prod x_i, y = \prod y_i$ then $x_j \neq y_j$ for some $j \in \Lambda$. Now consider

$x_j, y_j \in X_j$. Since (X_j, τ_j) is $L - T_1(ii)$, $\exists u_j, v_j \in \tau_j$ such that $u_j(x_j) = 1, u_j(y_j) = 0$ and $v_j(x_j) = 0, v_j(y_j) = 1$. Now take $u = \prod u'_j, v = \prod v'_j$ where $u'_j = u_j, v'_j = v_j$ and $u_i = v_i = 1$ for $i \neq j$. Then $u, v \in \prod \tau_i$ such that $u(x) =$

$1, u(y) = 0$ and $v(x) = 0, v(y) = 1$. Hence the product L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_1(ii)$.

Conversely, let the product L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_1(ii)$.

Take any coordinate space (X_j, τ_j) , choose $x_j, y_j \in X_j, x_j \neq y_j$. Now construct $x, y \in X$ such that $x = \Pi x'_i, y = \Pi y'_i$ where $x'_i = y'_i$ for $i \neq j$ and $x'_j = x_j, y'_j = y_j$. Then $x \neq y$ and using the product space $L - T_1(ii)$ $\exists u, v \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0$ and $v(x) = 0, v(y) = 1$. Now choose any L-fuzzy point x_r in u . Then \exists a basic open L-fuzzy set $\Pi u_j^r \in \Pi \tau_j$ such that $x_r \in \Pi u_j^r \subseteq u$ which implies that $r < \Pi u_j^r(x)$ or that $r < \inf_j u_j^r(x'_j)$ and hence $r < \Pi u_j^r(x'_j) \forall j \in \Lambda \dots \dots (i)$ and $u(y) = 0 \Rightarrow \Pi u_j(y) = 0 \dots \dots (ii)$. Similarly, corresponding to a fuzzy point $y_s \in v$ there exists a basic open L-fuzzy set $\Pi v_j^s \in \Pi \tau_j$ that $y_s \in \Pi v_j^s \subseteq v$ which implies that $s < v_j^s(j) \forall j \in \Lambda \dots \dots (iii)$ and $v_j^s(y) = 0 \dots \dots (iv)$. Further, $\Pi u_j^r(y) = 0 \Rightarrow u_i^r(y_i) = 0$, since for $j \neq i, x'_j = y'_j$ and hence from (i), $u_j^r(y_j) = u_j^r(x_j) > r$. Similarly, $\Pi v_j^s(x) = 0 \Rightarrow v_i^s(x_i) = 0$ using (iii). Thus we have $u_i^r(x_i) > r, u_i^r(y_i) = 0$ and $v_i^s(y_i) > s, v_i^s(x_i) = 0$. Now consider $\sup_r u_i^r = u_i, \sup_s v_i^s = v_i \in \tau_i$ then $u_i(x_i) = 1, u_i(y_i) = 0$ and $v_i(x_i) = 0, v_i(y_i) = 1$ showing that (X_i, τ_i) is $L - T_1(ii)$.

Moreover one can easily verify that

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(i) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(i)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(iii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(iii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(iv) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(iv)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(v) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(v)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(vi) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(vi)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(vii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(vii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_1(viii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_1(viii) .$$

Hence we see that $L - T_1(i), L - T_1(ii), L - T_1(iii), L - T_1(iv),$

$L - T_1(v), L - T_1(vi), L - T_1(vii), L - T_1(viii)$ Properties are productive and projective.

3.4 Mapping in L-topological spaces

We show that $L - T_1(j)$ property is preserved under one-one, onto and continuous mapping for $j = i, ii, iii, iv, v, vi, vii, viii$.

3.4.1 Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, onto and L-open map, then-

$$(a) (X, \tau) \text{ is } L - T_1(i) \Rightarrow (Y, s) \text{ is } L - T_1(i).$$

$$(b) (X, \tau) \text{ is } L - T_1(ii) \Rightarrow (Y, s) \text{ is } L - T_1(ii).$$

$$(c) (X, \tau) \text{ is } L - T_1(iii) \Rightarrow (Y, s) \text{ is } L - T_1(iii).$$

$$(d) (X, \tau) \text{ is } L - T_1(iv) \Rightarrow (Y, s) \text{ is } L - T_1(iv).$$

$$(e) (X, \tau) \text{ is } L - T_1(v) \Rightarrow (Y, s) \text{ is } L - T_1(v).$$

(f) (X, τ) is $L - T_1(vi) \Rightarrow (Y, s)$ is $L - T_1(vi)$.

(g) (X, τ) is $L - T_1(vii) \Rightarrow (Y, s)$ is $L - T_1(vii)$.

(h) (X, τ) is $L - T_1(viii) \Rightarrow (Y, s)$ is $L - T_1(viii)$.

Proof: Suppose (X, τ) is $L - T_1(ii)$. We shall prove that (Y, s) is $L - T_1(ii)$.

Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto, $\exists x_1, x_2 \in X$ such that $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one. Again since (X, τ) is $L - T_1(ii)$ $\exists u, v \in \tau$ such that $u(x_1) = 1, u(x_2) = 0$ and $v(x_1) = 0, v(x_2) = 1$. Now

$$f(u)(y_1) = \{supu(x_1): f(x_1) = y_1\} = 1$$

$$f(u)(y_2) = \{supu(x_2): f(x_2) = y_2\} = 0 \text{ and}$$

$$f(v)(y_1) = \{supv(x_1): f(x_1) = y_1\} = 0$$

$$f(v)(y_2) = \{supv(x_2): f(x_2) = y_2\} = 1.$$

Since f is L-open, $f(u), f(v) \in s$. Now it is clear that $\exists f(u), f(v) \in s$ such that $f(u)(y_1) = 1, f(u)(y_2) = 0$ and $f(v)(y_1) = 0, f(v)(y_2) = 1$. Hence it is clear that the L-topological space (Y, s) is $L - T_1(ii)$.

Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

3.4.2Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be L-continuous and one-one map, then-

(a) (Y, s) is $L - T_1(i) \Rightarrow (X, \tau)$ is $L - T_1(i)$.

(b) (Y, s) is $L - T_1(ii) \Rightarrow (X, \tau)$ is $L - T_1(ii)$.

(c) (Y, s) is $L - T_1(iii) \Rightarrow (X, \tau)$ is $L - T_1(iii)$.

(d) (Y, s) is $L - T_1(iv) \Rightarrow (X, \tau)$ is $L - T_1(iv)$.

(e) (Y, s) is $L - T_1(v) \Rightarrow (X, \tau)$ is $L - T_1(v)$.

(f) (Y, s) is $L - T_1(vi) \Rightarrow (X, \tau)$ is $L - T_1(vi)$.

(g) (Y, s) is $L - T_1(vii) \Rightarrow (X, \tau)$ is $L - T_1(vii)$.

(h) (Y, s) is $L - T_1(viii) \Rightarrow (X, \tau)$ is $L - T_1(viii)$.

Proof: Suppose (Y, s) is $L - T_1(ii)$. We shall prove that (X, τ) is $L - T_1(ii)$.

Let $x_1, x_2 \in X$ with $x_1 \neq x_2, \Rightarrow f(x_1) \neq f(x_2)$ as f is one-one. Since (Y, s) is $L - T_1(ii)$, $\exists u, v \in s$ such that $u(f(x_1)) = 1, u(f(x_2)) = 0$ and $v(f(x_1)) = 0, v(f(x_2)) = 1$. This implies that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0$ and $f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$ and hence $f^{-1}(u), f^{-1}(v) \in \tau$ as f is

L -continuous and $u, v \in s$. Now it is clear that $f^{-1}(u), f^{-1}(v) \in \tau$ such that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0$ and $f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$.

Hence the L -topological space (X, τ) is $-T_1(ii)$.

Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

Chapter-4

On T_2 Space in L-Topological Spaces

4.1 Introduction:

Hausdorff, [25] introduced the fundamental concept of T_2 space in general topology. T_2 space in fuzzy topology was introduced by Ghanim et.al. [23], Ganguly [22], Shinha [70] and Fora[1] etc. Later FT_2 space has been developed by Ali [2, 6], Cutler [14], Reilly [34] and Hossain [27]. Seven concepts of T_2 space in L-topological spaces are introduced and studied in this chapter. We showed that all these concepts satisfy “good extension” property. We also establish some relationships among them and study some other properties of these spaces.

4.2 T_2 -property in L-Topological Spaces

We now give the following definitions of T_2 -property in L-topological spaces.

4.2.1 Definition: An lts (X, τ) is called-

(a) $L - T_2(i)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) =$

$1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$.


- (b) $L - T_2(ii)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u, y_s \notin u$ and $x_r \notin v, y_s \in v$ and $u \cap v = 0$.
- (c) $L - T_2(iii)$ if for all pairs of distinct L-fuzzy singletons $x_r, y_s \in S(X)$ with $x_r \bar{q} y_s$ then $\exists u, v \in \tau$ such that $x_r \subseteq u, y_s \bar{q} u$ and $y_s \subseteq v, x_r \bar{q} v$ and $u \cap v = 0$.
- (d) $L - T_2(iv)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u, u \bar{q} y_s$ and $y_s \in v, v \bar{q} x_r$ and $u \cap v = 0$.
- (e) $L - T_2(v)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ then $\exists u, v \in \tau$ such that $x_r \in u \subseteq co y_s, y_s \in v \subseteq co x_r$ and $u \subseteq co v$.
- (f) $L - T_2(vi)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) > 0, u(y) = 0$ and $v(x) = 0, v(y) > 0$.
- (g) $L - T_2(vii)$ if $\forall x, y \in X, x \neq y$ then $\exists u, v \in \tau$ such that $u(x) > u(y)$ and $v(y) > v(x)$.

Here, we established a complete comparison of the definitions

$L - T_2(ii), L - T_2(iii), L - T_2(iv), L - T_2(v), L - T_2(vi)$ and $L - T_2(vii)$

with $L - T_2(i)$.

4.2.2Theorem: Let (X, τ) be an lts. Then we have the following implications:



The reverse implications are not true in general, except $L - T_2(vi)$ and $L - T_2(vii)$.

Proof: $L - T_2(i) \Rightarrow L - T_2(ii)$, $L - T_2(i) \Rightarrow L - T_2(iii)$ can be proved easily. Now $L - T_2(i) \Rightarrow L - T_2(iv)$ and $L - T_2(i) \Rightarrow L - T_2(v)$, since $L - T_2(ii) \Leftrightarrow L - T_2(iv)$ and $L - T_2(iv) \Leftrightarrow L - T_2(v)$. $L - T_2(i) \Rightarrow L - T_2(vi)$; It is obvious. $L - T_2(i) \Rightarrow L - T_2(vii)$, since $L - T_2(vi) \Rightarrow L - T_2(vii)$.

The reverse implications are not true in general, except $L - T_2(vi)$ and $L - T_2(vii)$, as can be seen through the following counter-examples:

Example-1: Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u, v\}$ where $u(x) = 0.5, u(y) = 0$ $v(x) = 0, v(y) = 0.6$, $L = \{0, 0.05, 0.1, 0.15, \dots, 0.95, 1\}$ and $r = 0.4, s = 0.3$.

Example-2: Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u, v\}$ where $u(x) = 0.5, u(y) = 0$ $v(x) = 0, v(y) = 0.4$, $L = \{0, 0.05, 0.1, 0.15, \dots, 0.95, 1\}$ and $r = 0.5, s = 0.4$.

Proof: $L - T_2(ii) \not\Rightarrow L - T_2(i)$: From example-1, we see that the lts (X, τ) is clearly $L - T_2(ii)$ but it is not $L - T_2(i)$. Since there is no L-fuzzy set in τ which grade of membership is 1.

$L - T_2(iii) \not\Rightarrow L - T_2(i)$: From example-2, we see the lts (X, τ) is clearly $L - T_2(iii)$ but it is not $L - T_2(i)$. Since $L - T_2(iii) \not\Rightarrow L - T_2(ii)$ and $L - T_2(ii) \not\Rightarrow L - T_2(i)$ so $L - T_2(iii) \not\Rightarrow L - T_2(i)$.

$L - T_1(i) \not\Rightarrow L - T_1(iv)$: As for the distinct L-fuzzy singletons x_1, y_1 in τ there does not exist $u, v \in \tau$ such that $x_1 \subseteq u, y_1 \bar{q} u$ and $y_1 \subseteq v, x_1 \bar{q} v$.

$L - T_2(iv) \not\Rightarrow L - T_2(i)$: This follows from the fact that

$L - T_2(ii) \Leftrightarrow L - T_2(iv)$ and it has already been shown that $L - T_2(ii) \not\Rightarrow L - T_2(i)$ so $L - T_2(iv) \not\Rightarrow L - T_2(i)$.

$L - T_2(v) \not\Rightarrow L - T_2(i)$: Since $L - T_2(iv) \Leftrightarrow L - T_2(v)$ and $L - T_2(iv) \not\Rightarrow L - T_2(i)$ so $L - T_2(v) \not\Rightarrow L - T_2(i)$. But $L - T_2(vii) \Rightarrow L - T_2(vi) \Rightarrow T_2(i)$ is obvious.

4.3 “Good extension”, Hereditary, Productive and Projective Properties in L-Topology

Here we showed that all definitions $L - T_2(i), L - T_2(ii), L - T_2(iii), L - T_2(iv), L - T_2(v), L - T_2(vi)$ and $L - T_2(vii)$ are ‘good extensions’ of T_2 – property, is shown below:

4.3.1Theorem: Let (X, T) be a topological space. Then (X, T) is T_2 iff $(X, \omega(T))$ is $L - T_2(j)$, where $j = i, ii, iii, iv, v, vi, vii$.

Proof: Let (X, T) be T_2 . Choose $x, y \in X$ with $x \neq y$. Then $\exists U, V \in T$ such that $x \in U, y \notin U$ and $y \in V, x \notin V$ and $U \cap V = \emptyset$. Now consider the lower semi continuous functions $1_U, 1_V$. Then $1_U, 1_V \in \omega(T)$ such that $1_U(x) = 1, 1_U(y) = 0$ and $1_V(x) = 0, 1_V(y) = 1$ and so that $1_U \cap 1_V = 0$. Thus $(X, \omega(T))$ is $L - T_2(i)$.

Conversely, let $(X, \omega(T))$ be $L - T_2(i)$. To show that (X, T) is T_2 . Choose $x, y \in X$ with $x \neq y$. Then $\exists u, v \in \omega(T)$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$, i.e., $u(y) < u(x)$ and $v(x) < v(y)$. Choose r and s such that $u(y) < r < u(x)$ and $v(x) < s < v(y)$ and consider $u^{-1}(r, 1]$ and $v^{-1}(s, 1]$. Then $u^{-1}(r, 1], v^{-1}(s, 1] \in T$ and is $x \notin u^{-1}(r, 1], y \in u^{-1}(r, 1], x \in v^{-1}(s, 1], y \notin v^{-1}(s, 1]$ and $u^{-1}(r, 1] \cap v^{-1}(s, 1] = \emptyset$ as $u \cap v = 0$. Hence (X, T) is T_2 .

Similarly, we can show that $L - T_2(ii), L - T_2(iii), L - T_2(iv), L - T_2(v), L - T_2(vi), L - T_2(vii)$ are also hold ‘good extension’ property.

4.3.2Theorem: Let (X, τ) be an lts, $A \subseteq X$ and $\tau_A = \{u|_A : u \in \tau\}$, then

- (a) (X, τ) is $L - T_2(i) \Rightarrow (A, \tau_A)$ is $L - T_2(i)$.
- (b) (X, τ) is $L - T_2(ii) \Rightarrow (A, \tau_A)$ is $L - T_2(ii)$.

(c) (X, τ) is $L - T_2(iii) \Rightarrow (A, \tau_A)$ is $L - T_2(iii)$.

(d) (X, τ) is $L - T_2(iv) \Rightarrow (A, \tau_A)$ is $L - T_2(iv)$.

(e) (X, τ) is $L - T_2(v) \Rightarrow (A, \tau_A)$ is $L - T_2(v)$.

(f) (X, τ) is $L - T_2(vi) \Rightarrow (A, \tau_A)$ is $L - T_2(vi)$.

(g) (X, τ) is $L - T_2(vii) \Rightarrow (A, \tau_A)$ is $L - T_2(vii)$.

Proof: The author proved only (a). Suppose (X, τ) is L -topological space and $L - T_2(i)$. We shall prove (A, τ_A) is $L - T_2(i)$. Let $x, y \in A$ with $x \neq y$, then $x, y \in X$ with $x \neq y$ as $A \subseteq X$. Since (X, τ) is $L - T_2(i)$, $\exists u, v \in \tau$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. For $A \subseteq X$ we find $u|A, v|A \in \tau_A$ and $u|A(x) = 1, u|A(y) = 0$ and $v|A(x) = 0, v|A(y) = 1$ and $u|A \cap v|A = (u \cap v)|A = 0$ as $x, y \in A$. Hence it is clear that the subspace (A, τ_A) is $L - T_2(i)$.

Similarly, (b), (c), (d), (e), (f), (g) can be proved.

4.3.3 Theorem: Given $\{(X_i, \tau_i): i \in \Lambda\}$ be a family of L -topological space. Then the product of L -topological space $(\prod X_i, \prod \tau_i)$ is $L - T_2(j)$ iff each coordinate space (X_i, τ_i) is $L - T_2(j)$ where $j = i, ii, iii, iv, v, vi, vii$.

Proof: Let each coordinate space $\{(X_i, \tau_i): i \in \Lambda\}$ be $L - T_2(i)$. We showed that the product space is $L - T_2(i)$. Suppose $x, y \in X$ with $x \neq y$, again suppose $x = \prod x_i, y = \prod y_i$ then $x_j \neq y_j$ for some $j \in \Lambda$. Now consider $x_j, y_j \in X_j$. Since (X_j, τ_j) is $L - T_2(i)$, $\exists u_j, v_j \in \tau_j$ such that $u_j(x_j) = 1, u_j(y_j) = 0, v_j(x_j) = 0, v_j(y_j) = 1$ and $u_j \cap v_j = 0$.

Now take $u = \Pi u'_j, v = \Pi v'_j$ where $u'_j = u_j, v'_j = v_j$ and $u_i = v_i = 1$ for $i \neq j$. Then $u, v \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. Hence the product of L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_2(i)$.

Conversely, let the product of L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - T_2(i)$. Take any coordinate space (X_j, τ_j) , choose $x_j, y_j \in X_j, x_j \neq y_j$. Now construct $x, y \in X$ such that $x = \Pi x'_i, y = \Pi y'_i$ where $x'_i = y'_i$ for $i \neq j$ and $x'_j = x_j, y'_j = y_j$. Then $x \neq y$ and using the product space $L - T_2(i) \exists u, v \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. Now choose any L-fuzzy point x_r in u . Then \exists a basic open L-fuzzy set $\Pi u_j^r \in \Pi \tau_j$ such that $x_r \in \Pi u_j^r \subseteq u$ which implies that $r < \Pi u_j^r(x)$ or that $r < \inf_j u_j^r(x'_j)$ and hence $r < \Pi u_j^r(x'_j) \forall j \in \Lambda \dots \dots (i)$ and $u(y) = 0 \Rightarrow \Pi u_j(y) = 0 \dots \dots (ii)$.

Similarly, corresponding to a fuzzy point $y_s \in v$ there exists a basic fuzzy open set $\Pi v_j^s \in \Pi \tau_j$ such that $y_s \in \Pi v_j^s \subseteq v$ which implies that

$s < v_j^s(j) \forall j \in \Lambda \dots \dots (iii)$ and

$\Pi v_j^s(y) = 0 \dots \dots (iv)$. Further, $\Pi u_j^r(y) = 0 \Rightarrow u_i^r(y_i) = 0$, since for $j \neq i, x'_j = y'_j$ and hence from (i), $u_j^r(y_j) = u_j^r(x_j) > r$. Similarly, $\Pi v_j^s(x) = 0 \Rightarrow v_i^s(x_i) = 0$ using (iii). Thus we have $u_i^r(x_i) > r, u_i^r(y_i) = 0$ and $v_i^s(y_i) > s, v_i^s(x_i) = 0$.

Now consider $\sup_r u_i^r = u_i, \sup_s v_i^s = v_i \in \tau_i$ then $u_i(x_i) = 1, u_i(y_i) = 0, v_i(x_i) = 0, v_i(y_i) = 1$ and $u_i \cap v_i = 0$, showing that (X_i, τ_i) is $L - T_2(i)$.

Moreover one can verify that

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(ii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(ii).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(iii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(iii).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(iv) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(iv).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(v) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(v).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(vi) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(vi).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - T_2(vii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - T_2(vii).$$

Hence it is seen that $L - T_2(i), L - T_2(ii), L - T_2(iii), L - T_2(iv),$

$L - T_2(v), L - T_2(vi), L - T_2(vii)$ Properties are productive and projective.

4.4 Mapping in L-topological spaces

We showed that $L - T_2(j)$ property is preserved under one-one, onto and continuous mapping for $j = i, ii, iii, iv, v, vi, vii$.

4.4.1 Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, onto and L-open map, then-

$$(a) (X, \tau) \text{ is } L - T_2(i) \Rightarrow (Y, s) \text{ is } L - T_2(i).$$

(b) (X, τ) is $L - T_2(ii) \Rightarrow (Y, s)$ is $L - T_2(ii)$.

(c) (X, τ) is $L - T_2(iii) \Rightarrow (Y, s)$ is $L - T_2(iii)$.

(d) (X, τ) is $L - T_2(iv) \Rightarrow (Y, s)$ is $L - T_2(iv)$.

(e) (X, τ) is $L - T_2(v) \Rightarrow (Y, s)$ is $L - T_2(v)$.

(f) (X, τ) is $L - T_2(vi) \Rightarrow (Y, s)$ is $L - T_2(vi)$.

(g) (X, τ) is $L - T_2(vii) \Rightarrow (Y, s)$ is $L - T_2(vii)$.

Proof: Suppose (X, τ) is $L - T_2(i)$. We shall prove that (Y, s) is $L - T_2(i)$.

Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$. Since f is onto, $\exists x_1, x_2 \in X$ such that $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one. Again since (X, τ) is $L - T_2(i) \exists u, v \in \tau$ such that $u(x_1) = 1, u(x_2) = 0, v(x_1) = 0,$

$v(x_2) = 1$ and $u \cap v = 0$.

Now

$$f(u)(y_1) = \{\sup u(x_1) : f(x_1) = y_1\} = 1$$

$$f(u)(y_2) = \{\sup u(x_2) : f(x_2) = y_2\} = 0$$

$$f(v)(y_1) = \{\sup v(x_1) : f(x_1) = y_1\} = 0$$

$$f(v)(y_2) = \{\sup v(x_2) : f(x_2) = y_2\} = 1$$

and

$$f(u \cap v)(y_1) = \{\sup(u \cap v)(x_1) : f(x_1) = y_1\}$$

$$f(u \cap v)(y_2) = \{\sup(u \cap v)(x_2) : f(x_2) = y_2\}$$

Hence $f(u \cap v) = 0 \Rightarrow f(u) \cap f(v) = 0$

Since f is L-open, $f(u), f(v) \in s$. Now it is clear that $\exists f(u), f(v) \in s$ such that $f(u)(y_1) = 1, f(u)(y_2) = 0, f(v)(y_1) = 0, f(v)(y_2) = 1$ and $f(u) \cap f(v) = 0$. Hence it is clear that the L-topological space (Y, s) is $L - T_2(i)$. Similarly (b), (c), (d), (e), (f), (g) can be proved.

4.4.2 Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be L-continuous and one-one map, then-

- (a) (Y, s) is $L - T_2(i) \Rightarrow (X, \tau)$ is $L - T_2(i)$.
- (b) (Y, s) is $L - T_2(ii) \Rightarrow (X, \tau)$ is $L - T_2(ii)$.
- (c) (Y, s) is $L - T_2(iii) \Rightarrow (X, \tau)$ is $L - T_2(iii)$.
- (d) (Y, s) is $L - T_2(iv) \Rightarrow (X, \tau)$ is $L - T_2(iv)$.
- (e) (Y, s) is $L - T_2(v) \Rightarrow (X, \tau)$ is $L - T_2(v)$.
- (f) (Y, s) is $L - T_2(vi) \Rightarrow (X, \tau)$ is $L - T_2(vi)$.
- (g) (Y, s) is $L - T_2(vii) \Rightarrow (X, \tau)$ is $L - T_2(vii)$.

Proof: Suppose (Y, s) is $L - T_2(i)$. We shall prove that (X, τ) is $L - T_2(i)$.

Let $x_1, x_2 \in X$ with $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$ as f is one-one. Since (Y, s) is $L - T_2(i), \exists u, v \in s$ such that $u(f(x_1)) = 1, u(f(x_2)) = 0, v(f(x_1)) = 0, v(f(x_2)) = 1$ and $u \cap v = 0$. This implies that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0, f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$ and $f^{-1}(u \cap v) = 0 \Rightarrow f^{-1}(u) \cap f^{-1}(v) = 0$. Hence $f^{-1}(u), f^{-1}(v) \in \tau$ as f is L-continuous and $u, v \in s$. Now it is clear that $f^{-1}(u), f^{-1}(v) \in \tau$ such

that $f^{-1}(u)(x_1) = 1$, $f^{-1}(u)(x_2) = 0$, $f^{-1}(v)(x_1) = 0$, $f^{-1}(v)(x_2) = 1$ and $f^{-1}(u) \cap f^{-1}(v) = 0$. Hence the L-topological space (X, τ) is $L - T_2(i)$. Similarly (b), (c), (d), (e), (f), (g) can be proved.

Chapter-5

On R_0 Space in L-Topological Spaces

5.1: Introduction

The concept of R_0 -property first defined by Shanin[64] and there after Dude[16], Naimpally[57], Dorsett[15], Caldas[10], Ekici[18], as earlier Keskin[38] and Roy[62] defined many characterizations of R_0 -properties. The concepts of fuzzy R_0 -propertise are established and discussed by Hutton[33,34], Srivastava[78], Ali[9], Khedr[40], Zhang[92] and many other fuzzy topologist. In this chapter we define possible eight definitions of R_0 space in L-topological spaces and we show that this space possesses many nice properties which are hereditary, productive and projective.

5.2 R_0 -property in L-Topological Spaces

We now give the following definitions of R_0 -property in L-topological spaces.

5.2.1Definition: An lts (X, τ) is called-

- (a) $L - R_0(i)$ if $\forall x, y \in X, x \neq y$ whenever $\exists u \in \tau$ with $u(x) \neq u(y)$ then $\exists v \in \tau$ such that $v(x) \neq v(y)$.
- (b) $L - R_0(ii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists u \in \tau$ with $u(x) = 1, u(y) = 0$ then $\exists v \in \tau$ such that $v(x) = 0, v(y) = 1$.

- (c) $L - R_0(iii)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ whenever $\exists u \in \tau$ with $x_r \in u, y_s \notin u$ then $\exists v \in \tau$ such that $x_r \notin v, y_s \in v$.
- (d) $L - R_0(iv)$ if for all pairs of distinct L-fuzzy singletons $x_r, y_s \in S(X)$ and $x_r \bar{q} y_s$ whenever $\exists u \in \tau$ with $x_r \subseteq u, y_s \bar{q} u$ then $\exists v \in \tau$ such that $y_s \subseteq v, x_r \bar{q} v$.
- (e) $L - R_0(v)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ whenever $\exists u \in \tau$ with $x_r \in u, u \bar{q} y_s$ then $\exists v \in \tau$ such that $y_s \in v, v \bar{q} x_r$.
- (f) $L - R_0(vi)$ if for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ whenever $\exists u \in \tau$ with $x_r \in u, y_s \cap u = 0$ then $\exists v \in \tau$ such that $y_s \in v, x_r \cap v = 0$.
- (g) $L - R_0(vii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists u \in \tau$ with $u(x) > 0, u(y) = 0$ then $\exists v \in \tau$ such that $v(x) = 0, v(y) > 0$.
- (h) $L - R_0(viii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists u \in \tau$ with $u(x) > u(y)$ then $\exists v \in \tau$ such that $v(y) > v(x)$.

5.3 “Good extension”, Hereditary, Productive and Projective Properties in L-Topology

Now all the definitions $L - R_0(i), L - R_0(ii), L - R_0(iii), L - R_0(iv),$

$L - R_0(v), L - R_0(vi), L - R_0(vii)$ and $L - R_0(viii)$ are ‘good extensions’ of

R_0 – property, is shown below:

5.3.1Theorem: Let (X, T) be a topological space. Then (X, T) is R_0 iff $(X, \omega(T))$ is $L - R_0(j)$, for $j = i, ii, iii, iv, v, vi, vii, viii$.

Proof: Let the topological space (X, T) be R_0 , we shall prove that the fuzzy topological space $(X, \omega(T))$ is $L - R_0(ii)$. Choose $x, y \in X$ with $x \neq y$. Let $u \in \omega(T)$ with $u(x) = 1, u(y) = 0$, then it is clear that $u^{-1}(r, 1] \in T$, for any $r \in I_1$ and $x \in u^{-1}(r, 1], y \notin u^{-1}(r, 1]$. Since (X, T) is R_0 , then there exist $V \in T$ with $x \notin V, y \in V$. Now consider the characteristics function 1_V . We see that $1_V \in \omega(T)$ with $1_V(x) = 0, 1_V(y) = 1$. Thus $(X, \omega(T))$ is $L - R_0(ii)$.

Conversely, let $(X, \omega(T))$ be $L - R_0(ii)$, we shall prove that (X, T) is R_0 . Choose $x, y \in X, x \neq y$ and $U \in T$, with $x \in U, y \notin U$, but we know that the characteristic function $1_U \in \omega(T)$. Also it is clear that $1_U(x) = 1, 1_U(y) = 0$. Since $(X, \omega(T))$ is $L - R_0(ii)$, then $\exists v \in \omega(T)$ such that $v(x) = 0, v(y) = 1$. Again since v is lower semi continuous function then $v^{-1}(0, 1] \in T$ and from above, we get $x \notin v^{-1}(0, 1], y \in v^{-1}(0, 1]$. Hence (X, T) is R_0 .

Similarly we can show that $L - R_0(i), L - R_0(iii), L - R_0(iv), L - R_0(v),$

$L - R_0(vi), L - R_0(vii), L - R_0(viii)$ are also hold 'good extension' property.

5.3.2Theorem: Let (X, τ) be an lts, $A \subseteq X$ and $\tau_A = \{u|A : u \in \tau\}$, then

(a) (X, τ) is $L - R_0(i) \Rightarrow (A, \tau_A)$ is $L - R_0(i)$.

(b) (X, τ) is $L - R_0(ii) \Rightarrow (A, \tau_A)$ is $L - R_0(ii)$.

- (c) (X, τ) is $L - R_0(iii) \Rightarrow (A, \tau_A)$ is $L - R_0(iii)$.
- (d) (X, τ) is $L - R_0(iv) \Rightarrow (A, \tau_A)$ is $L - R_0(iv)$.
- (e) (X, τ) is $L - R_0(v) \Rightarrow (A, \tau_A)$ is $L - R_0(v)$.
- (f) (X, τ) is $L - R_0(vi) \Rightarrow (A, \tau_A)$ is $L - R_0(vi)$.
- (g) (X, τ) is $L - R_0(vii) \Rightarrow (A, \tau_A)$ is $L - R_0(vii)$.
- (h) (X, τ) is $L - R_0(viii) \Rightarrow (A, \tau_A)$ is $L - R_0(viii)$.

Proof: We prove only (b). Suppose (X, τ) is L -topological space and $L - R_0(ii)$. We shall prove (A, τ_A) is $L - R_0(ii)$. Let $x, y \in A$, $x \neq y$, and $w \in \tau_A$ with $w(x) = 1$, $w(y) = 0$. Then $x, y \in X$ with $x \neq y$ as $A \subseteq X$. Consider u be the extension function of w on the set X , then it is clear that $u(x) = 1, u(y) = 0$. Since (X, τ) is $L - R_0(ii)$. Then $\exists v \in \tau$ such that $v(x) = 0, v(y) = 1$. For $A \subseteq X$, we find $\exists v|A \in \tau_A$ such that $v|A(x) = 0, v|A(y) = 1$ as $x, y \in A$. Hence it is clear that the subspace (A, τ_A) is $L - R_0(ii)$. Similarly, (a), (c), (d), (e), (f), (g), (h) can be proved.

5.3.3 Theorem: Given $\{(X_i, \tau_i): i \in \Lambda\}$ be a family of L -topological space. Then the product of L -topological space $(\Pi X_i, \Pi \tau_i)$ is $L - R_0(j)$ iff each coordinate space (X_i, τ_i) is $L - R_0(j)$ where $j = i, ii, iii, iv, v, vi, vii, viii$.

Proof: Let each coordinate space $\{(X_i, \tau_i): i \in \Lambda\}$ be $L - R_0(ii)$. Then we show that the product space is $L - R_0(ii)$. Suppose $x, y \in \Pi X_i$, $x \neq y$, and $u \in \Pi \tau_i$ with $u(x) = 1, u(y) = 0$. Choose $x = \Pi x_i$, $y = \Pi y_i$, but we have $u(x) = \min\{u_i(x_i), \text{ for } i \in \Lambda \text{ and } u_i \in \tau_i\}$,

$u(y) = \min\{u_i(y_i), \text{for } i \in \Lambda \text{ and } u_i \in \tau_i\}$, then there exist at least one $j \in \Lambda$, such that $x_j \neq y_j$ and $u_j(x_j) = 1, u_j(y_j) = 0$. Since (X_j, τ_j) is $L - R_0(ii)$ for each $j \in \Lambda$, then $\exists v_j \in \tau_j$ such that $v_j(x_j) = 0, v_j(y_j) = 1$. Now take, $v = \Pi v'_j$ where $v'_j = v_j$ and $v_i = 1$ for $i \neq j$. then $\exists v \in \Pi \tau_i$ such that $v(x) = 0, v(y) = 1$. Hence the product of L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - R_0(ii)$.

Conversely, let the product of L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - R_0(ii)$. We shall prove the each coordinate space (X_j, τ_j) is also $L - R_0(ii)$. Choose $x_j, y_j \in X_j, x_j \neq y_j$ and $u_j \in \tau_j$ with $u_j(x_j) = 1, u_j(y_j) = 0$. Now construct $x, y \in X$ such that $x = \Pi x'_i, y = \Pi y'_i$ where $x'_i = y'_i$ for $i \neq j$ and $x'_j = x_j, y'_j = y_j$. Then $x \neq y$. Further, let $\pi_j: X \rightarrow X_j$ be a projection map from X into X_j . Now, we observe that $u_j((\pi_j)(x)) = u_j(x_j) = 1, u_j((\pi_j)(y)) = u_j(y_j) = 0$, i.e for $u_j \circ \pi_j \in \Pi \tau_i$, with $(u_j \circ \pi_j)(x) = 1, (u_j \circ \pi_j)(y) = 0$. Since the product space $(\Pi X_i, \Pi \tau_i)$ is $L - R_0(ii)$. Then $\exists v \in \Pi \tau_i$ such that $v(x) = 0, v(y) = 1$. Now choose any L-fuzzy point y_r in v . Then \exists a basic open L-fuzzy set $\Pi v_j^r \in \Pi \tau_j$ such that $y_r \in \Pi v_j^r \subseteq v$ which implies that $r < \Pi v_j^r(y)$ or that $r < \inf_j v_j^r(y'_j)$ and hence $r < \Pi v_j^r(y'_j) \forall j \in \Lambda \dots (i)$ and $v(x) = 0 \Rightarrow \Pi v_j(x) = 0 \dots (ii)$. Further, $\Pi v_j^r(x) = 0 \Rightarrow v_i^r(x_i) = 0$, since for $j \neq i, x'_j = y'_j$ and hence from (i), $u_j^r(x_j) = v_j^r(y_j) > r$. Thus we have $v_i^r(y_i) > r, v_i^r(x_i) = 0$.

Now consider $\sup_r v_i^r = v_i$ then $v_i \in \tau_i$ with $v_i(y_i) = 1, v_i(x_i) = 0$. This showing that (X_i, τ_i) is $L - R_0(ii)$.

Moreover one can easily verify that

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(i) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(i)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(iii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(iii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(iv) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(iv)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(v) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(v)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(vi) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(vi)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(vii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(vii)$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_0(viii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_0(viii).$$

Hence we see that $L - R_0(i), L - R_0(ii), L - R_0(iii), L - R_0(iv),$

$L - R_0(v), L - R_0(vi), L - R_0(vii), L - R_0(viii)$ Properties are productive and projective.

5.4 Mapping in L-topological spaces

We show that $L - R_0(j)$ property is preserved under one-one, onto and continuous mapping for $j = i, ii, iii, iv, v, vi, vii, viii$.

5.4.1 Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, onto, L-continuous and L-open map, then-

- (a) (X, τ) is $L - R_0(i) \Rightarrow (Y, s)$ is $L - R_0(i)$.
- (b) (X, τ) is $L - R_0(ii) \Rightarrow (Y, s)$ is $L - R_0(ii)$.
- (c) (X, τ) is $L - R_0(iii) \Rightarrow (Y, s)$ is $L - R_0(iii)$.
- (d) (X, τ) is $L - R_0(iv) \Rightarrow (Y, s)$ is $L - R_0(iv)$.
- (e) (X, τ) is $L - R_0(v) \Rightarrow (Y, s)$ is $L - R_0(v)$.
- (f) (X, τ) is $L - R_0(vi) \Rightarrow (Y, s)$ is $L - R_0(vi)$.
- (g) (X, τ) is $L - R_0(vii) \Rightarrow (Y, s)$ is $L - R_0(vii)$.
- (h) (X, τ) is $L - R_0(viii) \Rightarrow (Y, s)$ is $L - R_0(viii)$.

Proof: Suppose (X, τ) is $L - R_0(ii)$. We shall prove that (Y, s) is

$L - R_0(ii)$. Let $y_1, y_2 \in Y$, $y_1 \neq y_2$ and $u \in s$ with $u(y_1) = 1, u(y_2) = 0$.

Since f is onto, $\exists x_1, x_2 \in X$ such that $f(x_1) = y_1, f(x_2) = y_2$ and $x_1 \neq x_2$ as f is one-one.

Now $f^{-1}(u)(x_1) = u(f(x_1)) = u(y_1) = 1$ and

$$f^{-1}(u)(x_2) = u(f(x_2)) = u(y_2) = 0$$

Since f is L -continuous then $f^{-1}(u) \in t$ and $f^{-1}(u)(x_1) = 1$,

$f^{-1}(u)(x_2) = 0$. Since (X, τ) is $L - R_0(ii)$, then $\exists v \in \tau$ such that $v(x) = 0, v(y) = 1$.

Now

$$f(v)(y_1) = \{\sup v(x_1) : f(x_1) = y_1\} = 0$$

$$f(v)(y_2) = \{\sup v(x_2) : f(x_2) = y_2\} = 1.$$

Since f is L -open, $f(u) \in s$. Now it is clear that $\exists f(v) \in s$ such that $f(v)(y_1) = 0, f(v)(y_2) = 1$. Hence it is clear that the L -topological space (Y, s) is $L - R_0(ii)$.

Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

5.4.2Theorem: Let (X, τ) and (Y, s) be two L -topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, L -continuous and L -open map, then-

- (a) (Y, s) is $L - R_0(i) \Rightarrow (X, \tau)$ is $L - R_0(i)$.
- (b) (Y, s) is $L - R_0(ii) \Rightarrow (X, \tau)$ is $L - R_0(ii)$.
- (c) (Y, s) is $L - R_0(iii) \Rightarrow (X, \tau)$ is $L - R_0(iii)$.
- (d) (Y, s) is $L - R_0(iv) \Rightarrow (X, \tau)$ is $L - R_0(iv)$.
- (e) (Y, s) is $L - R_0(v) \Rightarrow (X, \tau)$ is $L - R_0(v)$.
- (f) (Y, s) is $L - R_0(vi) \Rightarrow (X, \tau)$ is $L - R_0(vi)$.
- (g) (Y, s) is $L - R_0(vii) \Rightarrow (X, \tau)$ is $L - R_0(vii)$.
- (h) (Y, s) is $L - R_0(viii) \Rightarrow (X, \tau)$ is $L - R_0(viii)$.

Proof: Suppose (Y, s) is $L - R_0(ii)$. We shall prove that (X, τ) is

$L - R_0(ii)$. Let $x_1, x_2 \in X, x_1 \neq x_2$ and $u \in \tau$ with $u(x_1) = 1, u(x_2) = 0$.

Since f is one-one map then $f(x_1) \neq f(x_2)$.

Now $f(u)(f(x_1)) = \sup\{u(x_1)\} = 1$ as f is one-one

And $f(u)(f(x_2)) = \sup\{u(x_2)\} = 0$

So, we have $f(u) \in s$, with $f(u)(f(x_1)) = 1$, $f(u)(f(x_2)) = 0$, as f is L-open map. Since (Y, s) is $L - R_0(ii)$, then $\exists v \in s$ such that $v(f(x_1)) = 0, v(f(x_2)) = 1$. This implies that $f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$ and $f^{-1}(v) \in \tau$ as f is L-continuous and $v \in s$. Now it is clear that $\exists f^{-1}(v) \in \tau$ such that $f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$. Hence the L-topological space (X, τ) is $L - R_0(ii)$.

Similarly (a), (c), (d), (e), (f), (g), (h) can be proved.

Chapter-6

On R_1 Space in L-Topological Spaces

6.1: Introduction

The concept of R_1 -property first defined by Yang [87] and there after Murdeshwar[56], Dorset[15], Dude[17], Caldas[10], Ekici[18], as earlier Keskin[38] and Roy[62] defined many characterizations of R_1 -properties. The concepts of fuzzy R_1 -propertise are established and discussed by Hutton[31,32], Srivastava[76], Ali[8], Khedr[40], Kandil[36], Hossain[28] and many other fuzzy topologist.

In this chapter we define possible seven definitions of R_1 space in L-topological spaces. All these definitions satisfy ‘good extension’ property and we establish some implications among them. Finally we show that all these definitions are hereditary, productive and projective and preserved under one-one, onto and continuous maps.

6.2 R_1 -property in L-Topological Spaces

We now give the following definitions of R_1 -property in L-topological spaces.

6.2.1 Definition: An lts (X, τ) is called-

- (a) $L - R_1(i)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then $\exists u, v \in \tau$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$.
- (b) $L - R_1(ii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ and $\exists u, v \in \tau$ such that $x_r \in u, y_s \notin u$ and $x_r \notin v, y_s \in v, u \cap v = 0$.
- (c) $L - R_1(iii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then for all pairs of distinct L-fuzzy singletons $x_r, y_s \in S(X), x_r \bar{q} y_s$ and $\exists u, v \in \tau$ such that $x_r \subseteq u, y_s \bar{q} u$ and $y_s \subseteq v, x_r \bar{q} v$ and $u \cap v = 0$.
- (d) $L - R_1(iv)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ and $\exists u, v \in \tau$ such that $x_r \in u, u \bar{q} y_s$ and $y_s \in v, v \bar{q} x_r$ and $u \cap v = 0$.
- (e) $L - R_1(v)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ and for any pair of distinct L-fuzzy points $x_r, y_s \in S(X)$ and $\exists u, v \in \tau$ such that $x_r \in u \subseteq coy_s, y_s \in v \subseteq cox_r$ and $u \subseteq cov$.
- (f) $L - R_1(vi)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then $\exists u, v \in \tau$ such that $u(x) > 0, u(y) = 0$ and $v(x) = 0, v(y) > 0$.

(g) $L - R_1(vii)$ if $\forall x, y \in X, x \neq y$ whenever $\exists w \in \tau$ with $w(x) \neq w(y)$ then $\exists u, v \in \tau$ such that $u(x) > u(y)$ and $v(y) > v(x)$.


Here, we established a complete comparison of the definitions

$L - R_1(ii), L - R_1(iii), L - R_1(iv), L - R_1(v), L - R_1(vi)$ and

$L - R_1(vii)$ with $L - R_1(i)$.

6.2.2Theorem: Let (X, τ) be an lts. Then we have the following

implications:



The reverse implications are not true in general except $L - R_1(vi)$ and

$L - R_1(vii)$.

Proof: $L - R_1(i) \Rightarrow L - R_1(ii), L - R_1(i) \Rightarrow L - R_1(iii)$ can be proved easily.

Now $L - R_1(i) \Rightarrow L - R_1(iv)$ and $L - R_1(i) \Rightarrow L - R_1(v)$, since $L - R_1(ii) \Leftrightarrow$

$L - R_1(iv)$ and $L - R_1(iv) \Leftrightarrow L - R_1(v)$. $L - R_1(i) \Rightarrow L - R_1(vi)$; It is

obvious. $L - R_1(i) \Rightarrow L - R_1(vii)$, since $L - R_1(vi) \Rightarrow L - R_1(vii)$.

The reverse implications are not true in general except $L - R_1(vi)$ and

$L - R_1(vii)$, it can be seen through the following counter examples:

Example-1: Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u, v, w\}$ where $w(x) = 0.6, w(y) = 0.7, u(x) = 0.5, u(y) = 0, v(x) = 0, v(y) = 0.6, L = \{0, 0.05, 0.1, 0.15, \dots \dots 0.95, 1\}$ and $r = 0.4, s = 0.3$.

Example-2: Let $X = \{x, y\}$, τ be the L-topology on X generated by $\{\alpha: \alpha \in L\} \cup \{u, v, w\}$ where $w(x) = 0.8, w(y) = 0.9, u(x) = 0.5, u(y) = 0, v(x) = 0, v(y) = 0.4, L = \{0, 0.05, 0.1, 0.15, \dots \dots 0.95, 1\}$ and $r = 0.5, s = 0.4$.

Proof: $L - R_1(ii) \not\Rightarrow L - R_1(i)$: From example-1, we see that the lts (X, τ) is clearly $L - R_1(ii)$ but it is not $L - R_1(i)$. Since there is no L-fuzzy set in τ which grade of membership is 1.

$L - R_1(iii) \not\Rightarrow L - R_1(i)$: From example-2, we see the lts (X, τ) is clearly $L - R_1(iii)$ but it is not $L - R_1(ii)$. Since $L - R_1(iii) \not\Rightarrow L - R_1(ii)$ and $L - R_1(ii) \not\Rightarrow L - R_1(i)$ so $L - R_1(iii) \not\Rightarrow L - R_1(i)$.

$L - R_1(iv) \not\Rightarrow L - R_1(i)$: This follows automatically from the fact that

$L - R_1(ii) \Leftrightarrow L - R_1(iv)$ and it has already been shown that $L - R_1(ii) \not\Rightarrow$

$L - R_1(i)$ so $L - R_1(iv) \not\Rightarrow L - R_1(i)$.

$L - R_1(v) \not\Rightarrow L - R_1(i)$: Since $L - R_1(iv) \Leftrightarrow L - R_1(v)$ and $L - R_1(iv) \not\Rightarrow L - R_1(i)$ so $L - R_1(v) \not\Rightarrow L - R_1(i)$. But $L - R_1(vii) \Rightarrow L - R_1(vi) \Rightarrow$

$L - R_1(i)$ is obvious.

6.3 “Good extension”, Hereditary, Productive and Projective Properties in L-Topology

We show that all definitions $L - R_1(i), L - R_1(ii), L - R_1(iii),$

$L - R_1(iv), L - R_1(v), L - R_1(vi)$ and $L - R_1(vii)$ are ‘good extensions’ of R_1 – property, is shown below:

6.3.1Theorem: Let (X, T) be a topological space. Then (X, T) is R_1 iff $(X, \omega(T))$ is $L - R_1(j)$, where $j = i, ii, iii, iv, v, vi, vii$.

Proof: Let (X, T) be R_1 . Choose $x, y \in X, x \neq y$. Whenever $\exists W \in T$ with $x \in W, y \notin W$ or $x \notin W, y \in W$ then $\exists U, V \in T$ such that $x \in U, y \notin U$ and $y \in V, x \notin V$ and $U \cap V = \emptyset$. Suppose $x \in W, y \notin W$ since $W \in T$ then $1_w \in \omega(T)$ with $1_w(x) \neq 1_w(y)$. Also consider the lower semi continuous function $1_U, 1_V$, then $1_U, 1_V \in \omega(T)$ such that $1_U(x) = 1, 1_U(y) = 0$ and $1_V(x) = 0, 1_V(y) = 1$ and so that $1_U \cap 1_V = 0$ as $U \cap V = \emptyset$. Thus $(X, \omega(T))$ is $L - R_1(i)$.

Conversely, let $(X, \omega(T))$ be $L - R_1(i)$. To show that (X, T) is R_1 . Choose $x, y \in X$ with $x \neq y$. Whenever $\exists w \in T$ with $w(x) \neq w(y)$ then $\exists u, v \in \omega(T)$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. Since $w(x) \neq w(y)$, then either $w(x) < w(y)$ or $w(x) > w(y)$. Choose $w(x) < w(y)$, then $\exists s \in L$ such that $w(x) < s < w(y)$. So it is clear that $w^{-1}(s, 1] \in T$ and $x \notin w^{-1}(s, 1], y \in w^{-1}(s, 1]$. Let $U = u^{-1}\{1\}$ and $V =$

$v^{-1}\{1\}$, then $U, V \in T$ and is $x \in U, y \notin U, x \notin V, y \in V$, and $U \cap V = \emptyset$ as $u \cap v = 0$. Hence (X, T) is R_1 .

Similarly, we can show that $L - R_1(ii), L - R_1(iii), L - R_1(iv),$

$L - R_1(v), L - R_1(vi), L - R_1(vii)$ are also hold 'good extension' property.

6.3.2 Theorem: Let (X, τ) be an lts, $A \subseteq X$ and $\tau_A = \{u|A : u \in \tau\}$, then

- (a) (X, τ) is $L - R_1(i) \Rightarrow (A, \tau_A)$ is $L - R_1(i)$.
- (b) (X, τ) is $L - R_1(ii) \Rightarrow (A, \tau_A)$ is $L - R_1(ii)$.
- (c) (X, τ) is $L - R_1(iii) \Rightarrow (A, \tau_A)$ is $L - R_1(iii)$.
- (d) (X, τ) is $L - R_1(iv) \Rightarrow (A, \tau_A)$ is $L - R_1(iv)$.
- (e) (X, τ) is $L - R_1(v) \Rightarrow (A, \tau_A)$ is $L - R_1(v)$.
- (f) (X, τ) is $L - R_1(vi) \Rightarrow (A, \tau_A)$ is $L - R_1(vi)$.
- (g) (X, τ) is $L - R_1(vii) \Rightarrow (A, \tau_A)$ is $L - R_1(vii)$.

Proof: We prove only (a). Suppose (X, τ) is L-topological space and is also $L - R_1(i)$. We shall prove that (A, τ_A) is $L - R_1(i)$. Let $x, y \in A$ with $x \neq y$ and $\exists w \in \tau_A$ such that $w(x) \neq w(y)$, then $x, y \in X$ with $x \neq y$ as $A \subseteq X$. Consider m be the extension function of w on X , then $m(x) \neq m(y)$. Since (X, τ) is $L - R_1(i)$, $\exists u, v \in \tau$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. For $A \subseteq X$, we find $u|A, v|A \in \tau_A$ and $u|A(x) = 1, u|A(y) = 0$ and $v|A(x) = 0, v|A(y) = 1$ and $u|A \cap v|A = (u \cap v)|A = 0$ as $x, y \in A$. Hence it is clear that the subspace (A, τ_A) is $L - R_1(i)$.

Similarly, (b), (c), (d), (e), (f), (g) can be proved.

6.3.3Theorem: Given $\{(X_i, \tau_i): i \in \Lambda\}$ be a family of L-topological space. Then the product of L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - R_1(j)$ iff each coordinate space (X_i, τ_i) is $L - R_1(j)$, where $j = i, ii, iii, iv, v, vi, vii$.

Proof: Let each coordinate space $\{(X_i, \tau_i): i \in \Lambda\}$ be $L - R_1(i)$. Then we show that the product space is $L - R_1(i)$. Suppose $x, y \in X$ with $x \neq y$ and $w \in \Pi \tau_i$ with $w(x) \neq w(y)$, again suppose $x = \Pi x_i, y = \Pi y_i$ then $x_j \neq y_j$ for some $j \in \Lambda$. But we have $w(x) = \min \{w_i(x_i): i \in \Lambda\}$, and $w(y) = \min \{w_i(y_i): i \in \Lambda\}$. Hence we can find at least one $w_j \in \tau_j$ with $w_j(x_j) \neq w_j(y_j)$, since each $(X_i, \tau_i): i \in \Lambda$ be $L - R_1(i)$ then $\exists u_j, v_j \in \tau_j$ such that $u_j(x_j) = 1, u_j(y_j) = 0, v_j(x_j) = 0, v_j(y_j) = 1$ and $u_j \cap v_j = 0$. Now take $u = \Pi u'_j, v = \Pi v'_j$ where $u'_j = u_j, v'_j = v_j$ and $u_i = v_i = 1$ for $i \neq j$. Then $u, v \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. Hence the product of L-topological space is also L-topological space and $(\Pi X_i, \Pi \tau_i)$ is $L - R_1(i)$.

Conversely, let the product L-topological space $(\Pi X_i, \Pi \tau_i)$ is $L - R_1(i)$. Take any coordinate space (X_j, τ_j) , choose $x_j, y_j \in X_j, x_j \neq y_j$ and $w_i \in \Pi \tau_i$ with $w_i(x_i) \neq w_i(y_i)$. Now construct $x, y \in X$ such that $x = \Pi x'_i, y = \Pi y'_i$ where $x'_i = y'_i$ for $i \neq j$ and $x'_j = x_j, y'_j = y_j$. Then $x \neq y$ and using the product space $L - R_1(i)$, $\Pi w_i \in \Pi \tau_i$ with $\Pi w_i(x_i) \neq \Pi w_i(y_i)$, since

$(\Pi X_i, \Pi \tau_i)$ is $L - R_1(i)$ then $\exists u, v \in \Pi \tau_i$ such that $u(x) = 1, u(y) = 0, v(x) = 0, v(y) = 1$ and $u \cap v = 0$. Now choose any L-fuzzy point x_r in u . Then \exists a basic open L-fuzzy set $\Pi u_j^r \in \Pi \tau_j$ such that $x_r \in \Pi u_j^r \subseteq u$ which implies that $r < \Pi u_j^r(x)$ or that $r < \inf_j u_j^r(x'_j)$

and hence $r < \Pi u_j^r(x'_j) \forall j \in \Lambda \dots \dots (i)$ and

$$u(y) = 0 \Rightarrow \Pi u_j(y) = 0 \dots \dots (ii).$$

Similarly, corresponding to a fuzzy point $y_s \in v$ there exists a basic fuzzy open set $\Pi v_j^s \in \Pi \tau_j$ such that $y_s \in \Pi v_j^s \subseteq v$ which implies that

$$s < v_j^s(j) \forall j \in \Lambda \dots \dots (iii) \text{ and}$$

$\Pi v_j^s(y) = 0 \dots \dots (iv)$. Further, $\Pi u_j^r(y) = 0 \Rightarrow u_i^r(y_i) = 0$, since for $j \neq i, x'_j = y'_j$ and hence from (i), $u_j^r(y_j) = u_j^r(x_j) > r$. Similarly, $\Pi v_j^s(x) = 0 \Rightarrow v_i^s(x_i) = 0$ using (iii).

Thus we have $u_i^r(x_i) > r, u_i^r(y_i) = 0$ and $v_i^s(y_i) > s, v_i^s(x_i) = 0$. Now consider $\sup_r u_i^r = u_i, \sup_s v_i^s = v_i$, then $u_i(x_i) = 1, u_i(y_i) = 0, v_i(x_i) = 0, v_i(y_i) = 1$ and $u_i \cap v_i = 0$, showing that (X_i, τ_i) is $L - R_1(i)$.

Moreover one can easily verify that

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(ii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(ii).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(iii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(iii).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(iv) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(iv).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(v) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(v).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(vi) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(vi).$$

$$(X_i, \tau_i), i \in \Lambda \text{ is } L - R_1(vii) \Leftrightarrow (\Pi X_i, \Pi \tau_i) \text{ is } L - R_1(vii).$$

Hence, we see that $L - R_1(i), L - R_1(ii), L - R_1(iii), L - R_1(iv),$

$L - R_1(v), L - R_1(vi), L - R_1(vii)$ Properties are productive and projective.

6.4 Mapping in L-topological spaces

We show that $L - R_1(j)$ property is preserved under one-one, onto and continuous mapping for $j = i, ii, iii, iv, v, vi, vii$.

6.4.1 Theorem: Let (X, τ) and (Y, s) be two L-topological space and $f: (X, \tau) \rightarrow (Y, s)$ be one-one, onto L-continuous and L-open map, then-

- (a) $(X, \tau) \text{ is } L - R_1(i) \Rightarrow (Y, s) \text{ is } L - R_1(i).$
- (b) $(X, \tau) \text{ is } L - R_1(ii) \Rightarrow (Y, s) \text{ is } L - R_1(ii).$
- (c) $(X, \tau) \text{ is } L - R_1(iii) \Rightarrow (Y, s) \text{ is } L - R_1(iii).$
- (d) $(X, \tau) \text{ is } L - R_1(iv) \Rightarrow (Y, s) \text{ is } L - R_1(iv).$
- (e) $(X, \tau) \text{ is } L - R_1(v) \Rightarrow (Y, s) \text{ is } L - R_1(v).$
- (f) $(X, \tau) \text{ is } L - R_1(vi) \Rightarrow (Y, s) \text{ is } L - R_1(vi).$
- (g) $(X, \tau) \text{ is } L - R_1(vii) \Rightarrow (Y, s) \text{ is } L - R_1(vii).$

Proof: Suppose (X, τ) is $L - R_1(i)$. We shall prove that (Y, s) is $L - R_1(i)$. Let $y_1, y_2 \in Y$ with $y_1 \neq y_2$ and $w \in s$ with $w(y_1) \neq w(y_2)$. Since f is onto then $\exists x_1, x_2 \in X$ such that $f(x_1) = y_1$ and $f(x_2) = y_2$, also $x_1 \neq x_2$, as f is one-one. Now we have $f^{-1}(w) \in \tau$, Since f is L -continuous, also we have $f^{-1}(w)(x_1) = wf(x_1) = w(y_1)$ and $f^{-1}(w)(x_2) = wf(x_2) = w(y_2)$. Therefore $f^{-1}(w)(x_1) \neq f^{-1}(w)(x_2)$. Again since (X, τ) is $L - R_1(i)$ and $\exists f^{-1}(w) \in \tau$ with $f^{-1}(w)(x_1) \neq f^{-1}(w)(x_2)$ then $\exists u, v \in \tau$ such that $u(x_1) = 1, u(x_2) = 0, v(x_1) = 0, v(x_2) = 1$ and $u \cap v = 0$.

Now

$$f(u)(y_1) = \{\sup u(x_1) : f(x_1) = y_1\} = 1$$

$$f(u)(y_2) = \{\sup u(x_2) : f(x_2) = y_2\} = 0$$

$$f(v)(y_1) = \{\sup v(x_1) : f(x_1) = y_1\} = 0$$

$$f(v)(y_2) = \{\sup v(x_2) : f(x_2) = y_2\} = 1$$

And

$$f(u \cap v)(y_1) = \{\sup(u \cap v)(x_1) : f(x_1) = y_1\} = 0$$

$$f(u \cap v)(y_2) = \{\sup(u \cap v)(x_2) : f(x_2) = y_2\} = 0$$

$$\text{Hence } f(u \cap v) = 0 \Rightarrow f(u) \cap f(v) = 0$$

Since f is L -open, $f(u), f(v) \in s$. Now it is clear that $\exists f(u), f(v) \in s$ such that $f(u)(y_1) = 1, f(u)(y_2) = 0, f(v)(y_1) = 0, f(v)(y_2) = 1$ and

$f(u) \cap f(v) = 0$. Hence it is clear that the L-topological space (Y, s) is

$$L - R_1(i).$$

Similarly (b), (c), (d), (e), (f), (g) can be proved.

6.4.2Theorem: Let (X, τ) and (Y, s) be two L-topological spaces and $f: (X, \tau) \rightarrow (Y, s)$ be L-continuous and one-one map, then-

- (a) (Y, s) is $L - R_1(i) \Rightarrow (X, \tau)$ is $L - R_1(i)$.
- (b) (Y, s) is $L - R_1(ii) \Rightarrow (X, \tau)$ is $L - R_1(ii)$.
- (c) (Y, s) is $L - R_1(iii) \Rightarrow (X, \tau)$ is $L - R_1(iii)$.
- (d) (Y, s) is $L - R_1(iv) \Rightarrow (X, \tau)$ is $L - R_1(iv)$.
- (e) (Y, s) is $L - R_1(v) \Rightarrow (X, \tau)$ is $L - R_1(v)$.
- (f) (Y, s) is $L - R_1(vi) \Rightarrow (X, \tau)$ is $L - R_1(vi)$.
- (g) (Y, s) is $L - R_1(vii) \Rightarrow (X, \tau)$ is $L - R_1(vii)$.

Proof: Suppose (Y, s) is $L - R_1(i)$. We shall prove that (X, τ) is $L - R_1(i)$. Let $x_1, x_2 \in X$ with $x_1 \neq x_2$ and $w \in \tau$ with $w(x_1) \neq w(x_2)$, $\Rightarrow f(x_1) \neq f(x_2)$ as f is one-one, also $f(w) \in s$ as f is L-open. We have $f(w)(f(x_1)) = \sup \{w(x_1)\}$ and $f(w)(f(x_2)) = \sup \{w(x_2)\}$ and $f(w)(f(x_1)) \neq f(w)(f(x_2))$. Since (Y, s) is $L - R_1(i)$, $\exists u, v \in s$ such that $u(f(x_1)) = 1, u(f(x_2)) = 0, v(f(x_1)) = 0, v(f(x_2)) = 1$ and $u \cap v = 0$. This implies that $f^{-1}(u)(x_1) = 1, f^{-1}(u)(x_2) = 0, f^{-1}(v)(x_1) = 0, f^{-1}(v)(x_2) = 1$ and $f^{-1}(u \cap v) = 0 \Rightarrow f^{-1}(u) \cap f^{-1}(v) = 0$.

Now it is clear that $\exists f^{-1}(u), f^{-1}(v) \in \tau$ such that $f^{-1}(u)(x_1) = 1$, $f^{-1}(u)(x_2) = 0$, $f^{-1}(v)(x_1) = 0$, $f^{-1}(v)(x_2) = 1$ and $f^{-1}(u) \cap f^{-1}(v) = 0$.

Hence the L-topological space (X, τ) is $L - R_1(i)$.

Similarly (b), (c), (d), (e), (f), (g) can be proved.

BIBLIOGRAPHY

1. Abu safiya, A.S., Fora, A.A. and Warner, M.W.: Fuzzy separation axioms and fuzzy continuity in fuzzy bitopological spaces; *Fuzzy Sets and Systems*. 62 (1994) 367-373.
2. Ali, D.M.: A note on FT_0 , FT_1 , FT_2 spaces; *Ganit*. (1987) 35-40.
3. Ali, D.M.: Some weaker separation axioms in fuzzy topological spaces; *Math Applicata*. (1988) 1-7.
4. Ali, D.M.: A note on fuzzy regularity concepts; *Fuzzy Sets and Systems*. 35(1990) 101-104.
5. Ali, D.M.: Some remarks on regularity in fuzzy topology; *RU Studies*. 18(1990) 181-190.
6. Ali, D.M.: A note on some FT_2 concepts; *Fuzzy Sets and Systems*. 42(1991) no.3, 381-386.
7. Ali, D.M.: Some other types of fuzzy connectedness; *Fuzzy Sets and Systems*. 46 (1992) 55-61.

8. Ali, D.M.: On the R_1 -property in fuzzy topology; *Fuzzy Sets and Systems*; 50(1992) 97-101.
9. Ali, D.M., Wuyts, P. and Srivastava, A. K.: On the R_0 -property in fuzzy topology; *Fuzzy Sets and Systems*. 38(1990) no.1, 97-113.
10. Caldas, M., Jafari, S. and Noiri, T.: Characterization of pre R_0 and R_1 topological spaces; *Topology proceeding*. 25(2000) 17-30.
11. Chakravarty, K.K., Ahsanullah, T.M.G.: Fuzzy topology on fuzzy sets and tolerance topology; *Fuzzy sets and systems*. 45 (1992) 103-108.
12. Chang, C.L.: Fuzzy topological spaces; *J. Math. Anal Appl.* 24(1968) 182-192.
13. Choubey, A.: Some investigations in Categorical Fuzzy topology; Thesis, *Banaras Hindu University. Varanasi, 1995*.
14. Cutler, D.R. and Reilly, I.L.: A comparison of some Hausdorff notions in fuzzy topological spaces; *Comput. Math. Appl.* 19 (1990) no.11, 97-104.
15. Dorsett, C.: R_0 and R_1 topological spaces; *Math. Vesnik* 2(15) (30) (1978) 112-117.

16. Dude, K.K.: A note on R_0 -topological spaces; *Math.Vesnik* 11(36) (1974) 203-208.
17. Dude, K.K.: A note on R_1 -topological spaces; *Period. Math. Hungar* 13(4) (1982) 267-271.
18. Ekici, E.: On R spaces; *int. j. pure. Appl. Math.* 25(2) (2005) 163-172.
19. Eroglue, M.S.: Topological representation for fuzzy topological spaces; *Fuzzy Sets and Systems.* 42(1991) 335-362.
20. Fora, A.A.: Separation axioms, subspace and product spaces in fuzzy topology; *Arab Culf J. Sci. Res.* 8 (1990) no.3, 1-16.
21. Foster, D.H.: Fuzzy topological groups; *J. Math. Anal. Appl.* 67 (1979) 549-567.
22. Ganguly, S. and Saha, S.: On separation axioms and separations of connected sets in fuzzy topological spaces; *Bull.Cal. Math. Soc.* 79(1987) 215-225.
23. Ghanim, M.H. and Kerre, E. E., Mashhour, A. S.: Separation axioms, subspace and sums in fuzzy topology; *J. Math. Anal. Appl.* 102(1984) 189-202.

24. Goguen, J.A.: L-fuzzy sets; *J. Math. Anal. Appl.* 18(**1967**) 145-174.
25. Hausdorff, F.: *Grundzüge der Mengenlehre*; Leipzig, Germany: von Veit, (**1914**).
26. Hossain, M.S. and Ali, D.M.: On T_1 fuzzy topological spaces; *Ganit. J. Bangladesh Math. Soc.* 24(**2004**) 99-106.
27. Hossain, M.S. and Ali, D.M.: On T_2 fuzzy topological spaces; *J. Bangladesh Academy of Science.* 29(2) (**2005**) 201-208.
28. Hossain, M.S. and Ali, D.M.: On R_0 and R_1 fuzzy topological spaces; *R. U. Studies part-B. j. Sci* 33(**2005**) 51-63.
29. Hossain, M.S. and Ali, D.M.: On Fuzzy Regular topological spaces; *R.U. Studies. Part B. Journal of Science.* Vol.34 (**2006**) 25-36.
30. Hossain, M.S. and Ali, D.M.: On T_0 fuzzy topological spaces; *J. Math and Math. Sci.* Vol. 24(**2009**) 95-102.
31. Hutton, B.: Normality in fuzzy topological spaces; *J. Math. Anal. Appl.* (**1975**) 74-79.
32. Hutton, B.: Products of fuzzy topological spaces Topology; *J. Math. Anal. Appl.* 11(**1980**) 59-67.

33. Hutton, B. and Reilly, I.L.: Separation axioms in fuzzy topological spaces;
Dept. Math. Univ. of Auckland. Report No.55, March 1974.
34. Hutton, B. and Reilly, I.: Separation axioms in fuzzy topological spaces.
Fuzzy Sets and Systems.3 (1980) 93-104.
35. Jin-xuan, F. and Ren Bai-lin: A set of new separation axioms in L-fuzzy
topological spaces; *Fuzzy sets and Systems.96 (1998) 359-366.*
36. Kandil, A. and Sedahmed, E.H.: A fuzzy R_1 separation axioms;
Information Sciences 79(1994) 293-313.
37. Kaufmann, A.: Introduction to the theory of fuzzy subsets; Volume I;
Acamadic Press, I. N. C. (1975).
38. Keskin, A. and Nori, T.: On $\gamma - R_0$ and $\gamma - R_1$ spaces; *Miskole Mathematics
Notes 10(2) (2009) 137-143.*
39. Kerre, E.E. and Ottey, P.L.: On α -generated fuzzy topologies; *Fasc. Math.*
No. 19 (1990) 127-134.
40. Khedr, F.M. and Zeyada, F.M. and Sayed: On separation axioms in
fuzzyfing topology; *Fuzzy Sets and Systems 119(2001) 439-458.*

41. Kudri, S.R.T.: Countability in L-fuzzy topology; *Fuzzy Sets and Systems*. 71 (1995) 241-249.
42. Kudri, S.R.T., Warner, M.W.: L-fuzzy weak local compactness; *Fuzzy sets and Systems*. 81(1996) 275-279.
43. Li, S.G.: Remarks on product of L-fuzzy topological spaces; *Fuzzy Sets and Systems*. 94(1998) 121-124.
44. Li, S.G.: Separation axioms in L-fuzzy topological spaces (I): T₀ and T₁; *Fuzzy Sets and Systems*. 116(2000) 377-383.
45. Lipschutz, S.: General, Topology; Copyright © by the *Schaum publishing Company* 1965.
46. Liu Ying-Ming and Luo Mao-Kang: Fuzzy Topology; Copyright © by World Scientific Publishing Co. Pte. Ltd. 1997.
47. Lowen, R.: Topologies floues; *C. R. Acad Sc. Parish*. 278 series A (1974) 925-928.
48. Lowen, R.: Fuzzy topological spaces and fuzzy compactness; *J. Math. Anal. Appl.* 56(1976) 621-633.

49. Malghan, S.R. and Benchalli, S.S.: On fuzzy topological spaces; *Glasnic Mathematicki*. 16(36) (**1981**) 313-325.
50. Malghan, S.R. and Benchalli, S.S.: On open Maps, closed Maps and Local compactness in fuzzy topological spaces; *J. Math. Anal. Appl.* 99(2) (**1984**) 338-349.
51. Mashhour, A.S., Morsi, Nehad, N.,El. And Tantawy, O.A.: Fuzzy neighborhood syntopogenous structure; *Fuzzy Sets and Systems*. 43(**1991**) no.2, 219-234.
52. Mashhour, A.S. and Morsi, N.N.: On regularity axioms in fuzzy neighborhood spaces; *Fuzzy Sets and Systems*. 44(**1991**) 265-271.
53. Ming, Pu. Pao, Ming, Liu Ying: Fuzzy topology I. Neighborhood Structure of a fuzzy point and Moore-Smith convergence; *J. Math. Anal. Appl.* 76 (**1980**) 571-599.
54. Ming, Pu. Pao, Ming, Liu Ying: Fuzzy topology II. Product and Quotient Spaces; *J. Math. Anal. Appl.* 77 (**1980**) 20-37.

55. Mukherjee, M.N., Sinha, S.P.: On some near fuzzy continuous functions between fuzzy topological spaces; *Fuzzy Sets and Systems*. 34 (1990) no. 2, 245-254.
56. Murdeshwar, A.G. and Naimpally, S.A.: R_1 -topological spaces; *Canad. Math. Bull.* 9(1965) 521-423.
57. Naimpally, S.A.: On R_0 -topological spaces; *Ann. Univ. Sci. Budapest. Eotuos. Sect. Math* 10(1967) 53-54.
58. Nanda, S., Dang, A. Behera: The fuzzy topological complementation Theorem; *Jour. Fuzzy Math.* 1(2) (1993) 303-310.
59. Nouh, A.A.: C-closed sets in L-fuzzy topological space and some of its applications. *Turk J. Math* 26(2002) 245-261.
60. Petricevic, z. : R_0 and R_1 axioms in fuzzy topology; *Mat. Vesnik* 41(1989) 21-28.
61. Petricevic, z. : Some separation axioms in fuzzy spaces; *zb, Rad.* No. 4(1990) 25- 33

62. Roy, B. and Mukherjee, M.N.: A unified theory for R_0 , R_1 and certain other separation properties and their variant form; *Bol. Soc. Paran. Math* 28(2) (2010) 15-24.
63. Rudin, W.: Real and complex analysis; Copyright ©1966, by McGraw-hill Inc.1974.
64. Shanin, N.A.: On separation in topological spaces; *C. R.(Doklady) Acad. Sci. URSS (N. S.)* 38(1943) 110-113.
65. Shen, J.: Separation axiom in fuzzifying topology; *Fuzzy Sets and Systems*. 57 (1993) 111-123.
66. Shostak, A.P.: Two decades of fuzzy topology; the main ideas , concepts and results. (Russian). *Uspekhi. Mat. Nauk* 44(1989) 99-147.
67. Singal, M.K., Prakash, N.: Fuzzy pre-open sets and fuzzy pre separation axioms; *Fuzzy Sets and Systems*. 44(1991) 273-281.
68. Singal, M.K., Rajvanshi, N.: Regularly open sets in fuzzy topological spaces; *Fuzzy Sets and Systems*. 50(1992) 343-353.
69. Sinha, S.P.: Fuzzy normality and some of its weaker forms; *Bull. Korean. Math. Soc.* 28(1991) no.1, 89-97.

70. Sinha, S.P.: Separation axioms in fuzzy topological spaces; *Fuzzy Sets and Systems*. 45(**1992**) 261-270.
71. Song, C.L., Xie, L. and Xia, Z. Q.: L-fuzzifying topology; *J. Appl.Math. & Computing* 15(1) (**2004**) 323-33.
72. Sostak, A.P.: On the neighborhood structure of fuzzy topological spaces; *Zb, Rad.* No. 4(**1990**) 7-14.
73. Srivastava, R., Lal, S.N. and Srivastava, A.K.: Fuzzy Hausdorff topological space; *J. Math. Anal.Appl.* 81(**1981**) 497-506.
74. Srivastava, R. and Srivastava, A.K.: On fuzzy Hausdorff cconcepts; *Fuzzy Sets and Systems* 17(**1985**) 67-71.
75. Srivastava, A.K. and Ali, D.M.: A comparison of some FT2 concepts; *Fuzzy Sets and Systems* 23(**1987**) 289-294.
76. Srivastava, A.K.: R_1 -fuzzy topological spaces; *J.Math.Anal Appl.*127(**1987**) 151-154.
77. Srivastava, M. and Srivastava, R.: On fuzzy pairwise- T_0 and fuzzy Pairwise- T_1 Bitopological spaces; *Indian J. Pure appl. Math.*32 (3) (**2001**) 387-396.

78. Srivastava, R., Lal, S.N. and Srivastava, A.K.: On fuzzy T_0 and R_0 topological spaces; *J. Math. Anal Appl.* 136(1) (1988) 66-73.
79. Srivastava, R., Lal, S.N., Srivastava, A.K.: Fuzzy T_1 -topological spaces; *J. Math. Anal. Appl.* 102 (1984) 442-448.
80. Warner, M.W., Mclean, R.G.: On compact Hausdorff L-fuzzy spaces; *Fuzzy Sets and Systems*. 56 (1993) 103-110.
81. Warren, R.H.: Neighbourhoods, bases and continuity in fuzzy topological spaces; *Rocky Mountain J. Math.* 8 (1978) 459-470.
82. Weiss, M.D.: Fixed points, separation and induced topologies for fuzzy sets; *J. Math. Anal. Appl.* 50 (1975) 142-150.
83. Wong, C.K.: Fuzzy topology: Product and Quotient Theorem; *J. Math. Anal. Appl.* 45 (1974) 512-521.
84. Wong, C.K.: Fuzzy points and local properties of Fuzzy topology; *J. Math. Anal. Appl.* 46 (1974) 316-328.
85. Wuyts, P., Lowen, R.: On separation axioms in fuzzy topological spaces; Fuzzy neighborhood spaces and Fuzzy Uniform spaces; *J. Math. Anal. Appl.* 93 (1973) 27-41.

86. Xu, L.S.: The strong Hausdorff Separation property in L-fuzzy topological spaces; *Mohu. Xitong . Yu. Shuxue*. 5 (1991) no.2, 25-29.
87. Yang, C. T.: On Para compact spaces; *Proc. Amer. Math. Soc.* 5(1954) 185-189.
88. Ying, Ming Sheng: A new approach for fuzzy topology I; *Fuzzy Sets and Systems*. 39 (1991) no. 3, 303-321.
89. Yue, Y.: Lattice-valued induced fuzzy topological spaces; *Fuzzy Sets and Systems*. 158 (2007) 1461-1471.
90. Zadeh, L.A.: Fuzzy sets; *Information and Control* 8 (1965) 338-353.
91. Zhang, Ghong Quan: Fuzzy continuous function and its properties; *Fuzzy Sets and Systems*. 43 (1991) 159-171.
92. Zhang, J., Wang, W., Hu, M.: Topology on the set of R_0 semantics for R_0 algebra; *Soft Comput.* 12 (2008) 585-591.
93. ZHAO Bin: Inclusive Regular Separation in L-fuzzy topological Spaces; *J. of Mathematical Research and Exposition*. 25(4) (2005).