University of Rajshahi	Rajshahi-6205	Bangladesh.
RUCL Institutional Repository		http://rulrepository.ru.ac.bd
Department of Fisheries		PhD Thesis

2014

Development of low cost weed based aquaculture technology in ponds

Asadujjaman, Md.

University of Rajshahi

http://rulrepository.ru.ac.bd/handle/123456789/611 Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

DEVELOPMENT OF LOW COST WEED BASED AQUACULTURE TECHNOLOGY IN PONDS



A THESIS SUBMITTED TO THE UNIVERSITY OF RAJSHAHI, RAJSHAHI, BANGLADESH FOR THE DEGREE OF **DOCTOR OF PHILOSOPHY (Ph. D)**

By Md. Asadujjaman

B. Sc. Fisheries (Hons.) (KU), M. S. in Fisheries Management (BAU)

January, 2014

DEPARTMENT OF FISHERIES FACULTY OF AGRICULTURE UNIVERSITY OF RAJSHAHI RAJSHAHI-6205, BANGLADESH

Dedicated

То Му

Beloved Parents

DECLARATION

I do hereby declare that the research work submitted as a thesis entitled "Development of low cost weed based aquaculture technology in ponds" in the Department of Fisheries, University of Rajshahi, Bangladesh for the Degree of Doctor of Philosophy is the result of my own investigation. The thesis or part of it has not been submitted to any other University or institution for any degree.

Dated: January, 2014

(Md. Asadujjaman)

Ph. D. Fellow Department of Fisheries University of Rajshahi Session: 2009-2010 Reg. No. 0251 Roll No: 09803

CERTIFICATE

Certifying that, the thesis entitled "Development of low cost weed based aquaculture technology in ponds" is a *bona fide* research work of Mr. Md. Asadujjaman, Ph. D. Fellow, Roll no. 09803, Reg. no. 0251 and Session: 2009-2010, Department of Fisheries, University of Rajshahi, Bangladesh.

Principal Supervisor

Co-Supervisor

(Dr. Md. Akhtar Hossain) Professor

(Ex-Chairman) Department of Fisheries University of Rajshahi Rajshahi- 6205 Bangladesh Email: mahfaa@yahoo.com

(Dr. Mohammad Amirul Islam)

Professor Department of Biochemistry and Molecular Biology University of Rajshahi Rajshahi- 6205 Bangladesh Email: maislam06@gmail.com

ACKNOWLEDGEMENTS

The author is glad and grateful to the Almighty Allah who enabled him to pursue higher education in Fisheries science and to complete the research work for the degree of PhD in the Department of Fisheries under the University of Rajshahi, Bangladesh.

The author expresses to his profound sense of gratitude, sincere appreciation and indebtedness to his respected supervisor Dr. Md. Akhtar Hossain, Professor and Ex-Chairman, Department of Fisheries, University of Rajshahi for his scholastic guidance, valuable suggestions, affectionate feelings, constant encouragements, constructive criticisms & over all supervision throughout the study.

The author expresses his immense indebtedness to his honorable Co-supervisor Dr. Mohammad Amirul Islam, Professor, Department of Biochemistry and Molecular Biology, University of Rajshahi for his inspiration, valuable suggestions, constructive criticism and generous help for completion of the research work.

The author expresses his gratitude and hearty thanks to Dr. Md. Delwer Hossain, Associate Professor and Chairman, Department of Fisheries, University of Rajshahi, for his generous help to complete the research work.

The author expresses his deepest sense of gratitude to Dr. Saleha Jasmine, Dr. M. Afzal Hussain, Dr. Fawzia Adib Flowra, Dr. M. Manjurul Alam, Dr. ABM Mohsin, Dr. Md. Ishtiaque Hossain, Dr. Md. Abdur Razzaq Joadder, Dr. Md. Abdus Samad, Dr. Dr. Md. Abu Sayed Jewel, Dr. Md. Mostafizr Rahman Mondol and all other teachers of the Department of Fisheries, University of Rajshahi for their generous help for completion of the research work.

The author gratefully acknowledges the Ministry of Science and Technology, Bangladesh for providing him with the opportunity of NST fellowship for successfully completion of this PhD research work.

The author expresses his deep sense of gratitude to Mr. Md. Anisur Rahman Mridha for his valuable and cordial assistance during the period of study.

The author also expresses special thanks to Mr. Rafik Uddin Ahmed, Computer Operation Supervisor, Department of Fisheries, University of Rajshahi for composing and printing of this thesis.

The author gratefully expresses his sincere appreciation and boundless gratitude to his parents, wife Feroja Pervin and son Masnun Fahham Mafi for their support and sacrifice to complete the research work.

The Author

ABSTRACT

Three experiments were conducted during 2010-2012 towards the development of low cost weed based aquaculture technology in ponds. Experimental ponds were located at Kushtia district, Bangladesh. In experiment-1, six different conventional and non-conventional fish feed items like rice bran, wheat bran, mustard oilcake, Azolla, grass and banana leaves were tested to determine the nutrient contents (protein, lipid and carbohydrate) under 6 treatments as T₁, T_2 , T_3 , T_4 , T_5 and T_6 , respectively. Considering the nutrient content in experiment-1, experiment-2 evaluated the fish production and economics under 4 treatments of feed and weed based system with a similar stocking density of 11115 fish/ha in polyculture ponds as T₀: ponds fed with conventional feed like rice bran, wheat bran and mustard oilcake (no weed was used as fish feed), T1: Azolla (Azolla pinnata) fed ponds, T2: Grass (Cynodon dactylon) fed ponds and T₃: Banana (Musa acuminata) leaf fed ponds. Based on the performance of weeds and feeds in experiment-2, experiment-3 optimized the stocking density for Azolla based carp polyculture system in ponds under 3 different treatments of stocking densities like T₁: 9880 fish/ha, T₂: 11115 fish/ha and T₃: 12350 fish/ha. Nutrients in weeds and feeds were compared for a period of 6 months (April to September) in experiment-1. Fishes (Hypophthalmichthyes molitrix, Catla catla, Labeo rohita, Cirrhinus cirrhosus, Cyprinus carpio, Ctenopharyngodon idella and Barbonymus gonionotus) were also grown for a period of 6 months (April to September) in experiment-2 and experiment-3. Mean initial stocking weight of H. molitrix, C. catla, L. rohita, C. cirrhosus, C. carpio, C. idella and B. gonionotus were 62, 64, 57, 54, 63, 65 and 25 g, respectively for experiment-2 and 60, 65, 58, 52, 61, 70 and 22 g, respectively for experiment-3. There were 3 replications for each treatment under different experiments. Liming (250 kg/ha) and basal fertilization (cowdung: 1500 kg/ha, urea: 60 kg/ha and TSP: 60 kg/ha) were done for all the treatments under experiment-2 and experiment-3. Urea (2.5 kg/ha/day in all treatments except treatment T1 under experiment-2 and in no treatment under experiment-3) and TSP (2.5 kg/ha/day in all treatments under experiment-2 and experiment-3) were applied as periodic fertilization. In case of experiment-1, nutrient contents (protein, lipid and carbohydrate) were monitored monthly whereas in case of experiment-2 and experiment-3, water quality parameters (water temperature, transparency, DO, pH, alkalinity and free CO₂) were monitored fortnightly and fish growth parameters (weight gain and SGR) were monitored monthly. Economics (in terms of total cost, gross benefit, net profit margin and CBR) of fish farming were also evaluated for both experiment-2 and experiment-3. In experiment-1, significant variations (P<0.05) were found in the mean values of nutrient contents with different treatments of feed items but in case of same feed item no significant difference was found in the nutrient content at different months. Among the non-conventional feed items treatment T₄ (Azolla) varied more significantly (P<0.05) for the mean values of protein content. In experiment-2, no significant difference in the mean values of water quality parameters were found among the treatments. Treatment T_0 varied more significantly (P<0.05) for the mean values of SGR, weight gain, final weight, survival rate and total yield but in terms of total cost, gross benefit, net benefit, net profit margin and CBR, treatment T₁ (Azolla fed pond) was found best. In **experiment-3**, no significant difference in the mean values of water quality parameters were found among the treatments. Significant difference (P<0.05) with the treatments were found in all the growth parameters except survival rate. Treatment T_3 varied more significantly (P<0.05) for the mean values of total yield but in terms of total cost, gross benefit, net benefit, net profit margin and CBR, treatment T₁ (stocking density of 9880 fish/ha) was found best. Findings indicated that the stocking density of 9880 fish/ha could be a good option for low cost Azolla based fish farming in Bangladesh.

Title		Page no.
Acknowledgements	3	i
Abstract		ii
Contents		iii-iv
List of Tables		v-ix
List of Figures		х
List of Maps		xi
List of Plates		xii
List of Abbreviation	S	xiii
Glossary		xiv
Chapter One: Gen	eral Introduction	1-15
Chapter Two: Rev	iew of Literature	16-28
2.1: Nutritive	e value of weeds and conventional fish feeds	16
2.2: Water q	uality	18
2.3: Fish production		21
2.4: Econom	nics	26
Chapter Three:	Determination of protein, lipid and carbohydrate contents of conventional and non-conventional feed items used in carp polyculture pond	29-51
3.1: Introduc	tion	29
3.2: Materials and Methods		33
3.3: Results		40
3.4: Discussion		48
5.5: Conclus	sion	51
5.6: Recomr	nendation	51

CONTENTS

Title	Page no.

Chapter Four:	Evaluation of fish production and economics under conventional feed and weed based systems in polyculture ponds	52-106
4.1: Introduct	tion	52
4.2: Materials	s and Methods	54
4.3: Results		63
4.4: Discussi	on	92
5.5: Conclusi	ion	106
5.6: Recomm	nendation	106

Chapter Five:	Optimization of stocking density for <i>Azolla</i> based carp polyculture in pond	107-162
5.1: Introdu	uction	107
5.2: Materi	als and Methods	112
5.3: Result	S	120
5.4: Discus	ssion	149
5.5: Conclu	5.5: Conclusion	
5.6: Recor	nmendation	162
Chapter Six: Su	mmary, Conclusion and Recommendation	163-165
References		166-186
Appendices		187

LIST OF TABLES

Table No.	Title	Page
		No.
Table 1.1:	Top 20 food fish aquaculture producing countries	4
Table 1.2:	Geographical distribution in adoption and coverage of fish culture system by culture pattern in Bangladesh	7
Table 1.3:	Aquatic macrophytes used as fish feed in weed based aquaculture	10
Table 1.4:	Efforts for the promotion of weed based aquaculture	12
Table 3.1:	Dietary protein requirement for highest growth rate of fishes in carp polyculture	30
Table 3.2:	Nutritional status of major conventional feed items used in carp polyculture in Bangladesh	31
Table 3.3:	Nutritive status of non-conventional feed items	32
Table-3.4:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 1 st month (April, 2010)	40
Table-3.5:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 2 nd month (May, 2010)	41
Table-3.6:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 3 rd month (June, 2010)	41
Table-3.7:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 4 th month (July, 2010)	42
Table-3.8:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 5 th month (August, 2010).	42
Table-3.9:	Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 6 th month (September, 2010)	43
Table-3.10	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T ₁ (Rice, <i>Oryza sativa</i> bran)	43
Table-3.11	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T ₂ (Wheat, <i>Trticum aestivum</i> bran)	44
Table-3.12	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T_3 (Mustard, <i>Brassica napus</i> Oilcake)	44
Table-3.13	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T ₄ (<i>Azolla pinnata</i>)	45

Table No.	Title	Page
		No.
Table-3.14	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T_5 (Grass, <i>Cynodon dactylon</i>)	45
Table-3.15	: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T ₆ (Leaf of banana, <i>Musa acuminata</i>)	45
Table-3.16	: Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments during study period	46
Table 4.1:	Stocking density and stocking ratio/combination of the stocked fish species	56
Table-4.2:	Variations in the mean values of water quality parameters under different treatments at 1 st fortnight	64
Table-4.3:	Variations in the mean values of water quality parameters under different treatments at 2 nd fortnight	64
Table-4.4:	Variations in the mean values of water quality parameters under different treatments at 3 rd fortnight	65
Table-4.5:	Variations in the mean values of water quality parameters under different treatments at 4 th fortnight	65
Table-4.6:	Variations in the mean values of water quality parameters under different treatments at 5 th fortnight	66
Table-4.7:	Variations in the mean values of water quality parameters under different treatments at 6 th fortnight	66
Table-4.8:	Variations in the mean values of water quality parameters under different treatments at 7 th fortnight	67
Table-4.9:	Variations in the mean values of water quality parameters under different treatments at 8 th fortnight	67
Table-4.10	: Variations in the mean values of water quality parameters under different treatments at 9 th fortnight	68
Table-4.11	: Variations in the mean values of water quality parameters under different treatments at 10 th fortnight	68
Table-4.12	: Variations in the mean values of water quality parameters under different treatments at 11 th fortnight	69
Table-4.13	: Variations in the mean values of water quality parameters under different treatments at 12 th fortnight	69
Table 4.14	: Variations in the mean values of water quality parameters under different treatments during study period	71
Table-4.15	:Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 1 st month (April)	74
Table-4.16	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 2 nd month (May)	74
Table-4.17	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 3 rd month (June)	75

Table No.	Title	Page
		No.
Table-4.18	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 4 th month (July)	75
Table-4.19	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 5 th month (August)	76
Table-4.20	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 6 th month (September)	76
Table-4.21	: Variations in the mean weight gain (g/month) of fishes under different treatments at 1 st month (April)	78
Table-4.22	: Variations in the mean weight gain (g/month) of fishes under different treatments at 2 nd month (May)	78
Table-4.23	: Variations in the mean weight gain (g/month) of fishes under different treatments at 3 rd month (June)	79
Table-4.24	: Variations in the mean weight gain (g/month) of fishes under different treatments at 4 th month (July)	79
Table-4.25	: Variations in the mean weight gain (g/month) of fishes under different treatments at 5 th month (August)	80
Table-4.26	: Variations in the mean weight gain (g/month) of fishes under different treatments at 6 th month (September)	80
Table-4.27	: Variations in the mean SGR (%, bwd ⁻¹) of fishes under different treatments during the study period	84
Table-4.28	: Variations in the mean weight gain (g/month) of fishes under different treatments during the study period	84
Table-4.29	: Variations in the mean final weight (g) of fishes under different treatments during the study period	85
Table-4.30	: Variations in the mean survival rate (%) of fishes under different treatments during the study period	85
Table-4.31	: Variations in the mean yield (kg/ha) of fishes under different treatments during the study period	86
Table 4.32	Economics of fish farming under different treatments during study period	90
Table 5.1:	Stocking density and stocking ratio/combination of fish species	114
Table-5.2:	Variations in the mean values of water quality parameters under different treatments at 1 st fortnight	121
Table-5.3:	Variations in the mean values of water quality parameters under different treatments at 2 nd fortnight	121
Table-5.4:	Variations in the mean values of water quality parameters under different treatments at 3 rd fortnight	122
Table-5.5:	Variations in the mean values of water quality parameters under different treatments at 4 th fortnight	122

Table No.	Title	Page
		No.
Table-5.6:	Variations in the mean values of water quality parameters under different treatments at 5 th fortnight	123
Table-5.7:	Variations in the mean values of water quality parameters under different treatments at 6 th fortnight	123
Table-5.8:	Variations in the mean values of water quality parameters under different treatments at 7 th fortnight	124
Table-5.9:	Variations in the mean values of water quality parameters under different treatments at 8 th fortnight	124
Table-5.10	: Variations in the mean values of water quality parameters under different treatments at 9 th fortnight	125
Table-5.11	: Variations in the mean values of water quality parameters under different treatments at 10 th fortnight	125
Table-5.12	: Variations in the mean values of water quality parameters under different treatments at 11 th fortnight	126
Table-5.13	: Variations in the mean values of water quality parameters under different treatments at 12 th fortnight	126
Table 5.14	: Variations in the mean values of water quality parameters under different treatments during study period	128
Table-5.15	:Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 1 st month (April)	131
Table-5.16	:Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 2 nd month (May)	131
Table-5.17	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 3 rd month (June)	132
Table-5.18	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 4 th month (July)	132
Table-5.19	:Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 5 th month (August)	133
Table-5.20	: Variations in the mean values of SGR (%, bwd ⁻¹) of fishes under different treatments at 6 th month (September)	133
Table-5.21	: Variations in the mean weight gain (g/month) of fishes under different treatments at 1 st month (April)	135
Table-5.22	: Variations in the mean weight gain (g/month) of fishes under different treatments at 2 nd month (May)	135
Table-5.23	: Variations in the mean weight gain (g/month) of fishes under different treatments at 3 rd month (June)	136
Table-5.24	: Variations in the mean weight gain (g/month) of fishes under different treatments at 4 th month (July)	136
Table-5.25	: Variations in the mean weight gain (g/month) of fishes under different treatments at 5 th month (August)	137

Table No. Title	Page
	NO.
Table-5.26: Variations in the mean weight gain (g/month) of fishes under different treatments at 6 th month (September)	137
Table-5.27: Variations in the mean SGR (%, bwd ⁻¹) of fishes under different treatments during the study period	141
Table-5.28: Variations in the mean weight gain (g/6 month) of fishes under different treatments during the study period	141
Table-5.29: Variations in the mean final weight (g) of fishes under different treatments during the study period	^t 142
Table-5.30: Variations in the mean survival rate (%) of fishes under different treatments during the study period	142
Table-5.31: Variations in the mean yield (kg/ha) of fishes under different treatments during the study period	^t 143
Table 5.32: Economics of fish production under different treatments	147

LIST OF FIGURES

Figure N	lo. Title	Page No.
Fig. 1.1:	Aquatic regions of Bangladesh (Source: Dey et al., 2008)	2
Fig. 1.2:	The relative share of different inland waterbodies in Bangladesh (Source: DoF, 2013)	2
Fig. 1.3:	Percentage composition of fish production in capture, culture and marine fisheries (Source: DoF, 2013)	3
Fig. 1.4:	Fish Production trends from 2001-02 to 2011-12 (Source: DoF, 2013)	5
Fig. 1.5:	Species group based fish production in Bangladesh (Source: DoF, 2012)	6
Fig. 1.6:	Intensification of aquaculture and potential incremental yield	8
Fig. 1.7.	Areas of weed based aquaculture efforts	13
Fig. 3.1.	Variations in the mean values of nutrient contents under different fish feed items (a. Protein; b. Lipid; c. carbohydrate)	47
Fig. 4.1:	Variations in the mean values of water quality parameters (a; Water temperature, b; transparency, c; DO, d; pH, e; alkalinity, f; free CO ₂) under different treatments during study period	72
Fig. 4.2:	Variations in the mean values of different growth parameters under different treatments during the study period: a; SGR (%, bwd ⁻¹), b; weight gain (g/month), c; final weight (g) and d; survival rate (%)	87
Fig. 4.3:	Mean values of individual fish yield under different treatments	88
Fig. 4.5:	Mean values of total fish yield under different treatments	88
Fig. 4.6:	Economics of fish farming under different treatments: a; total cost (BDT), b; net benefit (BDT) and c; CBR	91
Fig. 5.1:	Variations in the mean values of water quality parameters (a, Water temperature; b, transparency; c, DO; d, pH; e, alkalinity; f, free CO_2) under different treatments during study period	129
Fig. 5.2.	Variations in the mean values of different growth parameters under different treatments during the study period: a; SGR (%, bwd ⁻¹), b; weight gain (g), c; final weight (g) and d; survival rate (%)	144
Fig. 5.3:	Variations in the mean values of fish yield under different treatments	145
Fig. 5.4:	Variations in the mean values of total fish yield under different treatments	145
Fig. 5.5:	Economics of fish farming under different treatments: a; total cost (BDT.), b; net benefit (BDT) and c; CBR	148

Map No.	Title	Page No.
Мар 3.1:	Study site (a: site for feed item collection and b: site for feed analysis)	34
Map 4.1:	Shows the study area	54
Map 5.1:	Shows the study area	112

LIST OF MAPS

Plate No.	Title	Page No.
Plate 1.1:	Aquatic macrophytes used as feed in weed based aquaculture	11
Plate 4.1:	Experimental ponds.	60
Plate 4.2:	Monitoring water quality parameters.	61
Plate 4.3:	Measurement of growth of fishes.	62
Plate 4.4:	Harvesting of fishes	89
Plate 5.1:	Experimental ponds	117
Plate 5.2:	Supply of Azolla in experimental pond	118
Plate 5.3:	Monitoring of water quality parameters	118
Plate 5.4:	Measurement of growth of fishes	119
Plate 5.5:	Harvesting of fishes	146

LIST OF PLATES

ABBREVIATIONS

ANOVA	=	Analysis of Variance
BDT	=	Bangladeshi Taka
CBR	=	Cost-Benefit Ratio
CO ₂	=	Carbon-dioxide
RCBD	=	Randomized Completely Block Design
DMRT	=	Duncan Multiple Range Test
DO	=	Dissolved Oxygen
DoF	=	Department of Fisheries
DWRP	=	Duckweed Research Project
GDP	=	Gross domestic production
SGR	=	Specific growth rate
SPSS	=	Statistical Package for Social Science
TSP	=	Triple Super Phosphate

GLOSSARY

Baor	=	Dead arms of river which is confined	
Bana	=	Bamboo fencing	
Beel	=	Natural shallow depression connected to open waters	
Haor	=	Saucer-shaped floodplain depressions	
Lac	=	100 thousand	
PRISM-Bangladesh =		Non-government Organization	
Upazila	=	Sub district	

CHAPTER ONE

GENERAL INTRODUCTION

1.1: Present status of the fisheries sector in Bangladesh

Fisheries sector plays an important role in providing income, employment, nutrition and earning foreign exchange in Bangladesh. It has significant role in the improvement of the socio-economic condition of poor fisherman. The fisheries sector contributes 4.39% of the gross domestic product (GDP), 22.76% of agricultural resources and 2.46% of foreign exchange earning of Bangladesh. Fish alone contributes about 60% of total animal protein intake. More than 10% of total populations of Bangladesh are directly or indirectly dependent on fisheries sector for their livelihoods. It provides full time employment to 14.6 million in fishing and other activities related to fisheries (DoF, 2013).

Fish and Fisheries have been an essential part of the life and culture of the people of Bangladesh. Bangladesh is uniquely rich and diverse in water resources. It has innumerable water bodies including ponds, tanks, lakes, rivers, *haors, baors, beels*, estuaries and inundated paddy fields. Due to favorable climatic condition, the water bodies of Bangladesh are found highly productive and aquaculture is found as an important commercially viable activity (DoF, 2011). The country is blessed with vast inland waterbodies comprising 2710,766 ha of floodplains, 8,53,863 ha of rivers and estuaries, 1,14,161 ha of beels (natural shallow depression connected to open waters), 68,800 ha of man-made reservoir (Kaptai Lake), 3,71,309 ha of ponds and ditches, 5,488 ha of oxbow lakes and 2,75,232 ha of shrimp farms. These water resources offer great scope and potential for augmenting fish production by adopting culture based fisheries techniques (DoF, 2013) (Fig. 1.1 and 1.2).



Fig. 1.1: Aquatic regions of Bangladesh (Source: Dey et al., 2008)



Fig. 1.2: The relative share of different inland waterbodies in Bangladesh (Source: DoF, 2013)

Total fish production in our country during the year 2011-2012 was about 3.26 million metric tons of which 2.68 million metric tons was produced from freshwater including culture fisheries and 0.57 million metric tons from marine water (DoF, 2013) (Fig. 1.3).



Fig. 1.3: Percentage composition of fish production in capture, culture and marine fisheries (Source: DoF, 2013)

There are indications that the production from capture fisheries decreased recently which increased pressure on aquaculture to fill up the gap. Causes for decreasing capture fisheries production include habitat destruction, unplanned construction of flood control barrages, water abstraction for irrigation, over-fishing and reclamation of land for agriculture. Concurrently, aquaculture production increased due to the development and implementation of improved culture techniques and expansion of the pond culture area (Gupta *et al.*, 1999; Alam and Thomson, 2001).

Inspite of having very rich fisheries resources, the present fish production of Bangladesh is unable to meet the demand of fish intake for the increasing population of the country. The need of annual per capita fish intake is 20.44 kg while the amount produced is 18.94 kg (DoF, 2013). This situation clearly indicates the necessity of creating more attention on the aquaculture operation of the country.

1.2 Potentials, problems and challenges of aquaculture in Bangladesh

Global production of fish from aquaculture has grown rapidly over the past four decades, contributing significant quantities to the world's supply of fish for human consumption. Aquaculture now accounts for almost half (45%) of the world's food fish. With its continued growth, it is expected that aguaculture will, in the near future, produce more fish for direct human consumption than capture fisheries. Aquaculture, which started as primarily an Asian freshwater food production system, has now spreaded over all continents, encompassing all aquatic environments and using a range of aquatic species. From an activity that was principally small scale, non-commercial and family based, aquaculture now includes large-scale commercial and industrial production of high-value species that are traded at national, regional and international levels. Although production remains predominantly in Asian countries and is still largely based on small-scale operations, there is a wide consensus that aquaculture has the potential to meet the growing global demand for nutritious food fish and to contribute to the growth of national economies, while supporting the sustainable livelihoods of many communities (FAO, 2006). Globally, Bangladesh reaches 6th position in producing fish (Table 1.1) and yearly total fish production of Bangladesh is also increasing smoothly (Fig. 1.4).

Country	Production (1000 tonnes)	Percentage
China	32 414	67.3
India	2838	5.9
Viet Nam	1437	3.0
Indonesia	1197	2.5
Thailand	1144	2.4
Bangladesh	882	1.8
Japan	746	1.5
Chile	698	1.5
Norway	657	1.4
Philippines	557	1.2
Egypt	540	1.1
Myanmar	475	1.0
United States of America	472	1.0
Republic of Korea	436	0.9
Taiwan, Prov. of China	305	0.6
France	258	0.5
Brazil	258	0.5
Spain	222	0.5
Italy	181	0.4
Malaysia	176	0.4
Rest of the world	225	7 4.7
World total	48150	100.0

Table 1.1: Top	20 food fish	aquaculture	producing	countries
----------------	--------------	-------------	-----------	-----------

Adapted from Subasinghe et al. (2009)





The different fish farming production systems are generally distinguished according to their degree of intensification which is itself usually defined according to the feeding practices as food represents more than 50% of the total operating costs in intensive systems. However, intensification (or, inversely, extensification) involves many other production factors, such as water, land, capital, and labor (Khan *et al.*, 2009).

The main carp species cultivated in the world are primarily seven in number and are often grouped on the basis of their natural geographical occurrence: the so-called Chinese carps, which include the grass carp, *Ctenopharyngodon idella*, the silver carp, *Hypophthalmichthys molitrix,* and the bighead carp, *Aristichthys nobilis,* and the so-called Indian major carps, which include catla, *Catla catla,* rohu, *Labeo rohita,* and mrigal, *Cirrhinus cirrhosus.* The seventh species is the common carp, *Cyprinus carpio.* Taxonomically, carps belong to the family of *Cyprinidae* (order, Cypriniformes) (Azad *et al.*, 2004).

Aquaculture in Bangladesh has rapidly progressed in recent years with a contribution of 44% to the annual fish production (Talukdar *et al.*, 2012). Among

different techniques of aquaculture, polyculture is found as one of the most important techniques. The basic principle of fish polyculture system depends on the idea as compatible species of different feeding habits are cultured together in the same pond, the maximum utilization of all natural food sources takes place without harmful effects. Polyculture or mixed culture of carps has been found as an economically viable and technically sustainable in perennial water bodies (Alikhuni, 1957; Chen, 1976). The selection of fish species is very important for polyculture systems. Generally silver carp (Hypophthalmichthys molitrix), catla (Catla catla), rohu (Labeo rohita), mrigal (Cirrhinus cirrhosus), common carp (Cyprinus carpio), grass carp (Ctenopharyngodon idella) and sharpunti (Barbonymus gonionotus) are selected for polyculture. Carps are the most important species for pond culture. Three major Indian carps rohu (Labeorohita), catla (Catla catla) and mrigal (Cirrhinus cirrhosus) and one exotic carp (silver carp, *Hypophthalmichthys molitrix*) accounted for about 80% of pond fish production. Carp species have a vital role in national fish production in Bangladesh. In 2010-11 fiscal year carp species comprised 34.61% of the total fish production (DoF, 2012) (Fig.1.5).



Fig. 1.5: Species group based fish production in Bangladesh (Source: DoF, 2012).

These species are suitable for low inputs culture system in small ponds and ditches for their quick growth and for maximum production within short period. Freshwater fish-culture systems practiced in Bangladesh can be categorized into four groups: carp polyculture, mixed culture, monoculture and integrated fish culture. Mixed culture of fish is a type of polyculture in which carp and some other species are cultured together. In integrated systems, fish culture is integrated with other agricultural enterprises, such as rice-fish farming, ricefish-duck farming, and fish-poultry farming. Polyculture of Indian and Chinese carps along with a few other exotic species is the most dominant system in Bangladesh. Other practices include pond monoculture of Thai pangus, mixed culture of Nile tilapia and carps in seasonal ponds or ditches, and culture of carps (mainly mirror carp) and silver barb in rice fields. Monoculture of genetically improved Nile tilapia in ponds is also becoming popular, particularly among commercial producers (Dey et al., 2008). The possibilities of increasing fish production per unit area, through carp polyculture are found highest when compared with other systems (Table 1.2).

Major	% of l	Upazilas			
cultural pattern	Extensive	Improved extensive	Semi- intensive	Intensive	adopting system (%)
Carp polyculture	13.7	64.8	41.0	1.1	100.0
Carp polyculture with pangus	11.9	42.5	32.6	2.6	82.2
Pangus monoculture	7.5	22.2	31.9	5.5	63.2
Carp-golda mixed culture	12.6	27.8	14.1	1.1	49.8
Tilapia monoculture	15.9	29.5	26.9	2.2	68.9
Other culture patterns	6.6	15.9	11.9	2.0	33.7

 Table 1.2: Geographical distribution in adoption and coverage of fish culture system by culture pattern in Bangladesh.

Source: Dey et al., 2008.

Different species combination in polyculture system effectively contributes also to improve the pond environment. Algal blooming is common in most tropical manure fed ponds. By stocking phytoplanktophagus Silver carp in appropriate density certain algal blooming can be controlled. Grass carp on the other hand keeps the macrophyte abundance under control due to its macrovegetation feeding habit and it adds increased amount of partially digested excreta which becomes the feed for the bottom dweller coprofagous common carp. The bottom dwelling mrigal, common/mirror carp help re-suspension of bottom nutrients to water while stirring the bottom mud in search of food. Such an exercise of bottom dwellers also aerates the bottom sediment. All these facts suggest that carp polyculture is the most suitable proposition for fish culture in tropical ponds (Dey *et al.*, 2008).

Carp polyculture in Bangladesh may be characterized as semi-intensive, with high fingerling stocking rates and low use of feed and fertilizer (Fig. 1.6). Fish farmers generally use less supplementary feed and other inputs compared with their counterparts in countries with more advanced aquaculture systems, such as China and Thailand (Dey *et al.*, 2008). It is true that feed expenditure is about 60%-80% of total fish production cost. To decrease this production cost it is essential to minimize the cost for fish feed (DoF, 2010). The major challenge for further promotion of carp polyculture is to reduce the fish feed cost which indicates for the development of low cost aquaculture technology using terrestrial and aquatic weeds as fish feed.

Major aquaculture problems in Bangladesh includes the lack of quality seed and low cost feed, poor water quality, disease and parasites (DoF, 2003).



Fig. 1.6: Intensification of aquaculture and potential incremental yield {Source: Dey *et al.* (2008) which is modified from Edwards (1993)}.

1.3: Importance of low cost aquaculture technique in Bangladesh

Fish feed generally constitutes 60-70% of the operational cost in intensive and semi- intensive aquaculture system (Singh et al., 2006). The fish feed used in aquaculture is quite expensive, irregular and short in supply in many third world countries. These feeds are sometimes adulterated, contaminated with pathogen as well as containing harmful chemicals for human health. Naturally there is a need for the development of healthy, hygienic fish feed which influences the production as well as determines the quality of cultured fish (Bag et al., 2011). Considering the importance of nutritionally balanced and costeffective alternative diets for fish, there is a need for research effort to evaluate the nutritive value of different non-conventional feed resources, including terrestrial and aquatic macrophytes (Edwards et al., 1985, Wee and Wang, 1987, Mondal and Ray, 1999). The aquatic weeds have been shown to contain substantial amounts of protein and minerals (Ray and Das, 1995). Aquatic and terrestrial macrophytes have been used as supplementary feeds in fish farming since the early times of freshwater fish culture (Bardach et al., 1972) and still play an important role as fish feed in extensive culture systems (Edwards, 1987). Since the fish production is found not yet sufficient to meet up the existing demand, further promotion of aquaculture is required specially for the poor who are unable to receive the high cost feed based aquaculture technology. Hence weed based system may be a good option as a low cost, environment friendly sustainable aquaculture technique in Bangladesh (Grover et al., 2000).

1.4: Weed based aquaculture system and its history

The weed based system refers to the use of some inputs from plant sources, eg., weeds or grasses or leaves or macrophytes like duckweeds, Azolla etc. (Table 1.3 and Plate 1.1) as supplemental feed in fish production. These inputs are consumed first as feed by herbivorous fish and subsequently a part of the semi digested faecal matter of the macrophytes feeding fishes are consumed by the other fishes and the remaining part will be recycled in food chain as nutrients for primary production, thus they have potentiality to increase the total

fish production of aquaculture system. Aquatic plants have been utilized as food components and thus have played an important role in culture of herbivorous fish since 4000 years ago in Egypt and 2500 years ago in the Orient, including Indian subcontinent (Bardach et al., 1972). Duckweeds might be having as much potential as fish foods that could be utilized in preparation of suitable fish feed essential in expansion of low-cost aquaculture system in the tropics (Hassan and Edwards, 1992). Since long Azolla pinnata utilized as biofertilizer in agriculture has been popular among farmers. Nowadays, its utility in pisciculture has come to limelight and has been proven worthy of note because of its two unique activities: capable of nitrogen fixation from atmosphere that enhances nitrogen in semi-intensive pisciculture systems (Ayyappan et al., 1993) and used as direct food by some macrophagous fish (Cassani, 1981; Antonie et al., 1987). Fresh duckweed (and also the dried meal) is suited to intensive production of herbivorous fish (Gaiger et al., 1984) and duckweed is converted efficiently to live weight gain by carp and tilapia (Van-Dyke and Sutton, 1977; Hepher and Pruginin, 1979; Robinette et al., 1980; Hassan and Edwards, 1992; Skillicorn et al., 1993).

SI. No.	Scientific name	Local name	Characteristics
1	Azolla pinnata	Azolla	The species is typically triangular measuring about 1.5 to 3.0 cm in length, 1 to 2 cm in breadth. Newly form leaves are green but aged leaves are brown in color. With roots.
2	Spirodela polyrriza	Sonapana	Leaves are flat or oval, 6-10.5 mm in length, 5-10 mm wide and 0.6-1.5 mm thick. Deep green above but with deep brown/reddish ventral. It contains 10-15 roots which are 10-40 mm long.
3	Lemna minor	Tetulipana	Leaves are flat and elongated, like tamarind tree leaves, 3-4.5 mm in length, 2-2 mm wide and 0.2-0.3 mm thick. Deep green or green in color. It contains single root which is 10-15 mm long.
4	Wolffia arrhiza	Sujipana, Dimpana	Leaves are minute and rounded, 0.6-1.2 mm in length and 0.5-1 mm wide. Deep green in color and without roots.

Table 1.3: Aquatic macrophytes used as fish feed in weed based aquaculture

Source: Das (1997)







IIRR-ICLARM (1992) reported some farmers' proven weed based integration to aquaculture farming in different countries like China, Malaysia, Vietnam, Philippines and India. The Indian NGO, Sulabh International conducted two duckweed fish production projects in rural areas of the states of Haryana and Orissa (Iqbal, 1999). Since 1989 the NGO, PRISM- Bangladesh has set up an impressive programme to develop and disseminate duck weed aquaculture in Bangladesh. PRISM has developed three centres located at Mirzapur, Manikgonj and Khulna. The centres served as demonstration farms and training institutions for the promotion of integrated duckweed-fish production in the surrounding villages (DWRP, 1996 and DWRP, 1997).

The aforesaid history clearly indicates the necessity of realizing the efforts made for the development of weed based aquaculture, finding out the lacks in our research system and thereby providing strategy of developing low cost aquaculture technique specially for the resource poor farmers.

1.5.: Efforts made in promoting the weed based aquaculture

Some efforts are already taken to promote the weed based aquaculture (Table 1.4 and Fig. 1.7) in home and abroad.

Efforts taken by	Type of study	Place of Study (Country)	Weed used	Fish species used	Major thrust	Remarks
Talukdar e <i>t al.</i> (2012)	Experiment	Bangladesh	Lemna minor	Tilapia, sharpunti, grass carp, catla, mrigal	Use of weed in fish culture	Economics and nutritional aspects are not explored
Kabir <i>et al.</i> (2009)	Experiment	Bangladesh	Lemna minor	Tilapia, catla, rui	Fish production performance	Economics and nutritional aspects are not explored
Chowdhury et al. (2008)	Experiment	Bangladesh	Lemna minor	Nile Tilapia	Use of weed in monoculture	Economics and nutritional aspects are not explored
Ferdoushi et al. (2008)	Experiment	Bangladesh	Lemna and Azolla	Sharpunti, catla, mrigal, rui	Water quality	Economics and nutritional aspects are not explored
Abou <i>et al.</i> (2007)	Experiment	Benin West Africa	Azolla	Nile Tilapia	Fish growth and economics	Economics and nutritional aspects are not explored
Ansal and Dhawan (2007)	Experiment	India	Spirodela sp.	Catla, mrigal, rui	Low-cost carp feed production	Economics and nutritional aspects are not explored
Majhi <i>et al</i> . (2006)	Experiment	India	Azolla	Grass carp	Organic fish production	Economics and nutritional aspects are not explored
El-Shafai <i>et al.</i> (2004)	Experiment	Egypt	<i>Lemna</i> sp.	Nile Tilapia	Use of weed as feed for tilapia culture	Economics and nutritional aspects are not explored
Thy <i>et al.</i> (2004)	Experiment	Combodia	Water spinach, duck weed.	Common carp, mrigal	Fish growth	Economics and nutritional aspects are not explored
Azim and Wahab (2003)	Experiment	Bangladesh	<i>Lemna</i> sp.	Rui, catla, common carp	Fish growth, pond ecology	Economics and nutritional aspects are not explored
lqbal (1995)	Pilot project	Bangladesh	Lemna minor	Tilapia, rui, catla	Fish growth	Economics and nutritional aspects are not explored
Hassan and Edwards (1992)	Experiment	Vietnam	Lemna, Spirodella	Nile Tilapia	Fish growth	Economics and nutritional aspects are not explored
PRISM- Bangladesh (1990)	Pilot project	Bangladesh	<i>Lemna</i> sp	Rui, catla, mrigal, silver carp	Use of weed in fish culture	Only one weed is used
Mibagwv and Adeniji (1988)	Experiment	Nigeria	Lemna sp	Tilapia	Nutritional value of weed	Only one weed is used

Table 1.4: Efforts for the promotion of weed based aquaculture.



Fig. 1.7. Areas of weed based aquaculture efforts. {(Source: Talukdar *et al.* (2012), Abou *et al.* (2007), Ansal and Dhawan (2007), Majhi *et al.* (2006)}

Efforts made indicate that

- (i) Weeds are used as fish feed at home and abroad.
- (ii) Azolla, Lemna, Spirodela and water spinach are used as fish feed for fish farming.
- (iii) Fish species- tilapia and Indian major carps are used in weed based aquaculture system.
- (iv) Major thrusts are fish production, pond ecology and nutritive value of weed.
- (v) Comparative growth performance of fishes using weeds and supplementary feeds (for semi-intensive culture) was not found accordingly in the efforts made in Bangladesh.
- (vi) Economics of the weed based system was not explored well in Bangladesh.
- (vii) Maximum fish species was not introduced in carp-polyculture system under weed based efforts in Bangladesh.
- (viii) Nutritive values of the weeds were also not explored in the efforts made in Bangladesh.

1.6: Research questions

Based on the indications, some questions are also raised:

- (i) Can weed based system be considered as a low cost technique as compared to the conventional feed based system?
- (ii) Can weed based system be considered as suitable model for a typical carp poly culture including maximum species?
- (iii) Which weed performs better than others?
- (iv) Which weed is better in terms of maintaining good water quality?
- (v) Which weed can be used by most of the species directly or indirectly in polyculture system?
- (vi) What is the suitable stocking density of fishes to get higher production for weed based system?

1.7: Research need for weed based aquaculture development in Bangladesh

In recent years, organic aquaculture has been gaining considerable importance. Many farmers have begun shifting from traditional method to organic cultivation as means of producing safe foodstuff and respecting environment. Organic farming favours lower input costs, conserves nonrenewable resources, improves quality of the product and thereby increases farm income. Organic fish farming system rely on biological management in most cases and virtually prohibits utilization of synthetic chemicals in fish production. Earlier efforts are done by using one or two weed species and limited fish species. A technology has little or no value without its economic benefit. In the previous studies, comparative economic analysis of feed and weed system was not found. Nutritive values of the weeds are not also explored well in the study made in Bangladesh.

Based on those arguments, the present study emphasized on determining the nutritional values of weeds and comparing production and economics of fish farming towards the development of low-cost weed based aquaculture technique in Bangladesh.

14

1.8: Objectives

The general objectives of the present study were as follows:

- to determine the nutritional value of conventional and nonconventional feed items used in carp polyculture;
- to evaluate the growth performance of fishes under different weed and feed based systems;
- to monitor the water quality parameters in ponds under different treatments of weed and feed based systems;
- to evaluate the economics of fish production under different treatments of weed and feed based systems; and
- to recommend suitable weed based technology with suitable stocking density for carp polyculture in ponds.

CHAPTER TWO REVIEW OF LITERATURE

Review of related literature is a necessity in the sense that it provides as scope for reviewing the stock of knowledge and information relevant to the proposed research. However, the following literatures were reviewed in favour of the present study.

2.1: Nutritive value of weeds and conventional fish feeds

Pullin and Almazan (1983) estimated the nutritive value of Azolla and reported the crude protein, crude lipid, crude fiber, ash and starch as 24-30%, 3-33%, 9.1%, 10.5% and 6.5%, respectively.

Van-Hove and Lopez (1983) reported that Azolla contained 28% crude protein.

Tacon and Jackson (1985) reported that aquatic fern (dry basis) could be used as shrimp and fish feed, as it revealed a good nutritive value in terms of protein 25.3%, crude fat 3.8%, crude fiber 9.3% and ash 12.5%.

Mibagwv and Adeniji (1988) worked with *Lemna* sp. in the Kainji lake, Nigeria and reported crude protein ranging from 26.3-45.5% of dry weight.

Hepher (1988) reported the protein in ricebran, wheat bran, oil cake and Azolla as 11.88%, 14.57%, 30-33% and 19.27%, respectively.

Banerjee and Matai (1990) worked on the nutritive status of *Azolla pinnata* and reported ash as 15.4%, protein as 21.9% and Lipid as 3.8%.

Gavina (1994) reported the percentage composition (dry weight) of Azolla as moisture 5.06%, ash 22.21%, crude protein 20.98%, crude fat 5.17%, crude fiber 19.30%, calcium 1.08% and phosphorous 0.21%.

Suresh and Mandal (2000) worked on the determination of nutritive value of rice bran, mustard oilcake and Azolla. In rice bran they found crude protein and crude fibre as 12.6% and 21.9%, respectively. In mustard oilcake crude protein and crude fibre as 38.6% and 6.8%, respectively and in Azolla crude protein and crude fibre as 26.5% and 20.4%, respectively.

Fasakin and Balogun (2001) made a study on nutritional analysis of *Azolla* sp. and found crude protein as 28.9%, crude fibre as 12.2% and lipid contents as 4.6%. They also noted that nutritive values of *Azolla* sp. were higher than the values of *Spirodela* sp.

El-Shafai *et al*. (2004) reported the protein content of duckweed as 21.1% and wheat bran as 11.0%.

Alalade and Iyayi (2006) conducted an experiment on the chemical composition of Azolla (*Azolla pinnata*) and they reported crude protein, ether extract, crude fibre and ash as 21.4%, 2.7%, 12.7% and 16.2%, respectively.

Abou *et al.* (2007) studied the effects of stocking density on growth and profitability of farming of Nile tilapia (*Oreochromis niloticus*) in Azolla fed ponds and found crude protein, crude fat and crude fibre as 28.51%, 3.97% and 9.60%, respectively in Azolla meal.

Ansal and Dhawan (2007) worked on *Spirodela* sp. for low-cost carp feed formulation and they found crude protein, fat and ash as 30.52%, 1.97% and 9.45%, respectively.

Tavares *et al.* (2008) observed 38.8% crude protein, 3.8% crude fat and 13.2% crude fiber in dried duck weed. They also reported that protein content of duckweeds growing on nutrient poor and nutrient rich water varied between 15-25% and 35-45% (Dry matter basis), respectively.

Ahmed (2011) worked on the use of Azolla as fish feed and found protein, ash, carbohydrate and fat as 20-25%, 10%, 6-6.5% and 3-3.5%, respectively.

Bag *et al.* (2011) worked with aquatic weed as potential feed for Nile tilapia (*Oreochromis niloticus*) and reported as dry matter, organic matter, crude protein, crude lipid, ash and crude fibre as 92.56%, 82.80%, 20.56%, 9.89%, 9.76% and 9.24%, respectively.

17
2.2: Water quality

Wurts and Durbrow (1992) worked on interactions of pH, carbon dioxide, alkalinity and hardness in fish ponds and they reported the desirable range of pH as 7.5-8.5, CO₂ as 1-2 ppm, alkalinity as 50-150 mg/l and hardness as 75-200 mg/l for pond aquaculture.

Azim *et al.* (1995) recorded mean values of temperature as 26.2°C, transparency as 36.2 cm and pH as 7.1 in fertilized fish pond.

Wahab *et al.* (1995) recorded water temperature from 27.2 to 32.4°C, secchi depth between 26 to 50 cm, pH around 6.0, total hardness from 45 to 108 mg/l and dissolved oxygen from 2.2 to 7.1 mg/l in carp polyculture ponds.

DWRP (1996 and 1997) made a study on weed-based carp polyculture at Tangail and found water temperature, transparency, pH and DO as 28.1 to 32°C, 30 to 35cm, 6.5 to 7.3 and 4 to 6 mg/l, respectively.

Rahman (1999) studied the effect of duckweed on pond ecology and recorded the values of water temperature from 26.2 to 34.5°C, transparency from 12.0 to 46.5 cm, total alkalinity from 71.0 to 175.0 mg/l, pH from 6.5 to 8.8 and dissolved oxygen from 6.0 to 8.0 mg/l.

Dhawan and Kaur (2002a) studied the water quality in polyculture ponds under different treatments of feed utilization (T₁: 0% rice bran + 100% mustard oilcake, T₂: 25% rice bran + 75% mustard oilcake , T₃: 50% rice bran + 50% mustard oilcake, T₄: 75% rice bran + 25% mustard oilcake, T₅: 100% rice bran + 0% mustard oilcake, T₆: 18 L/ha/yr pig dung and T₇: 36 L/ha/yr pig dung) at Punjab, Ludhiana, India. They reported pH in T₁, T₂, T₃, T₄, T₅, T₆ and T₇ as 8.40, 8.36, 8.40, 8.47, 8.56, 8.56 and 8.45, respectively in winter; 8.93, 8.92, 8.94, 8.89, 8.83, 8.93 and 8.55, respectively in summer, DO (mg/l) in T₁, T₂, T₃, T₄, T₅, T₆ and T₇ as 11.41, 11.62, 12.00, 12.11, 12.19, 12.75 and 12.31, respectively in winter; 7.85, 7.60, 7.44, 7.56, 7.57, 7.91 and 7.61, respectively in summer, total alkalinity (mg/l) in T₁, T₂, T₃, T₄, T₅, T₆ and T₇ as 228, 236, 209, 233, 245, 237 and 219, respectively in winter; 310.25, 287.75, 285.50, 319.00, 280.25, 290.50 and 264.75, respectively in summer.

Azim and Wahab (2003) worked on the development of duckweed-fed carp polyculture system in Bangladesh and they stated that the water temperature, transparency, pH, dissolved oxygen and alkalinity ranged from 26°C to 31°C, 29.8 to 35.9 cm, 6.9 to 7.5, 4.1 to 5.25 mg/l and 110.25 to 145.70 mg/l, respectively.

El-Shafai *et al.* (2004) observed the water quality parameters in three different types of treated ponds (sewage-wheat bran fed pond, sewage-duck weed fed pond and freshwater duckweed fed pond). In case of sewage-wheat bran fed pond, values for temperature (°C), pH and DO (mg/l) as 23-35, 8-9.7 and 4.9±4.7, respectively. In case of sewage-duck weed fed pond temperature (°C), pH and DO (mg/l) values were 23-35, 8-9.4 and 5.9±4.7, respectively. In case of freshwater duckweed fed pond temperature (°C), pH and DO (mg/l) values were 23-35, 8-9.4 and 5.9±4.7, respectively. In case of freshwater duckweed fed pond temperature (°C), pH and DO (mg/l) values were 23-35, 8-9.4 and 5.9±4.7, respectively. In case of freshwater duckweed fed pond temperature (°C), pH and DO (mg/l) values were 23-35, 8.4-9.9 and 5.3±4.1, respectively.

Thy *et al.* (2004) reported the mean values of pH, water transparency (cm) and dissolved oxygen (mg/l) as 8.48, 30.7 and 4.15, respectively in effluent plus duckweed fed pond.

Majhi *et al.* (2006) reported that water temperature, pH, dissolved oxygen and total alkalinity varied from 20-22°C, 7.5-8.2, 7.0-7.8 mg/l and 40 to 45 ppm, respectively in Azolla fed pond.

Rahman *et al.* (2006) observed the water quality parameters in polyculture ponds under 3 treatments of fish stocking densities i.e. T_1 (9,880 nos/ha), T_2 (14,820 nos/ha) and T_3 (19,760 nos/ha). In case of T_1 , they reported the mean values of water temperature (°C), transparency (cm), and DO (mg/l) as 29.43, 26.37 and 6.40, respectively. In case of T_2 , the mean values of water temperature (°C), transparency (cm), and DO (mg/l) were as 29.36, 23.82 and 6.37, respectively. In case of T_3 , the mean values of water temperature (°C), transparency (cm), and DO (mg/l) were as 29.36, 23.82 and 6.37, respectively. In case of T_3 , the mean values of water temperature (°C), transparency (cm), and DO (mg/l) were as 29.28, 20.71 and 6.29, respectively.

Abou *et al.* (2007) observed the water quality parameters in three different types of treated ponds (30% Azolla fed pond, 35% Azolla fed pond and 40% Azolla fed pond). In case of 30% Azolla fed pond, transparency (cm),

temperature (°C), pH, DO (mg/l), nitrates (mg/l), nitrites (mg/l), and ammonium (mg/l) values were 42.2 ± 8.8 , 28.8 ± 0.2 , 6.61 ± 0.11 , 4.48 ± 0.24 , 0.12 ± 0.01 , 0.01 ± 0.00 and 0.46 ± 0.07 , respectively. In case of 35% Azolla fed pond, transparency (cm), temperature (°C), pH, DO (mg/l), nitrates (mg/l), nitrites (mg/l), and ammonium (mg/l) values were 40.4 ± 7.2 , 28.9 ± 0.1 , 6.40 ± 0.29 , 4.66 ± 0.26 , 0.10 ± 0.02 , 0.01 ± 0.00 and 0.42 ± 0.01 , respectively. In case of 40% Azolla fed pond, transparency (cm), temperature (°C), pH, DO (mg/l), nitrates (mg/l), nitrates (mg/l), nitrites (mg/l), nitrates (mg/l), nitr

Chowdhury *et al.* (2008) conducted an experiment on monoculture of Nile tilapia with duckweed (*Lemna minor*) as supplementary feed for a period of 90 days under 2 treatments viz. T_1 (Duckweed fed pond) and T_2 (without duckweed). In case of treatment T_1 , they stated the water temperature (°C), transparency (cm), DO (mg/l), pH, free CO₂ (mg/l) and alkalinity (mg/l) were 28.30, 28.66, 7.64, 7.53 and 59.75, respectively. In case of treatment T_2 , they stated the water temperature (°C), transparency (cm), DO (mg/l) were 28.59, 34.50, 7.29, 7.58 and 64.58, respectively.

Ferdoushi *et al.* (2008) studied the effects of two aquatic floating macrophytes (Lemna and Azolla) as biofilters of nitrogen and phosphate in fish ponds under three treatments T₁ (Lemna fed pond), T₂ (Azolla fed pond) and T₃ (without any macrophytes). In case of Lemna fed pond, temperature (°C), transparency (cm), dissolved oxygen (mg/l), pH, alkalinity (mg/l), NO₃-N (mg/l) and PO₄-P (mg/l) were found as 29.60±2.97, 34.69±11.93, 4.80±1.93, 7.53b±0.034, 133.87±18.95, 1.29±0.39 and 1.04±0.75, respectively. In case of Azolla fed pond, temperature (°C), transparency (cm), dissolved oxygen (mg/l) and PO₄-P (mg/l) were found as 29.30±2.63, 36.32±19.16, 4.84±1.63, 7.61±0.39, 144.93±25.09, 1.33±0.40 and 1.04±0.76, respectively. In case of control pond, temperature (°C), transparency (cm), dissolved oxygen (mg/l), pH, alkalinity (mg/l), NO₃-N (mg/l) and PO₄-P (mg/l) were found as 29.57±2.59, 26.27±13.47, 5.27±1.86, 8.09±0.46, 224.41±47.36, 1.87±0.63 and 1.64±1.02, respectively.

Bag *et al.* (2011) reported the mean values of water temperature, pH, free CO₂ and DO as 30°C, 7.2, 0.5 mg/l and 6 mg/l, respectively in ponds under aquatic weed based tilapia farming in India.

Talukdar *et al.* (2012) worked on the suitability of duckweed (*Lemna minor*) as feed for fish in polyculture system under two treatments i.e. T_1 (*Lemna minor* fed pond) and T_2 (without *Lemna minor*). In case of treatment T_1 , they reported water temperature (°C), transparency (cm), DO (mg/I), pH, free CO₂ (mg/I) and alkalinity (mg/I) were 28.02, 30.50, 6.63, 7.45, 2.85 and 75.33, respectively. In case of treatment T_2 , water temperature (°C), transparency (cm), DO (mg/I), pH, free CO₂ (mg/I), pH, free CO₂ (mg/I) and alkalinity (mg/I) were 27.92, 32.08, 6.23, 7.43, 2.90 and 74.00, respectively.

2.3: Fish production

Gaiger *et al.* (1984) studied the production of tilapia in three different types of ponds (only duckweed fed pond, only pellete fed pond and duckweed + pellete fed pond). They reported a good feed conversion ratio (1:1) and poor growth rate (0.67% of body mass d⁻¹) in ponds fed with only duckweed. When the fish were fed pellets in addition to duckweed the rate of duckweed consumption decreased and growth rate of the fish doubled with feed conversion ratios between 1.2 and 1.8. Fish grown on the mixed diet performed similarly to fish grown on pellets but had a better feed conversion ratio.

Edwards (1990) conducted an experiment on the use of terrestrial vegetation and aquatic macrophytes in aquaculture and reported tilapia yield of 3.7 t/ha/year from the fertilization of the water with excreta and 13.4 t/ha/year with the addition of Azolla as a nutritional supplement.

Shanmugasundaram and Balusamy (1993) reported the use of Azolla as feed to raise Indian major carps (catla: rohu: mrigal = 1:1:1) in low-lying wetlands with a stocking density of 3000/ha and thereby found the fish yield as 154 kg/ha.

Skillicorn *et al.* (1993) stated that the average yield of fish per hectare was estimated at around 10 tons annually using only duckweed as the supplement to the naturally available fish feed.

Gavina (1994) worked on pig-duck-fish-Azolla integration under 3 treatments i.e., T_1 (commercial feed alone), T_2 (fresh Azolla+60% commercial feed), T_3 (fresh Azolla+40% commercial feed) with a same stocking density (20,000 no/ha) for all the treatments and reported the fish production after four months of culture period, were 581, 436 and 374 kg/ha in treatment T_1 , T_2 and T_3 , respectively

Ahmed *et al.* (1995) studied the impact of improved aquaculture technologies in ponds fed with duckweed. They reported the carp production as 11.04 kg/40 m^2 , silver barb production as 6.75 kg/40 m^2 and Nile tilapia production as 13.43 kg/40 m^2 with a stocking density of 25-30, 60-65 and 80-85/40 m^2 for carp, Silver barb and Nile tilapia respectively.

Gopakumar *et al.* (1999) reported the fish production as 3-4 tonnes/ha/year in ponds under weed based carp polyculture in India.

Grover *et al.* (2000) recommended the Azolla based carp polyculture in ponds with a stocking density of 9880 nos/ha (catla : silver carp : grass carp : rajpunti : mrigal : kalbaus : common carp = 4:6:3:8:10:3:3:3) and thereby made assumption of fish production as 5575 kg/ha.

Cagauan *et al.* (2000) studied the growth of Nile tilapia (initial stocking size of 10-20 g) stocked at a density of 10,000 fishes/ha in integrated fish and Azolla farming pond in Philippines and observed the fish production as 3381.75 kg/ha/yr.

Abdelghany and Ahmad (2002) studied the growth and production of fish (Nile tilapia, common carp and silver carp) in Azolla fed pond for a period of 19 weeks under five different treatments. Feeding dose were 0.0% (T₁), 0.5% (T₂), 1.0% (T₃), 3.0% (T₄), 5.0% (T₅) fish biomass per day and apparent satiation (T₆). Each pond was stocked with 1000 Nile tilapia, 1000 silver carp and 200 common carp with average initial weights of 13.7g, 1.9g and 10.9g,

respectively. Fish yield (kg/ha) was 2770, 3226, 3571, 4598, 4225 and 4637 with treatment T_1 , T_2 , T_3 , T_4 , T_5 and T_6 , respectively,

Dhawan and Kaur (2002b) evaluated the fish growth in pig dung treated polyculture pond during winter and summer and reported that the growth of Indian major carps viz *Catla catla*, *Labeo rohita* and *Cirrhina mrigala* was higher during summer and that of exotic carps viz *Cyprinus carpio* and *Ctenophayrengodon idella* was higher during winter.

Jena *et al.* (2002a) conducted another experiment to evaluate the fish production performance under 3 treatments of polyculture stocked with six carp species, viz., catla, rohu, mrigal, silver carp, grass carp and common carp in the ratio of 2:2:2:2:0.5:1.5, at a combined density of 10,000 fingerlings/ha with average stocking weight of 17.5 g. The treatments were T₁ (single cropping-the fishes were harvested after rearing period of one year), T₂ (single stocking and multiple harvesting- the fishes of larger size were harvested at monthly intervals from the seventh month onwards) and T₃ (multiple cropping system-the fishes were harvested again after due pond preparation for second crop of 5.5 months). The net production recorded in treatments T₁, T₂ and T₃ were 5843.8 kg/ha/year, 6320.3 kg/ha/year and 6828.4 kg/ha/year, respectively.

Jena *et al.* (2002b) evaluated the production performance of carps under 4 treatments i.e. T_1 (5000 fingerlings/ha with six carp species: catla, rohu, mrigal, silver cap, grass carp and common carp, in the ratio of 2 : 2.5 : 2.0 : 1.5 : 0.5 : 1.5), T_2 (5000 fingerlings/ha with three Indian major carp species: catla, rohu, mrigal, in the ratio of 3 : 4 : 3), T_3 (10000 fingerlings/ha with six carp species: catla, rohu, mrigal, rohu, mrigal, silver cap, grass carp and common carp, in the ratio of 2 : 2.5 : 2.0 : 1.5 : 0.5 : 1.5) and T_4 (10000 fingerlings/ha with three Indian major carp species: catla, rohu, mrigal, in the ratio of 3 : 4 : 3). The mean initial stocking sizes of catla, rohu, mrigal, silver carp, grass carp and common carp were 15 g, 5 g, 4 g, 9 g, 33 g and 8 g, respectively. They reported mean net production levels in T_1 , T_2 , T_3 and T_4 as 1791 kg/ha, 2385 kg/ha, 2532 kg/ha and 3472 kg/ha, respectively in 6 months.

Azim and Wahab (2003) studied the effects of duckweed (*Lemma* sp) application on fish production. They reported that a stocking density of Thai silver barb at the rate of 6000 fingerlings ha⁻¹ with 3300 rohu, 3300 catla and 3400 common carp fingerlings ha⁻¹ was found appropriate species mix for duckweed fed poly culture system and the net fish production was 2020 kg ha⁻¹ during a 4-month culture period.

El-Shafai *et al.* (2004) recommended that fresh duck weed is superior to wheat bran as supplementary feed in tilapia pond farming. In their experiment the stocking density was 50 fish/dec (initial stocking weight of 20g) per pond under continuous aeration and regular removal of feaces for a period of 150 days and feeding rate was 25 g fresh Azolla/100g fish. They reported that duckweed fed ponds provided higher net fish yield (11.8 ton/ha/y) than that of wheat bran fed ponds (8.9 ton/ha/y).

Thy *et al.* (2004) worked on the effect of water spinach and duckweed on fish growth in poly culture ponds in Cambodia under 3 treatments i.e. T_1 (Effluent), T_2 (Effluent + water spinach) and T_3 (Effluent + duckweed) stocked with three fish species at a density of 3 fishes per m² with the species tilapia (*Oreochromis niloticus*), common carp and mrigal (*Cirrhinus mrigala*) with a composition of 40%, 35% and 25%, respectively. The water spinach and duckweed were supplied daily at 3 to 5% (Dry matter basis) of fish bodyweight. They reported the survival rate, daily weight gain and total yield as 88.3%, 6.17 kg/ha/day and 1,450 kg/ha/4 months for common carp, 90.6%, 15.2 kg/ha/day and 2,470 kg/ha/4 months for mrigal and 92.4%, 21.2 kg/ha/day and 3,120 kg/ha/4 months for tilapia, respectively.

Rahman *et al.* (2006) conducted an experiment on the culture potential of Thai sharpunti with major carps in seasonal ponds for a period of 90 days under 3 treatments i.e. T_1 (stocking density of rohu, catla, mrigal, silver carp and Thai sharpunti fingerlings was 494, 494, 1,482, 1,235 and 9,880/ha, respectively), T_2 (stocking density of rohu, catla, mrigal, silver carp and Thai sharpunti fingerlings was 494, 494, 1,482, 1,235 and 14,820/ha, respectively) and T_3 (stocking density of rohu, catla, mrigal, silver carp and Thai sharpunti fingerlings was 494, 494, 1,482, 1,235 and 14,820/ha, respectively) and T_3 (stocking density of rohu, catla, mrigal, silver carp and Thai sharpunti fingerlings was 494, 494, 1,482, 1,235 and 19,760/ha, respectively). All the

fishes were fed with supplementary diet comprising of rice bran and mustard oil cake (1:1) at the rate of 6-7% of the estimated body weight twice daily. They found the total fish production as 1,248.34, 1,343.19 and 1,592.67 kg/ha in treatment T_1 , T_2 and T_3 , respectively.

Majhi *et al.* (2006) worked on the effect of Azolla feeding on growth performance of grass carp under 2 treatments i.e T_1 (Azolla fed pond) and T_2 (Control) with the stocking density of 0.75 fish/m² in each pond for a period of 150 days. The fish were fed everyday at 9.30 am at the rate of 10% of fish biomass. They found growth of grass carp fed with T_1 (Azolla fed pond) was remarkably high (1.65 g/fish/day) and the total production of fish was 1850.76 kg/ha.

Abou *et al.* (2007) conducted an experiment to observe the effect of stocking density on growth, yield and profitability of farming Nile tilapia fed with Azolla diet in earthen ponds under 3 treatments i.e. T_1 (30% Azolla), T_2 (35% Azolla) and T_3 (40% Azolla) for a period of 90 days with a stocking density of 1 fish/m² for all the treatments. They reported the fish production (kg/acre/y) as 70.75±1.48, 67.95±0.71 and 65.01±2.21 in T_1 , T_2 and T_3 , respectively.

Chowdhury *et al.* (2008) conducted an experiment on monoculture of Nile tilapia with duckweed (*Lemna minor*) as supplementary feed for a period of 90 days under 2 treatments viz. T_1 (Duckweed fed pond) and T_2 (Control) stocked with 80 fingerlings/dec. They reported the mean survival rate (%), SGR (%, bwd⁻¹), net production (kg/ha/yr) as 94.37 and 93.75, 1.16 and 0.80, 4021.16 and 2203.24 in treatment T_1 and T_2 , respectively.

Ferdoushi *et al.* (2008) conducted an experiment on the effects of two aquatic floating macrophytes (Lemna and Azolla) as biofilters of nitrogen and phosphate in fish ponds for a period of four months under 3 treatments i.e. T_1 (Lemna fed pond), T_2 (Azolla fed pond) and T_3 (Control) with the similar stocking densities in all treatments. The stocking densities were three native carps: *Labeo rohita*, *Cirrhina mrigala*, *Catla catla* and one exotic carp, Thai sharpunti (*Barbonymus gonionotus*) were 2,000, 2,000, 2,000 and 5,000/ha, respectively. Lemna and Azolla were periodically supplied at the rate of 4% of

body weight of Thai sharpunti (*B. gonionotus*) and made available for 24 hours, per day. Commonly available ingredients such as rice bran and mustard oil cake were also used as supplementary feed daily at the rate 4% of body weight of total fish in all the treatments. They reported that these two macrophytes increase fish production.

Bag *et al.* (2011) worked on making Azolla as potential feed for Nile tilapia under 4 treatments i.e. T_1 (Lemna meal), T_2 (Water hyacinth meal), T_3 (Azolla meal) and T_4 (Control) for a period of 90 days with the stocking density of 260 Nile tilapia/dec (average weight of 5.5 g and length of 4.5 cm). They reported the final weight as 80.66±0.73 g and final length as 12.9±0.14 cm and feed intake as 2.01±0.20 g Azolla/fish/day.

Talukdar *et al.* (2012) worked on the suitability of duckweed (*Lemna minor*) as feed for fish in polyculture system under two treatments i.e. T_1 (*Lemna minor* fed pond) and T_2 (without *Lemna minor*) with the stocking density of 40 fishes/40 m² (tilapia : sharpunti : grass carp : catla : mirgal = 45:38:15:38:15). They reported the survival rate (%) of tilapia, sharpunti, grass carp, catla and mrigal as 82, 88, 96, 82, 95 with treatment T_1 and 82, 88, 86, 90, 96 with treatment T_2 . They mentioned the initial weight (g) of tilapia, sharpunti, grass carp, catla and mrigal as 32.67, 30.33, 74.33, 28.47, 18.83 with treatment T_1 and 32.67, 30.33, 74.33, 28.47, 18.83 with treatment T_2 and the end of the experiment they reported the final weight (g) of tilapia, sharpunti, grass carp, catla and mrigal as 68.92, 65.67, 231.03, 26.82, 84.69 with treatment T_1 and 53.11, 56.72, 115.38, 28.40, 74.00 with treatment T_2 .

2.4: Economics

Shanmugasundaram and Balusamy (1993) worked on rice-fish-Azolla integration in India. They stocked 3 Indian major carps (catla : rohu : mrigal = 1 : 1 : 1) at a stocking density of 3000 nos/ha. Azolla was applied twice a day at 2 tonnes/ha. Banana pseudostem, cowdung and rice bran were supplied as supplementary feed at 5% bwd⁻¹. They stated that gross return ((US\$/ha) was 985, net return (US\$/ha) was 463 and benefit cost ratio was 1.88.

Ahmed *et al.* (1995) made a comparative study on economics of carp, silver barb and tilapia faming and reported that average net income of carp farmers was almost triple (2.92 times) than that of silver barb farmers, and more than double (2.12 times) than that of Nile tilapia farmers. They also reported that cost of production in silver barb culture was 34% higher than carp polyculture, and 32% higher than Nile tilapia culture.

Abdelghany and Ahmad (2002) worked on the effects of feeding rates on growth and production of Nile tilapia, common carp and silver carp reared in polyculture system under 6 treatments i.e. 0.0% (T₁), 0.5% (T₂), 1.0% (T₃), 3.0% (T₄), 5.0% (T₅) and apparent satiation (T₆). Total cost (US\$/ha) was 628.07, 1064.67, 1461.51, 2759, 5180.51 and 2520.31 in treatments T₁, T₂, T₃, T₄, T₅ and T₆, respectively. Income from fish sale (US\$/ha) was 3968.26, 4738.54, 5157.93, 6986.33, 6365.42, 7064.37 in treatments T₁, T₂, T₃, T₄, T₅ and T₆, respectively. Net profit (US\$/ha) was 3340.19, 3673.87, 3696.42, 4226.85, 1184.91 and 4544.06 in treatments T₁, T₂, T₃, T₄, T₅ and T₆, respectively.

Roy *et al.* (2003) studied economics of carp-SIS polyculture and their economic analysis under 3 treatments i.e. T_1 (only carps), T_2 (carps with mola) and T_3 (carps with chela) in which the stocking density of Indian major carps, rohu (*Labeo rohita*), catla (*Catla catla*) and mirgal (*Cirrhinus mrigala*) was 9,500 fish/ha with a ratio of 1:1:1, respectively, grass carp (*Ctenopharyngodon idella*) 500 fish/ha, mola (*Amblypharyngodon mola*) and chela (*Chela cachius*) 25,000 fish/ha. They reported the total operational cost as 32,450.00, 39,950.00 and 42,450.00 Tk./ha/7 months in treatments T_1 , T_2 and T_3 , respectively. The total revenue was 128,000.00, 128,280.00 and 110,720.00 Tk./ha/7 months in treatments T_1 , T_2 , and T_3 , respectively. Benefit-cost ratio (BCR) was obtained as 3.94, 3.21 and 2.61 in treatments T_1 , T_2 and T_3 , respectively. BCR was higher in only carp polyculture system because the operational cost was comparatively lower due to the absence of small fish.

Majhi *et al.* (2006) studied the economic evaluation of grass carp production with Azolla feeding and they reported the net profit of \$118.00 ($(0.12/m^2)$) was obtained from 1000 m² pond with an investment of \$314.

Abou *et al.* (2007) conducted an experiment on cost, returns and profitability of feeding Nile tilapia under 3 treatments i.e. T_1 (30% Azolla), T_2 (35% Azolla) and T_3 (40% Azolla) for a period of 90 days. They reported investment cost (US\$) as 37.5, 36.8 and 36.6, gross return (US\$) as 85.7, 82.8 and 79.0, net return (US\$) as 48.2, 46.0 and 43.4 and profitability (%) as 1.3, 1.3 and 1.2 with treatments T_1 , T_2 and T_3 , respectively.

Ebrahim *et al.* (2007) worked on the economical analysis of the different levels of Azolla meal on Nile tilapia fingerlings under 5 treatments i.e. T_1 (0% Azolla meal), T_2 (10.6% Azolla meal), T_3 (21.2% Azolla meal), T_4 (31.8% Azolla meal) and T_5 (42.4% Azolla meal). They found the return IE (Egyptian pound) as 2.24, 1.90, 2.43, 2.17, 1.30 in treatments T_1 , T_2 , T_3 , T_4 and T_5 , respectively. They mentioned that the treatment T_3 got highest benefit among the treatments.

Khan *et al.* (2009) conducted an experiment of the economics of pangasiid catfish (*Pangasius hypophthalamus*) and silver carp (*Hypophthalmichthys molitrix*) polyculture under 3 treatments of feeding frequencies i.e. T_1 (feeding once a day), T_2 (feeding two times a day) and T_3 (feeding three times a day) with the stocking density of 25,000 fishes/ha. They reported that the treatment T_3 having generated the maximum benefit as compared to others.

CHAPTER THREE

Determination of protein, lipid and carbohydrate contents of conventional and non-conventional feed items used in carp polyculture pond

3.1: Introduction

The technique of polyculture of fish is based on the concept of utilization of different trophic and spatial niches of a pond in order to obtain maximum fish production per unit area. Different compatible species of fish of different trophic and spatial niches are raised together in the same pond to utilize all sorts of natural food available in the pond (Rahman *et al.*, 1992). Since the idea of polyculture is based on the principle that each species stocked has its own feeding niche that does not completely overlap with the feeding niches of other species therefore, a more comprehensive use is made of the food resources and space available in polyculture than in monoculture. In South Asia, especially in Bangladesh, several culture combinations of indigenous and exotic carp species are commonly practiced (Wahab *et al.*, 1994; Miah *et al.*, 1997). In some cases, one species enhances the food availability for other species and thus increases the total fish yield per unit area (Hepher *et al.*, 1989; Miah *et al.*, 1993; Azad *et al.*, 2004).

Growth and reproduction of fish and other aquatic animals are primarily dependent upon an adequate supply of nutrient, both in terms of quantity and quality, irrespective of the culture system in which they are grown. Supply of inputs (feeds, fertilizers etc.) has to be ensured so that the nutrients and energy requirements of the species under cultivation are met and the production goals of the system are achieved. Complete data on nutrient requirements are only available for a limited number of species. Although, dietary proteins. lipids and carbohydrates utilization is relatively well investigated for several fishes, available data on nutrient requirements for carp species are presented in table 3.1.

Table 3.1:Dietary protein requirement for highest growth rate of fishes in carp polyculture

Carp species	Crude dietary protein (%)
Common carp (<i>Cyprinus carpio</i>)	31-40.6
Rohu (<i>Labeo rohita</i>)	34-36
Grass carp (Ctenopharyngodon idella)	23-28
Silver carp (Hypophthalmichthys molitrix)	22-29
Catla (<i>Catla catla</i>)	24-27
Mrigal (Cirrhinus cirrhosus)	22-28
Thai punti (<i>Barbonymus gonionotus</i>)	20-25

Source: Hepher (1990), Tacon (1990), De Silva and Anderson (1995) and Hasan *et al*. (1996).

Supplementary feed plays an important role in achieving higher fish production. Unfortunately lack of low cost supplementary feed is found as one of the major problems in aquaculture in Bangladesh (DoF, 2011). Conventional supplementary feed for carps comprises rice bran, wheat bran and oil cakes (Table 3.2). Although rice bran and mustard oil cake are being used as supplementary fish feed in Bangladesh, these materials are costly. Fish meal and other animal byproducts, as source of protein, are not available for use in fish feeds. Commercial fish feeds are not easily available and unaffordable to poor fish farmers in Bangladesh. Consequently, there is no regular organized supplementary feeding practice and the fish production is found as low as 0.5-1.5 t/ha/year (Suresh and Mandal, 2000). It was thus considered necessary to look for cheaper and locally available materials as substitutes.

The optimal protein requirements of carp are affected by the nutritional value of the dietary protein and level of non-protein energy in the carp diet. When sufficient energy sources such as lipids and carbohydrates are available in the diet, most of the ingested protein goes to protein synthesis. Adult Indian major carps require 30% dietary protein for proper growth and survival. Lipids or fats are required as sources of energy and essential fatty acids, and serve as carriers for fat-soluble vitamins. The gross lipid requirement of Indian major carp is 7-8% of the diet, and young fish require relatively more fat and protein than adults. Carbohydrate is the least-expensive nutrient and also a less expensive energy source for carp. Indian major carp, being herbivorous/ omnivorous feeders, easily digest appreciable quantities of carbohydrates in their diets. A dietary level up to 30% carbohydrate does not affect the growth of carp and growth retardation and reduced feed efficiency are observed, however, when carbohydrate levels exceeded 35% of diet (Murthy, 2003).

Feed items	Crude protein (%)	Crude lipid (%)	Crude fibre (%)
Rice bran	12.6	16.5	16.3
Wheat bran	18.2	4.4	14.0
Mustard oilcake	36.5	11.2	11.6
Coconut oilcake	18.2	10.2	11.7
Fishmeal	56.4	19.7	2.2
Blood meal	92.9	0.3	0.5
Bone meal	17.5	5.2	3.5
Soybean meal	45.2	20.5	5.0

Table 3.2: Nutritional status of major conventional feed items used in carp polyculture in Bangladesh

Source: Hasan *et al.* (1989); Hasan *et al.* (1994); Zaher and Mazid (1994); Hossain (1996)

Fish culture is induced primarily by the need for increased protein supply. One of the most essential prerequisites for the successful management of fish culture programme is a comprehensive understanding of feeding (Halver, 1972). The increase in cost and demand of feed protein from conventional sources necessitates fish culturists of the developing countries to incorporate cheap and locally available ingredients in fish feeds. Recently the utilization of aquatic plants having high food value are used to supplement fish food has taken a new dimension for producing the much required animal protein at low cost (Lakshmanan *et al.*, 1967).

Aquatic macrophytes have been known to have potential food value (Edwards, 1980). A perusal of the available literature shows that some of the aquatic weeds are highly nutritive and, therefore, one alternative solution to check the massive population of these weeds might be their utilization through incorporation as components of feedstuff for fish. In fact, significant effort has been directed towards

evaluating the nutritive value (Table 3.3) of different non-conventional feed resources, including terrestrial and aquatic macrophytes, to formulate nutritionally balanced and cost-effective diets for fish and poultry (Edwards *et al.*, 1985; Patra and Ray, 1988; Ray and Das, 1995; Wee and Wang, 1987). Most of these nutritional studies are carried out abroad and no comprehensive studies are found in comparing the nutritional quality of both conventional and non-conventional feeds for fish farming in Bangladesh. However, before advocating the utilization of these aquatic weeds for supplementation of fish feeds, there is an urgent need to explore their nutritional quality, throughout the major culture season in ponds under carp polyculture system. Therefore, the present study aimed at evaluating the protein, lipid and carbohydrate content in conventional and non-conventional feed items used for carp polyculture system in Bangladesh.

Species	Protein (%)	Lipid (%)	Carbohydrate (%)
Azolla pinnata	21.9	3.8	-
Chara sp.	17.5	1.63	23.0
Eichhornia crassipes	16.5	3.4	28.1
Ipomoea aquatica	20.6	4.6	-
Lemna minor	20.4	3.8	-
Nymphae spp.	16.4	2.6	-
Trapa natans	10.8	5.1	-
Wolffia arrhiza	21.5	5.50	10.6
Banana leaves	6.0	-	-
Cassava leaves	2.0	-	-
Sweet potato leaves	2.0	-	-

Table 3.3: Nutritive status of non-conventional feed items	Table 3.3: Nutritive	status of non-o	conventional f	feed items.
--	----------------------	-----------------	----------------	-------------

Source: Boyd (1968, 1969), Culley and Epps (1973), Banerjee and Matai (1990) and Boeck (1996)

3.2: Specific objectives

The specific objectives included in this study were as follows-

- to compare the protein, lipid and carbohydrate contents in feed items (rice bran, wheat bran, oilcake, Azolla, grass and banana leaves); and
- to recommend suitable strategy in selecting feed item for the development of weed based fish farming in carp polyculture pond.

3.3: Materials and methods

3.3.1: Duration and location of the study

The study was conducted for a period of six months from April 2010 to September 2010. Feed items were collected from the fish farming study site located at Alampur village under Kushtia district of Bangladesh (Map 3.1). Whereas nutrient analysis was done at the Protein and Enzyme Research Laboratory under the Department of Bio-Chemistry and Molecular Biology, Rajshahi University, Rajshahi, Bangladesh.



Map 3.1: Study site (a: site for feed item collection and b: site for feed analysis)

3.3.2: Experimental design

The current experiment was carried out under six treatments of feed items each with three replications. The treatment assignments were designated as T_1 , T_2 , T_3 , T_4 , T_5 and T_6 for rice bran, wheat bran, mustard oilcake, Azolla, grass and banana leaves, respectively.

Conventional feed items (rice bran, wheat bran, mustard oilcake) were collected from local market during the experimental period. Non-conventional feed item like Azolla was collected from Azolla ponds adjacent to the research area whereas grass and banana leaf were collected from adjacent grass field and banana garden. Both conventional and non-conventional feed items were collected once a month for nutritional analysis during the experimental period.

3.3.3: Nutrient analysis of the collected samples

Determination of total protein content

Total protein contents of samples were determined by the micro-kjeldahl method (Rangama, 1979).

Apparatus required

Kjeldahl digestion flask: (250 ml capacity), Distillation apparatus, 100 ml conical flask, 50 ml beaker, 50 ml burette and 100 ml volumetric flask.

Reagents required

- a) Mixed indicator: Prepared 0.1% bromocresol green and 0.1% methyl red indicators in 95% alcohol separately. 10 ml of the bromocresol green was mixed with 2 ml of the methyl red solution in a bottle provided with a dropper, which delivered 0.05 ml per 4 drops.
- b) 2% Boric acid: 10 g of boric acid (crystals) was dissolved in 500 ml of boiling distilled water. After cooling, the solution was transferred into a glass-stoppered bottle.

- c) 30% Sodium hydroxide solution: 150 g of sodium hydroxide pellets was dissolved in 375 ml of distilled water. The solution was stored in a bottle closed with rubber stopper.
- d) Catalysts for digestion: 2.5 g of powdered selenium dioxide (SeO₂), 100 g of potassium sulphate (K₂SO₄) and 20 g of copper sulphate (CuSO₄, 5H₂O) were mixed.
- e) 0.01 N Hydrochloric acid: The concentration of the solution was checked against pure sodium carbonate.

Procedure

Digestion

The samples were weighed accurately and transferred to a 250 ml Kjeldahl flask. 1 g of catalyst mixture and 25 ml of conc. H₂SO₄ were added to it. The flask was placed in an inclined position on the stand in the digestion chamber. The flask was heated gently over a low flame until the initial frothing was ceased and the mixture was boiled briskly at a moderate rate. During heating the flask was rotated several times. The heating was continued until the color of the digest was pale blue. The digest was cooled and 30 ml of water was added to 5 ml portion with mixing. The digest was transferred to a 100 ml volumetric flask. The flask was then rinsed 3 times with water and the washings were transferred to the volumetric flask. The solution was made up to volume with water. A blank digestion was carried out without the sample and the digest was made up to 100 ml in total.

Distillation and titration

The distillation apparatus was placed under the condenser. The distilled water was boiled in the steam generator using a Bunsen burner. Stop crock and pink clamp were closed. Cold water was run through the condenser, from which 5 ml of distillate was collected per minute. The burner was removed, where upon the condensate in the distilling flask was sucked back into the steam trap. Funnel was filled with distilled water, and the stopcock was opened momentarily to drain the water into flask.

The burner was then replaced under the steam generator for about 20 seconds and it was removed again. 20 ml of 2% boric acid was pipetted into a clean conical flask and the mixed indicator was added to it. The micro burette was filled with 0.01 N HCI to the zero mark by this time; the distilling flask had become empty. The burner was replaced under the steam generator and pinch clamp was opened to remove liquid from the steam trap. The pinch clamp was left on the glass tubing through which the steam was escaped. The beaker was replaced under the condenser with the conical flask containing boric acid, and the flask was supported in an oblique position, so that the tip of the condenser was completely immersed into the liquid. The stopcock was opened with one hand and with the other hand 10 ml of the digest was pipetted. The funnel was rinsed twice with 3 ml portions of distilled water. Then necessary amount of 30% NaOH was introduced and stopcock was closed. The pinck cock was replaced on the digestion mixture and sodium hydroxide, and the ammonia was liberated which escaped with steam through the condenser into the boric acid solution.

The boric acid was changed from bluish purple to bluish green as soon as it came in contact with ammonia. The change, which was very sharp, was taken place between 20 to 30 seconds after the pinch clamp was closed. Boric acid had changed color within 5 minutes; the conical flask was lowered sot that the condenser tip was 1 cm above the liquid. The end of the condenser was washed with a little distilled water. Distillation was continued until sufficient distillate was collected. The burner was then removed. The distillate was titrated with standard hydrochloric acid until the blue color was disappeared. The titrated was done in daylight. The blank distillation and titration were carried out as in the case of the sample. The percentage of protein in sample was calculated using the following formula:

Percentage of nitrogen = $\frac{(V_{A}-V_{B})\times N\times 14\times V_{M}\times 100}{A\times W\times 100}$

Where,

- W = Weight of the sample taken
- V_A = Volume of HCI in actual titration

- V_B = Volume of HCI in blank titration
- N = Normality of HCI
- V_M = Volume made up of the digest
- A = Aliquot of the digest taken

Thus, % Protein = % Nitrogen × 6.25

Determination of lipid content

Lipid content of the samples were determined by Bligh and Dyer (1989) method.

Reagent used

A mixture of chloroform and ethanol (2:1V/V).

Procedure

5 g of sample were first grinded in a morter and pasted with 10 ml of distilled water. The grinded sample was transferred to a separating funnel and 30 ml of chloroform-ethanol mixture was added. The mixture was mixed well. It was then kept overnight at room temperature in the dark. At the end of this period 20 ml of chloroform and 20 ml of water were further added and mixed. Generally three layers were seen. A clear lower layer of chloroform containing the entire lipid, a colored aqueous layer of ethanol with all water soluble materials and a thick pasty inter phase were seen.

The chloroform layer was carefully colleted in a pre-weighed beaker (50 ml) and then placed on a steam bath for evaporation. After evaporation of the chloroform, the weight of the beaker was determined again. The difference in weight gives the amount of the lipid.

Calculation

Percent of lipid content (g per 100 g of sample) = $\frac{\text{Amount of lipid obtained}}{\text{Weight of sample}} \times 100$

Determination of carbohydrate content

The starch content of samples were determined by the Anthrone method (Boel *et al.* 1988)

Reagents used

Anthrone regent (0.2% in concentrated H_2SO_4), standard starch solution (10 mg / 100 ml distilled water) and 1(M) HCl

Procedure

5 g of samples were homogenized well with 10 ml of water. It was then filtered through double layer of muslin cloth. To filtrate twice the volume of ethanol was added to precipitate the polysaccharide, mainly starch. After kept it over night in cold the precipitate was collected by centrifugation in a clinical centrifuge at 3000 r.p.m. for 15 minutes. The precipitate was then dried over a steam bath. Then 40 ml of 1 (M) hydrochloric acid was added to the dried precipitate and heated to 70°C. It was then transferred to a volumetric flask and diluted to 100 ml with 1 (M) HCI. Then 2 ml of diluted solution was taken in another 100 ml volumetric flask and diluted to 100 ml with 1 (M) HCI.

Aliquot of 1 ml of the extract of each part was pipetted into different test tubes and 4 ml of anthrone reagent was added to the solution of each tube and mixed well. Glass marbles were placed on top of each tube to prevent loss of water by evaporation. The tubes were placed in a boiling water bath for 10 minutes, then removed and cooled. A reagent black was prepared by taking 1 ml of water 4 ml of anthrone reagent in test tube at 680 nm in a colorimeter. The amount of starch present in the sample was calculated by the following calculation.

Calculation

The percentage of starch content (g per 100 g of samples) = $\frac{\text{Amount of starch}}{\text{Weight of sample}} \times 100$

3.3.4: Statistical analysis

Data on nutrient contents (protein, lipid, carbohydrate) under different treatments were subjected to one way ANOVA (Analysis of Variance) using computer software SPSS (Statistical Package for Social Science, version-11). The mean values were also compared to see the significant difference from the DMRT (Duncan Multiple Range Test) after Gomez and Gomez (1984).

3.4: Results

Monthly variations

Monthly variations in the mean values of nutrient contents (protein, lipid and carbohydrate) under different treatments of feed items are presented in table 3.4 to 3.9.

Protein content significantly varied from $6.05\pm0.45\%$ with T₆ (banana leaf) at 6th month (September, 2010) to $31.20\pm0.32\%$ with treatment T₃ (mustard oilcake) at 2nd month (May, 2010). Lipid content significantly varied from 2.95±0.21% with treatment T₆ (banana leaf) at 5th month (August, 2010) to $13.72\pm0.36\%$ with treatment T₃ (mustard oilcake) at 4th month (July, 2010). Carbohydrate significantly varied from $32.85\pm0.14\%$ with treatment T₃ (mustard oilcake) at 4th month (July, 2010) to $66.35\pm0.32\%$ with T₂ (wheat bran) at 3rd month (June, 2010). In the same feed item no significant difference in the nutrient content was found during the study period (Table 3.10 to 3.15).

Table-3.4: Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 1st month (April, 2010).

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T2T3(Wheat, Trticum(Mustard, Brassica aestivum bran)Dilcake)		T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.60±0.22 ^d	17.20±0.05°	30.65±0.18ª	18.65±0.08 ^b	7.28±0.35°	6.25±0.11 ^f				
Lipid (%)	10.42±0.31⁵	6.75±0.41°	13.34±0.31ª	3.25±0.09 ^d	6.35±0.05°	3.05±0.04 ^d				
Carbohydrate (%)	44.25±0.41°	66.20±0.36ª	32.86±0.18 ^f	50.36±0.75 [♭]	46.58±0.12 ^d	48.85±0.36°				

Table-3.5:	Variations	in	the	mean	values	of	nutrient	(protein,	lipid	and
	carbohydra 2010).	ite)	conte	ents un	der diffe	rent	treatmen	ts at 2 nd m	onth (May,

			Treat	ments		
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T ₂ (Wheat, <i>Trticum</i> aes <i>tivum</i> bran)	T₃ (Mustard, <i>Brassica</i> <i>napus</i> Oilcake)	T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)
Protein (%)	13.92±0.19 ^d	17.05±0.12℃	31.20±0.32ª	18.45±0.41 ^b	7.32±0.25 ^e	6.20±0.21 ^f
Lipid (%)	_ipid (%) 10.50±0.25 ^b 6.66±0.69 ^c 13.24±0.47 ^a 3.		3.15±0.12 ^d	6.28±0.06 ^c	3.12±0.11 ^d	
Carbohydrate (%)	43.72±0.19 ^e	65.75±0.32ª	32.90±0.25 ^f	50.45±0.61 ^b	46.30±0.41 ^d	47.98±0.26 ^c

Table-3.6: Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 3rd month (June, 2010).

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T2T3(Wheat, Trticum(Mustard Brassica aestivum bran)Dilcake)		T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.65±0.19 ^d	17.25±0.12°	30.50±0.32ª	18.35±0.41 ^b	7.45±0.25 ^e	6.32±0.21 ^f				
Lipid (%)	Lipid (%) 10.64±0.25 ^b 6.80±0.69 ^c 13.2		13.25±0.47ª	3.12±0.12 ^d	6.45±0.06 ^c	3.10±0.11 ^d				
Carbohydrate (%)	43.85±0.19°	66.35±0.32ª	33.10±0.25 ^f	50.20±0.61 ^b	45.95±0.41 ^d	48.10±0.26°				

Table-3.7:	Variations	in	the	mean	values	of	nutrient	(protein,	lipid	and
	carbohydra	ite)	conte	ents un	der diffe	rent	treatmen	ts at 4 th m	onth (July,
	2010).									

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T2T3(Wheat, Trticum(Mustard, Brassica aestivum bran)Dilcake)		T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.50±0.36d	16.95±0.24°	30.25±0.15ª	18.45±0.32 ^b	7.15±0.14 ^e	6.12±0.31 ^f				
Lipid (%)	10.20±0.21 ^b	7.12±0.46°	13.72±0.36ª	3.35±0.18 ^d	6.23±0.12°	3.20±0.17 ^d				
Carbohydrate (%)	44.20±0.24 ^e	66.32±0.26ª	32.85±0.14 ^f	50.15±0.54 ^b	46.85±0.38 ^d	48.30±0.31°				

Table-3.8: Variations in the mean values of nutrient (protein, lipid and carbohydrate) contents under different treatments at 5th month (August, 2010).

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T2T3(Wheat, Trticum(Mustard, Brassica aestivumaestivumnapus Dilcake)		T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.22±0.28 ^d	17.10±0.34°	30.15±0.11ª	18.75±0.24 ^b	7.25±0.19°	6.14±0.36 ^f				
Lipid (%)	10.24±0.15 ^b	6.47±0.32 ^c	13.22±0.18ª	3.14±0.34 ^d	6.21±0.18 ^c	2.95±0.21 ^d				
Carbohydrate (%)	44.32±0.20 ^e	66.12±0.15ª	32.98±0.31 ^f	50.20±0.17⁵	46.70±0.19 ^d	48.90±0.35°				

Table-3.9:	Variations in	the me	an valu	es of	nutrient	(protein,	lip	id and
	carbohydrate)	content	s under	differe	ent treatr	nents at	6 th	month
	(September, 2	010).						

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T ₂ (Wheat, <i>Trticum</i> aes <i>tivum</i> bran)	T₃ (Mustard, <i>Brassica</i> <i>napus</i> Oilcake)	T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₅ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.50±0.24 ^d	17.22±0.18°	30.45±0.17ª	18.80±0.26 ^b	7.12±0.23 ^e	6.05±0.45 ^f				
Lipid (%)	10.45±0.26 ^b	6.32±0.38°	13.20±0.19ª	3.10±0.41 ^d	6.32±0.28°	2.96±0.41 ^d				
Carbohydrate (%)	44.20±0.16 ^e	65.99±0.23ª	33.02±0.46 ^f	49.88±0.27 ^b	45.76±0.14 ^d	48.85±0.24°				

Table-3.10:Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T₁ (Rice, *Oryza sativa* bran).

	Months									
Nutrients	April	Мау	June	June July		September				
Protein (%)	14.60±0.22ª	13.92±0.19ª	14.65±0.19ª	14.50±0.36ª	14.22±0.28ª	14.50±0.24ª				
Lipid (%)	10.42±0.31ª	10.50±0.25ª	10.64±0.25ª	10.20±0.21ª	10.24±0.15ª	10.45±0.26ª				
Carbohydrate (%)	44.25±0.41ª	43.72±0.19ª	43.85±0.19 ^a	44.20±0.24ª	44.32±0.20ª	44.20±0.16ª				

Table-3.11: Monthly	variations	in	nutrient	(protein,	lipid	and	carbohydrate)
contents	with treatm	ent	T ₂ (Wheat	, Trticum a	estivun	n bran).

	Months									
Nutrients	April	April May		July	August	September				
Protein (%)	17.20±0.05ª	17.05±0.12ª	17.25±0.12ª	16.95±0.24ª	17.10±0.34ª	17.22±0.18ª				
Lipid (%)	6.75±0.41ª	6.66±0.69ª	6.80±0.69ª	7.12±0.46ª	6.47±0.32ª	6.32±0.38ª				
Carbohydrate (%)	66.20±0.36ª	65.75±0.32ª	66.35±0.32ª	66.32±0.26ª	66.12±0.15ª	65.99±0.23ª				

Table-3.12: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T_3 (Mustard, *Brassica napus* Oilcake).

	Months									
Nutrients	April	April May		July	August	September				
Protein (%)	30.65±0.18ª	31.20±0.32ª	30.50±0.32ª	30.25±0.15ª	30.15±0.11ª	30.45±0.17ª				
Lipid (%)	13.34±0.31ª	13.24±0.47ª	13.25±0.47ª	13.72±0.36ª	13.22±0.18ª	13.20±0.19ª				
Carbohydrate (%)	32.86±0.18ª	32.90±0.25ª	33.10±0.25ª	32.85±0.14ª	32.98±0.31ª	33.02±0.46ª				

	Months									
Nutrients	April	Мау	June	July	August	September				
Protein (%)	18.65±0.08ª	18.45±0.41ª	18.35±0.41ª	18.45±0.32ª	18.75±0.24ª	18.80±0.26ª				
Lipid (%)	3.25±0.09ª	3.15±0.12ª	3.12±0.12ª	3.35±0.18ª	3.14±0.34ª	3.10±0.41ª				
Carbohydrate (%)	50.36±0.75ª	50.45±0.61ª	50.20±0.61ª	50.15±0.54ª	50.20±0.17ª	49.88±0.27ª				

Table-3.13: Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T_4 (*Azolla pinnata*).

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-3.14:Monthly variations in nutrient (protein, lipid and carbohydrate) contents with treatment T₅ (Grass, *Cynodon dactylon*).

	Months									
Nutrients	April May		June	July	August	September				
Protein (%)	7.28±0.35ª	7.32±0.25ª	7.45±0.25ª	7.15±0.14ª	7.25±0.19ª	7.12±0.23ª				
Lipid (%)	6.35±0.05ª	6.28±0.06ª	6.45±0.06ª	6.23±0.12ª	6.21±0.18ª	6.32±0.28ª				
Carbohydrate (%)	46.58±0.12 ^a	46.30±0.41ª	45.95±0.41ª	46.85±0.38ª	46.70±0.19ª	45.76±0.14ª				

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-3.15: Monthly	variations	in	nutrient	(protein,	lipid	and	carbohydrate)
contents	with treatm	ent	T ₆ (Leaf o	f banana, <i>N</i>	lusa ad	cumina	ata).

	Months									
Nutrients	April	Мау	June	July	August	September				
Protein (%)	6.25±0.11ª	6.20±0.21ª	6.32±0.21ª	6.12±0.31ª	6.14±0.36ª	6.05±0.45ª				
Lipid (%)	3.05±0.04ª	3.12±0.11ª	3.10±0.11ª	3.20±0.17ª	2.95±0.21ª	2.96±0.41ª				
Carbohydrate (%)	48.85±0.36ª	47.98±0.26ª	48.10±0.26ª	48.30±0.31ª	48.90±0.35ª	48.85±0.24ª				

Mean variations

The variations in the mean values of nutrient contents (protein, lipid and carbohydrate) with different treatments of feed items are presented in table 3.16 and fig. 3.1.

Protein content significantly varied from $6.18\pm0.13\%$ with treatment T₆ (banana leaf) to $30.53\pm0.40\%$ with treatment T₃ (mustard oilcake). Lipid content significantly varied from $3.06\pm0.09\%$ with treatment T₆ (banana leaf) to $13.33\pm0.10\%$ with treatment T₃ (mustard oilcake). Carbohydrate significantly varied from $32.95\pm0.29\%$ with treatment T₃ (mustard oilcake) to $66.12\pm0.47\%$ with treatment T₂ (wheat bran).

Table-3.16:Variations in	the	mean	value	es of	nut	trient	(prote	ein, lipi	d and
carbohydrate)	cont	tents	under	differ	ent	treatr	nents	during	study
period.									

	Treatments									
Nutrients	T₁ (Rice, <i>Oryza</i> sativa bran)	T₂ (Wheat, <i>Trticum</i> aestivum bran)	T₃ (Mustard, <i>Brassica</i> <i>napus</i> Oilcake)	T₄ (Azolla pinnata)	T₅ (Grass, Cynodon dactylon)	T₀ (Leaf of banana, <i>Musa</i> acuminata)				
Protein (%)	14.40±0.32 ^d	17.13±0.07°	30.53±0.40ª	18.58±0.09 ^b	7.26±0.18 ^e	6.18±0.13 ^f				
Lipid (%)	10.41±0.31 ^b	6.69±0.30°	13.33±0.10ª	3.19±0.10 ^d	6.31±0.13°	3.06±0.09 ^d				
Carbohydrate (%)	44.09±0.67 ^e	66.12±0.47ª	32.95±0.29 ^f	50.21±0.54 ^ь	46.36±0.16 ^d	48.50±0.51°				







Fig. 3.1. Variations in the mean values of nutrient contents under different fish feed items (a. Protein; b. Lipid; c. carbohydrate).

3.5: Discussion

Monthly variations of the nutrient contents

Protein content varied from $6.05\pm0.45\%$ with (T₆ at 6th month) to $31.20\pm0.32\%$ (T₃ at 2nd month). Lipid content ranged from $2.95\pm0.21\%$ (T₆ at 5th month) to $13.72\pm0.36\%$ (T₃ at 4th month). Carbohydrate content ranged from $32.85\pm0.14\%$ (T₃ at 4th month) to $66.35\pm0.32\%$ (T₂ at 3rd month). Suresh and Mandal (2000) worked on the determination of nutritive value of rice bran, mustard oil cake and Azolla for a period of 4 months from July to October. In rice bran they found crude protein and crude fibre as 12.6% and 21.9%, respectively. In mustard oilcake, crude protein and crude fibre was 38.6% and 6.8%, respectively and in Azolla, crude protein and crude fibre dwas 26.5% and 20.4%, respectively. Sithara and Kamalaveni (2008) worked on the formulation of low cost fish feed using Azolla as a protein supplement during September to March and reported 20-25.5% protein in Azolla. Ebrahim *et al.* (2007) used Azolla as tilapia diet for a period of 90 days in summer season and reported 20% protein in Azolla. Fasakin and Balogan (2001) worked on the nutritional aspects of Azolla in August, 1997 and reported 20.9% protein in Azolla.

Present findings also indicated that in case of same feed item, no significant difference was found in the nutrient content at different months (Table 3.10 to 3.15). This might be due to no major change in the temperature was found to affect the growth and composition of Azolla during the study period (Appendix-1). This statement was almost agreed with Lumpkin and Plucknett (1982) who reported that change in Azolla composition was subjected to change in environment. Statement also agreed with Van-Hove *et al.* (1987) and Ebrahim *et al.* (2007) who reported that change in Azolla composition was subjected to change in species.

Mean variation of the nutrient contents

In the present study the protein content varied from $6.18\pm0.13\%$ (T₆, banana leaf) to $30.53\pm0.40\%$ (T₃, mustard oilcake), lipid content varied from $3.06\pm0.09\%$ (T₆, banana leaf) to $13.33\pm0.10\%$ (T₃, mustard oilcake) and carbohydrate content varied from $32.95\pm0.29\%$ (T₃, mustard oilcake) to $66.12\pm0.47\%$ (T₂, wheat bran). The highest protein and lipid content was found in treatment T₃ (mustard oilcake) whereas the highest carbohydrate content was found in treatment T₂, wheat bran ($66.12\pm0.47\%$) followed by T₄, Azolla

(50.21±0.54%), T₆, banana leaf (48.50±0.51%), T₅, grass (46.36±0.16%), T₁, rice bran (44.09±0.67%), T₃, mustard oilcake (32.95±0.29%). Hepher (1988) reported the protein content of ricebran, wheat bran, oil cake and Azolla as 11.88%, 14.57%, 30-33% and 19.27%, respectively. Banerjee and Matai (1990) determined the nutritive status of *Azolla pinnata* and reported protein as 21.9% and Lipid as 3.8%. Gavina (1994) reported crude protein of 20.98%, crude fat of 5.17% and crude fiber of 19.30% in Azolla. Tavares *et al.* (2008) observed 38.8% crude protein, 3.8% crude fat and 13.2% crude fiber in dried duck weed. They also reported that the protein content of duckweeds growing on nutrient poor and nutrient rich water varied between 15-25% and 35-45% (Dry matter basis), respectively.

In case of conventional feed items the major nutrient like protein varied from $14.40\pm0.32\%$ (rice bran) to $30.53\pm0.40\%$ (mustard oilcake). Whereas in case of nonconventional feed items the protein varied from $6.18\pm0.13\%$ (banana leaf) to $18.58\pm0.09\%$ (Azolla). Being an omnivore, the fish can also feed on vegetation (Santhanam *et al.*, 1990) and may be able to assimilate Azolla in the diets.

The chemical composition of Azolla species varies with ecotypes and with the ecological conditions and the phase of growth. The crude protein content is about 19-30 percent dry matter basis during the optimum conditions for growth (Peters *et al.*, 1979; Becking, 1979). The protein contents of Azolla species are comparable to or higher than that of most other aquatic macrophytes. Aquatic weeds' are highly nutritious with protein content of 20-30%, when cultivated in nutrient rich waters (Culley *et al.*, 1981). Importantly, they are preferred food of a wide range of herbivorous fish such as grass carp (*Ctenopharyngodon idella*), silver barb (*Barbonymus gonionotus, Puntius jerdoni*), tilapias (*Oreochromis niloticus, Tilapia rendalli, Tilapia zillii*) and rohu (*Labeo rohita*) (Singh *et al.*, 1967; Gaiger *et al.*, 1984).

Overall findings indicated that inspite of having variations in nutrient contents, monthly supply of nutrients was almost same respective feed item under nonconventional feeds as with conventional feeds. Mean values of the nutrient

contents under non-conventional feed items are found potentials for the development of low cost aquaculture.

Fish feed generally constitutes 60–70% of the operational cost in intensive and semi- intensive aquaculture system (Singh *et al.*, 2006). The fish feed used in aquaculture is quite expensive, irregular and short in supply in many third world countries. These feeds are sometimes adulterated, contaminated with pathogen as well as containing harmful chemicals for human health. Naturally there is a need for the development of healthy, hygienic fish feed which influences the production as well as determines the quality of cultured fish. Considering the importance of nutritionally balanced and cost-effective alternative diets for fish, almost similar expression to evaluate the nutritive value of different non-conventional feed resources, including terrestrial and aquatic macrophytes was found with Wee and Wang (1987) and Mondal and Ray (1999). However potentials roles of aquatic and terrestrial macrophytes as supplementary feeds in fish farming were also found to be expressed with Bardach *et al.* (1972) and Edwards (1990).

3.6: Conclusion

In case of conventional feed items, protein, lipid and carbohydrate varied from $14.40\pm0.32\%$ to $30.53\pm0.40\%$, $6.69\pm0.30\%$ to $13.33\pm0.10\%$ and $32.95\pm0.29\%$ to $66.12\pm0.47\%$. In case of non-conventional feed items, protein, lipid and carbohydrate varied from $6.18\pm0.13\%$ to $18.58\pm0.09\%$, $3.06\pm0.09\%$ to $6.31\pm0.13\%$ and $46.36\pm0.16\%$ to $50.21\pm0.54\%$. Inspite of variations weeds are moderately nutritive and low cost effective diets for fish. However, the present study did not evaluate the fish production and economics of feed and weed based systems.

3.7: Recommendation

Present findings explored the nutritive aspects of both conventional and nonconventional feed items and question raised about the response of utilizing the feed specially of aquatic weeds to fish growth and economics. Therefore, it is recommended to conduct further study on the evaluation of fish production and economics under different feed and weed based systems in polyculture ponds.

CHAPTER FOUR

Evaluation of fish production and economics under conventional feed and weed based systems in polyculture ponds

4.1: Introduction

Supplementary feeds are the major expenditure in any fish culture operation (Mukhopadhyay and Jena, 1999; Fasakin, 1999) which contributes to cost increase per unit production (New and Csavas, 1993). To avoid increasing feed cost with conventional feed items, efforts are being made towards the use of various non-conventional feed sources as ingredients in fish feed (Ali *et al.*, 2006). The weed based system refers to the use of some inputs from plant sources, eg. weeds or grasses or leaves or macrophytes like duckweeds, Azolla etc. as supplemental feed in fish production. These inputs are consumed first as feed by herbivorous fish and subsequently a part of the semi digested faecal matter of the macrophytes feeding fishes are consumed by the other fishes and the remaining part will be recycled in food chain as nutrients for primary production, thus they have potentiality to increase the total fish production of aquaculture system (Grover *et al.*, 2000).

Aquatic weeds have been utilized as food components and thus have played an important role in culture of herbivorous fish since 4000 years ago in Egypt and 2500 years ago in the Orient, including Indian subcontinent (Bardach *et al.*, 1972). Okeyo (1989) listed a number of aquatic macrophytes (*Azolla pinnata*, *Chara* sp., *Eichhornia crassipes*, *Ipomoea aquatica*, *Lemna minor*, *Nymphae* sp., *Trapa natans*, *Wolffia arrhiza*, *Lactuca sativa*, *Typha latifolia*) as potential source of nutrients which are directly used as food components by a number of herbivorous fish. *Azolla pinnata* enhances nitrogen in semi-intensive pisciculture systems (Ayyappan *et al.*, 1993) and acts direct food by some macrophytophagous fish (Cassani, 1981; Antonie *et al.*, 1987). Duckweeds might be having as much potential as fish foods that could be utilized in preparation of suitable fish feed essential in expansion of low-cost aquaculture system in the tropics (Hassan and Edwards, 1992). Fresh duckweed (and also the dried meal) is suited to intensive production of herbivorous fish (Gaiger *et* *al.*, 1984) and duckweed is converted efficiently to live weight gain by carp and tilapia (Van-Dyke and Sutton, 1977; Hepher and Pruginin, 1979; Robinette *et al.*, 1980; Hassan and Edwards, 1992; Skillicorn *et al.*, 1993).

Most of the fish farmers of Bangladesh are resource poor. These resource poor farmers need suitable low cost aquaculture technology to constituting fish farming. Carp polyculture contributes 34.61% of total fish production (DoF, 2012). Weed can be grown abundantly with minimum cost and can be made available for carp polyculture with much cheaper price than other alternative protein sources in Bangladesh (Grover et al., 2000). Researches carried out on the application of weed have so far been centered on the wastewater treatment (Oron, 1994; Alaerts et al., 1996; Vander Steen et al., 1998). However, these research efforts also indicate that comprehensive studies are required to explore the nutrients, production and economics of fish farming based on both conventional and non-conventional feed items specially for the promotion of weed based aquaculture in Bangladesh. Already popularities of locally available weeds to have the major nutrients throughout the major carp polyculture systems are explored well (experiment-1). Unfortunately no effort is found to explore the growth and economics of fish farming of weed based system with conventional feed system. Therefore, the present study aimed at evaluating the fish production and economics under different treatments of weed and feed based systems in polyculture ponds.

4.2 Specific objectives

The specific objectives of this study were as follows-

- to monitor the water quality parameters (in terms of temperature, transparency, dissolved oxygen, pH, alkalinity and CO₂) in ponds under different treatments of weed and feed based systems;
- to evaluate fish production under different treatments of weed and feed based systems;
- to evaluate the economics of fish farming under different treatments of weed and feed based systems; and
- to recommend suitable weed as feed for the application in the carp polyculture ponds.
4.3: Materials and methods

4.3.1: Duration and location of the study

The study was conducted for a period of six months (April 2011 to September 2011) at Alampur village of Sadar Upazila under Kushtia district, Bangladesh (Map 4.1).



Map 4.1 : Shows the study area.

4.3.2: Description of the study ponds

A total of 12 ponds (average water area of 0.18ha and depth of 1.9 m) were selected for the present study. All the ponds were rain-fed and well exposed to sunlight (Plate 4.1).

4.3.3: Experimental design

The experiment was designed under Randomized Completely Block Design (RCBD) with four treatments (T_0 , T_1 , T_2 and T_3) of fish feeds, each with three replications. The treatment assignment was as follows-

- To : Ponds fed with conventional feed like rice bran, wheat bran and mustard oilcake (no weed was used as fish feed).
- T₁ : Azolla (*Azolla pinnata*) fed ponds.
- T₂ : Grass (*Cynodon dactylon*) fed ponds.
- T₃ : Banana (*Musa acuminata*) leaf fed ponds.

Stocking density and stocking ratio/combination of fish species (Table 4.1) were same for all the treatments.

4.3.4: Pond management

Aquatic weeds were removed from all the ponds manually. Predatory fish and other unwanted species were removed through repeated netting.

Liming was done at a rate of 250 kg/ha before 7 days of fertilization. All the ponds were fertilized with cowdung (1500 kg/ha), urea (60 kg/ha) and Triple Super Phosphate (TSP) (60 kg/ha) as basal dose.

For Azolla fed ponds, one tenth area of the research ponds were used as Azolla bank according to Grover *et al.* (2000). Azolla bank was prepared by bamboo fencing (locally called 'Bana'). In Azolla bank, compost manure was deposited (2470 kg/ha). Azolla seeds were introduced in Azolla bank at the rate of 1000 kg/ha for available supply of Azolla during culture period.

Fishes were stocked (11115 fish/ha) in all ponds after five days of basal fertilization. All the ponds were stocked with seven species of fishes (Table 4.1). Mean initial weight of *Hypophthalmichthyes molitrix*, *Catla catla*, *Labeo rohita*, *Cirrhinus cirrhosus*, *Cyprinus carpio*, *Ctenopharyngodon idella* and *Barbonymus gonionotus* were 62, 64, 57, 54, 63, 65 and 25 g, respectively.

Fish seeds were collected from local government fish farm. Stocking of fish seeds were done at early morning.

Pond layer	Fish species	No. of fish/decimal	No. of fish/ha
Surface layer	1. Hypophthalmichthyes molitrix	9	2223
	2. Catla catla	3	741
Middle layer	3. Labeo rohita	10	2470
Bottom layer	4. Cirrhinus cirrhosus	5	1235
	5. Cyprinus carpio	3	741
All layer	6. Ctenopharyngodon idella	5	1235
	7. Barbonymus gonionotus	10	2470
	Total	45	11115

 Table 4.1: Stocking density and stocking ratio/combination of the stocked fish species.

After stocking urea was applied (2.5 kg/ha/day) for ponds under all the treatments except the treatment T_1 (Azolla fed ponds). However, TSP was applied (2.5 kg/ha/day) for ponds under all treatments as periodic fertilization.

In ponds under treatment T_0 commonly available conventional feed ingredients such as rice bran (30%), wheat bran (30%) and oilcake (40%) were used as supplementary feed daily at the rate of 4% of body weight of total fish for the first 2 months (April and May), 3% for the next 2 months (June and July) and finally 2% for the last 2 months (August and September) of culture period. Fish were fed twice a day at 09:00-10:00 hours and at 03:00-04:00 hours with 50% of the ration allocated at each time.

In ponds under treatments T_1 , T_2 and T_3 , Azolla (100%), grass (100%) and banana leaf (100%) were supplied as supplementary feed daily at the rate of 100% of the body weight of herbivorous fishes (*C. idella* and *B. gonionotus*).

Azolla was supplied from Azolla bank, whereas grass and banana leaves were collected locally and chopped into very small pieces during application.

4.3.6: Water quality monitoring

Some important physico-chemical parameters of water such as water temperature, transparency, dissolved oxygen (DO), pH, alkalinity and carbondioxide (CO₂) were monitored fortnightly between 10:00 AM to 11:00 AM for the present study (Plate 4.2).

4.3.6.1: Water temperature

During the study period, water temperature was recorded with the help of a celsius thermometer at 20-30 cm depth of water. The data were expressed as °C.

4.3.6.2: Water transparency

Measurement of limit of visibility i.e. penetration of light in water was done by a Secchi disc. The data thus obtained, were expressed as secchi disc depth in centimeter.

4.3.6.3: Dissolved oxygen (DO)

The dissolved oxygen concentration of water was determined by the aid of a water quality test kit (HACH kit FF-2, USA). Alkaline iodide-azide powder pillows, manganese sulfate powder pillows, sodium thiosulfate titration cartridge (0.2000 N), starch indicator solution and sulfamic acid powder pillows were used for determination of dissolved oxygen. The concentration of dissolved oxygen thus estimated was expressed in milligram per litter (mg/l) of water.

4.3.6.4: Hydrogen ion concentration (pH)

pH was measured by using HACH kit (FF-2, USA). Wide range pH 4 pH indicator solution (1919-00) were used for determination of water pH. A colour comparator disc ranging from 1-14 was also used for this purpose.

4.3.6.5: Alkalinity

Alkalinity was determined through digital titration by the help of a HACH kit (FF-2, USA). Bromcresol green-methyl red powder pillows, phenolphthalein powder pillows and sulfuric acid titration cartridge (0.1600 N) were used for total alkalinity determination. It was also expressed as mg/l of water.

4.3.6.6: Free carbon dioxide (CO₂)

Free carbon dioxide was determined through digital titration by the help of a HACH kit (FF-2, USA). Phenolphthalein powder pillows and sodium hydroxide titration cartridge (0.3636 N) were used for determination of carbon dioxide. It was also expressed as mg/l of water.

4.3.7: Growth monitoring of fishes

At least 10% (by number) of the fish in each pond were randomly sampled on a monthly basis with a cast net. On each sampling day, individual fish from each pond were weighed and measured (Plate 4.3). The purpose was to determine fish growth in weight and to adjust the ration. Following growth parameters were used for the present study.

Weight gain (g) = Mean final weight (g) - Mean initial weight (g)

Final weight (g) = Weight of fish at harvest (g)

Specific Growth Rate, SGR (% bwd⁻¹) = [L_n (final weight) – L_n (initial weight)] / culture period (days) \times 100

(Brown, 1957)

Survival rate (%) = $\frac{\text{Number of fish harvested}}{\text{Number of fish stocked}} \times 100$

Fish yield (kg/ha) = Fish biomass at harvest – Fish biomass at stock

4.3.8: Economics

Simple economical analysis of the different treatments was performed during the study period. On the basis of the fixed and variable expenditure, the total cost (BDT/ha) was estimated. At the end of the experiment, fishes were sold locally and the gross benefit (BDT/ha) was estimated. Net benefit (BDT/ha) was estimated by deducting the total cost from gross benefit. The following parameters were used to explore the economics of different treatments:

Net profit margin (%) = $\frac{\text{Net benefit}}{\text{Total investment}} \times 100$ CBR (Cost-Benefit Ratio) = $\frac{\text{Net benefit}}{\text{Total investment}}$

4.3.9: Statistical analysis

Data on water quality parameters, fish production and economics under different treatments were subjected to one way ANOVA (Analysis of Variance) using computer software SPSS (Statistical Package for Social Science, version-11). The mean values were also compared to see the significant difference from the DMRT (Duncan Multiple Range Test) after Gomez and Gomez (1984).





Plate 4.1: Experimental ponds.



Plate 4.2 Monitoring water quality parameters.





Plate 4.3 Measurement of growth of fishes.

4.4: Results

4.4.1: Water quality

4.4.1.1: Fortnightly variations

The variations in the mean values of water quality parameters in different treatments at different fortnights are presented in table 4.2 to table 4.13.

Water temperature

During the study period, water temperature varied from 29.58 ± 0.03 to 32.23 ± 0.08 °C. The minimum value was recorded in treatment T₀ at 1st fortnight whereas the maximum value was recorded in treatment T₀ at 6th fortnight.

Water transparency

Water transparency value was found to range from 30.00 ± 0.58 to 35.00 ± 0.00 cm. The minimum value was recorded in treatment T₂ at 7th fortnight whereas the maximum value was recorded in treatment T₁ at 1st fortnight.

Dissolved Oxygen (DO)

Dissolved oxygen of water varied from 4.90 ± 0.06 to 6.17 ± 0.12 mg/l. The minimum value was recorded in treatment T₂ at 6th fortnight whereas the maximum value was recorded in treatment T₁ at 10th fortnight.

Hydrogen ion concentration (pH)

pH value of water varied from 7.03 \pm 0.03 to 7.87 \pm 0.09. The minimum value was recorded in treatment T₃ at 6th fortnight whereas the maximum value was recorded in treatment T₀ at 12th fortnight.

Alkalinity

Alkalinity of water was found to range from 109.22 ± 1.33 to 120.00 ± 1.15 mg/l. The minimum value was recorded in treatment T₃ at 1st fortnight whereas the maximum value was recorded in treatment T₀ at 9th fortnight.

Free carbon dioxide (CO₂)

Free CO₂ value was found to range from 2.49 ± 0.07 to 3.46 ± 0.03 mg/l. The minimum value was recorded in treatment T₁ at 4th fortnight whereas the maximum value was recorded in treatment T₃ at 12th fortnight.

Among the water quality parameters the values of transparency at 1^{st} , 10^{th} and 12^{th} fortnights, DO at 6^{th} , 9^{th} , 11^{th} and 12^{th} fortnights, pH at 12^{th} fortnight, alkalinity at 8^{th} and 9^{th} fortnights and CO₂ at 10^{th} fortnight differed significantly with the treatments.

Treatments				
	To	T₁	T ₂	T ₃
Parameters				
Water temperature (°C)	29.58±0.03ª	30.12±0.21ª	30.23±0.14ª	30.05±0.09ª
Transparency (cm)	32.67±0.88a ^b	35.00±0.00ª	32.00±1.15 ^b	32.33±0.67 ^{ab}
DO (mg/l)	5.18±0.12ª	5.32±0.06ª	5.13±0.09ª	5.17±0.09ª
рН	7.20±0.06ª	7.40±0.06ª	7.10±0.06ª	7.20±0.10 ^ª
Alkalinity (mg/l)	116.00±1.15ª	118.23±0.45ª	114.33±4.18ª	109.22±1.33ª
Free CO ₂ (mg/l)	3.23±0.02ª	2.52±0.07 ^a	3.12±0.03ª	3.02±0.06ª

Table-4.2: Variations in the mean values of water quality parameters under different treatments at 1st fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.3: Variations	in th	e mean	values	of	water	quality	parameters	under	different
treatments a	t 2 nd	fortnigh	t						

Treatments	-	-	-	-
Parameters	To	T ₁	12	13
Water temperature (°C)	30.46±0.14ª	30.69±0.23ª	30.85±0.06ª	30.58±0.06ª
Transparency (cm)	31.67±0.67ª	31.00±0.00ª	31.33±0.33ª	31.67±1.67ª
DO (mg/l)	5.27±0.09ª	5.35±0.15ª	5.00±0.12ª	5.27±0.03ª
рН	7.13±0.09ª	7.30±0.09ª	7.13±0.09ª	7.17±0.09ª
Alkalinity (mg/l)	113.33±1.45ª	112.00±1.73ª	118.00±3.51ª	111.33±3.28ª
Free CO ₂ (mg/l)	3.31±0.11ª	2.85±0.06ª	3.37±0.02ª	3.33±0.12ª

Treatments Parameters	Τo	Τ1	T2	Тз
Water temperature (°C)	31.61±0.09ª	30.98±0.05ª	31.25±0.11ª	31.54±0.09ª
Transparency (cm)	34.00±0.58ª	32.00±1.00ª	32.33±1.45ª	31.33±0.33ª
DO (mg/l)	5.12±0.17ª	5.36±0.15ª	5.13±0.03ª	5.17±0.09ª
рН	7.20±0.12ª	7.32±0.15ª	7.21±0.06ª	7.17±0.09ª
Alkalinity (mg/l)	110.00±1.73ª	111.00±2.08ª	113.33±0.33ª	112.33±2.03ª
Free CO ₂ (mg/l)	3.12±0.03 ^a	2.95±0.01ª	3.02±0.23 ^a	3.09±0.11ª

Table-4.4: Variations in the mean values of water quality parameters under different treatments at 3rd fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.5: Variations in the mean values of water quality parameters under different treatments at 4th fortnight

Treatments Parameters	T₀	T1	T2	T₃
Water temperature (°C)	31.98±0.09ª	31.74±0.09ª	31.67±0.02ª	31.59±0.06ª
Transparency (cm)	33.33±0.67ª	33.00±0.58ª	33.67±0.88ª	33.00±1.53ª
DO (mg/l)	5.03±0.09ª	5.20±0.06ª	4.97±0.12ª	5.07±0.12ª
рН	7.17±0.07ª	7.27±0.07ª	7.20±0.12ª	7.10±0.15ª
Alkalinity (mg/l)	112.67±1.45ª	109.67±0.88ª	113.00±3.61ª	117.00±2.52ª
Free CO ₂ (mg/l)	2.61±0.03ª	2.49±0.07ª	2.58±0.03ª	2.85±0.07ª

Treatments Parameters	Τo	Τı	T2	T₃
Water temperature (°C)	32.15±0.11ª	31.92±0.06ª	32.08±0.14ª	31.96±0.02ª
Transparency (cm)	32.67±0.88ª	33.00±1.53ª	33.00±1.53ª	33.00±1.15ª
DO (mg/l)	5.10±0.10ª	5.23±0.03ª	5.17±0.07ª	4.93±0.13ª
рН	7.30±0.06ª	7.40±0.06ª	7.27±0.09ª	7.33±0.09ª
Alkalinity (mg/l)	114.00±2.31ª	113.33±4.26ª	110.33±0.88ª	117.00±2.08ª
Free CO ₂ (mg/l)	2.72±0.05ª	2.55±0.06ª	2.74±0.08ª	2.77±0.05ª

Table-4.6: Variations in the mean values of water quality parameters under different treatments at 5th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.7: Variations in the mean values of water quality parameters under different treatments at 6th fortnight

Treatments Parameters	Τo	Τı	T2	T₃
Water temperature (°C)	32.23±0.08ª	32.04±0.24ª	32.16±0.08ª	31.98±0.02ª
Transparency (cm)	31.67±0.33ª	32.33±0.88ª	31.33±0.88ª	32.33±1.45ª
DO (mg/l)	5.23±0.09ª	5.30±0.06ª	4.90±0.06 ^b	5.10±0.00ª
рН	7.13±0.09ª	7.15±0.03ª	7.13±0.15ª	7.03±0.03ª
Alkalinity (mg/l)	115.00±1.53ª	113.67±3.84ª	109.33±0.88ª	112.67±0.88ª
Free CO ₂ (mg/l)	2.84±0.04ª	2.72±0.11ª	2.80±0.12ª	2.84±0.05ª

Treatments Parameters	Τo	Τı	T2	T₃
Water temperature (°C)	31.85±0.14ª	31.92±0.11ª	32.05±0.15ª	31.98±0.06ª
Transparency (cm)	31.00±0.58ª	30.67±0.67ª	30.00±0.58ª	31.33±0.33ª
DO (mg/l)	5.03±0.09ª	5.23±0.03ª	5.16±0.14ª	5.07±0.09ª
рН	7.13±0.09ª	7.20±0.12ª	7.07±0.12ª	7.10±0.06ª
Alkalinity (mg/l)	114.33±1.67ª	116.00±2.52ª	114.67±1.33ª	114.67±2.03ª
Free CO ₂ (mg/l)	2.87±0.09ª	2.79±0.06ª	2.89±0.03ª	2.86±0.06ª

Table-4.8: Variations in the mean values of water quality parameters under different treatments at 7th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.9: Variations in the mean values of water quality parameters under different treatments at 8th fortnight

Treatments Parameters	Τo	Τı	T2	T₃
Water temperature (°C)	31.71±0.08ª	31.62±0.02ª	31.55±0.12ª	31.39±0.05ª
Transparency (cm)	33.00±0.58ª	31.67±0.33ª	33.00±1.15ª	32.67±1.33ª
DO (mg/l)	5.07±0.07ª	5.20±0.10ª	5.13±0.07ª	5.13±0.03ª
рН	7.22±0.09ª	7.33±0.03ª	7.17±0.03ª	7.20±0.06ª
Alkalinity (mg/l)	115.33±1.45 ^{ab}	117.67±1.33ª	110.33±2.73⁵	110.33±2.03 [♭]
Free CO ₂ (mg/l)	2.91±0.12ª	2.85±0.06ª	2.92±0.09ª	2.90±0.06ª

Treatments Parameters	Τo	Τı	T2	T₃
Water temperature (°C)	31.37±0.09ª	31.36±0.18ª	31.26±0.12ª	31.24±0.10ª
Transparency (cm)	32.67±0.88ª	33.33±0.33ª	32.33±0.88ª	33.00±0.58ª
DO (mg/l)	5.73±0.07 ^{ab}	5.90±0.06ª	5.87±0.03ª	5.67±0.03 ^b
рН	7.20±0.12ª	7.31±0.12ª	7.24±0.07ª	7.17±0.07ª
Alkalinity (mg/l)	120.00±1.15ª	116.33±0.88 ^b	118.67±0.88ªb	118.00±1.15 ^{ab}
Free CO ₂ (mg/l)	2.98±0.03ª	2.88±0.06ª	2.96±0.03ª	2.95±0.00ª

Table-4.10: Variations in the mean values of water quality parameters under different treatments at 9th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.11: Variations in the mean values of water quality parameters under different treatments at 10th fortnight

Treatments Parameters	Τo	T ₁	T2	T ₃
Water temperature (°C)	31.44±0.09ª	31.23±0.03ª	31.31±0.17ª	31.21±0.05ª
Transparency (cm)	32.00±0.58 ^b	34.67±0.33ª	34.00±0.58ª	34.00±0.58ª
DO (mg/l)	6.13±0.09ª	6.17±0.12ª	6.13±0.09ª	6.13±0.03ª
рН	7.40±0.03ª	7.45±0.09ª	7.20±0.06ª	7.33±0.09ª
Alkalinity (mg/l)	115.67±1.20ª	115.00±1.00ª	117.00±1.15ª	114.33±0.88ª
Free CO ₂ (mg/l)	3.14±0.02ª	2.88±0.03 ^b	3.10±0.03 ^b	2.95±0.03 ^b

Treatments Parameters	Τo	T ₁	T ₂	T ₃
Water temperature (°C)	30.48±0.06ª	30.27±0.02ª	30.26±0.06ª	30.29±0.12ª
Transparency (cm)	31.67±0.88ª	33.00±0.58ª	32.00±1.00ª	31.67±0.88ª
DO (mg/l)	6.00±0.06 ^{ab}	6.05±0.03ª	5.87±0.03 ^{bc}	5.80±0.06 ^c
рН	7.27±0.13ª	7.56±0.09ª	7.50±0.12ª	7.33±0.03ª
Alkalinity (mg/l)	114.33±2.33ª	115.67±0.67ª	114.75±0.67ª	114.38±0.88ª
Free CO ₂ (mg/l)	3.22±0.03ª	3.05±0.03ª	3.10±0.03ª	3.20±0.06ª

Table-4.12: Variations in the mean values of water quality parameters under different treatments at 11th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.13: Variations in the mean values of water quality parameters under different treatments at 12th fortnight

Treatments Parameters	T₀	T₁	T2	T₃
Water temperature (°C)	30.63±0.08ª	30.47±0.06ª	30.23±0.12ª	30.45±0.03ª
Transparency (cm)	31.67±0.67⁵	34.33±0.33ª	32.33±0.88 ^{ab}	33.00±0.58 ^{ab}
DO (mg/l)	5.80±0.06ª	5.83±0.03ª	5.67±0.09 ^{ab}	5.53±0.03 ^b
рН	7.87±0.09ª	7.37±0.18 ^{bc}	7.73±0.12 ^{ab}	7.07±0.03°
Alkalinity (mg/l)	114.32±1.45ª	116.21±2.67ª	114.81±1.53ª	115.25±0.54ª
Free CO ₂ (mg/l)	3.45±0.03ª	3.30±0.01ª	3.38±0.03ª	3.46±0.03ª

4.4.1.2: Mean variations

The variations in the mean vales of different water quality parameters in different treatments by the total of all fortnights are presented in table 4.14 and fig. 4.1.

Water temperature

During the study period the mean values of water temperature ranged from 31.19 ± 0.20 to 31.29 ± 0.24 °C. The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₀.

Water transparency

The mean value of water transparency varied from 32.28 ± 0.32 to 32.83 ± 0.40 cm. The minimum value was recorded in treatment T₂ whereas the maximum value was recorded in treatment T₁.

Dissolved Oxygen (DO)

The mean value of DO varied from 5.33 ± 0.09 to 5.51 ± 0.11 mg/l. The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₁.

Hydrogen ion concentration (pH)

The mean value of water pH varied from 7.18 ± 0.03 to 7.38 ± 0.05 . The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₁.

Alkalinity

The mean value of total alkalinity varied from 113.61 ± 0.74 to 114.58 ± 0.67 mg/l. The minimum value was recorded in treatment T₁ whereas the maximum value was recorded in treatment T₀.

Free carbon-dioxide (CO₂)

The mean value of free CO₂ varied from 2.82 ± 0.07 to 3.03 ± 0.07 . The minimum value was recorded in treatment T₁ whereas the maximum value was recorded in treatment T₀.

No significant difference was found among the treatments for mean values of water quality parameters.

Treatments	To	T ₁	T ₂	T ₃
Parameters				
Water temperature (°C)	31.29±0.24ª	31.20±0.20ª	31.24±0.21ª	31.19±0.20ª
Transparency (cm)	32.34±0.25ª	32.83±0.40ª	32.28±0.32ª	32.44±0.24ª
DO (mg/l)	5.39±0.12ª	5.51±0.11ª	5.34±0.12ª	5.33±0.09ª
рН	7.22±0.03ª	7.38±0.05ª	7.25±0.05ª	7.18±0.03ª
Alkalinity (mg/l)	114.58±0.67ª	113.61±0.74ª	114.05±0.87ª	113.94±0.80ª
Free CO ₂ (mg/l)	3.03±0.07 ^a	2.82±0.07 ^a	3.00±0.07 ^a	3.02±0.06 ^a

Table 4.14: Variations in the mean values of water quality parameters under different treatments during study period















Fig. 4.1: Variations in the mean values of water quality parameters (a; Water temperature, b; transparency, c; DO, d; pH, e; alkalinity, f; free CO₂) under different treatments during study period.

4.4.2: Growth of fishes

4.4.2.1: Monthly variations

Specific growth rate (SGR, %, bwd⁻¹)

The specific growth rate (%, bwd⁻¹) varied with *H. molitrix* from 0.30 ± 0.01 (T₃ at 6th month) to 3.13 ± 0.03 (T₀ at 1st month), *C. catla* from 0.28 ± 0.01 (T₂ and T₃ at 6th month) to 2.49 ± 0.09 (T₁ 1st month), *L. rohita* from 0.23 ± 0.04 (T₁ at 6th month) to 2.87 ± 0.03 (T₀ at 1st month), *C. cirrhosus* from 0.31 ± 0.05 (T₃ at 6th month) to 2.91 ± 0.04 (T₀ 1st month), *C. carpio* from 0.30 ± 0.03 (T₃ at 6th month) to 2.70 ± 0.03 (T₀ at 1st month), *C. idella* from 0.34 ± 0.01 (T₀ 6th month) to 3.15 ± 0.06 (T₁ at 1st month) and *B. gonionotus* from 0.14 ± 0.01 (T₂ at 6th month) to 3.01 ± 0.06 (T₁ at 1st month) (Table 4.15 to 4.20).

Among the different species, the highest SGR was observed in *C. idella* as 3.15 ± 0.06 %, bwd⁻¹ and the lowest was in *B. gonionotus* as 0.14 ± 0.01 %, bwd⁻¹.

The lowest combined SGR value was found as 0.30 ± 0.03 (T₂ at 6th month) and the highest combined SGR was found as 2.62 ± 0.17 (T₀ at 1st month). The highest SGR value was found in the month of April and the lowest value was found in the month of September for all the species.

Significant variation in SGR values with the treatments was found in all months except the 2nd month (May) for *C. carpio*, 2nd month (May) and 3rd month (June) for *C. idella*, 4th month (July), 5th month (August) and 6th month (September) for *C. cirrhosus* and 4th month (July) for *B. gonionotus*.

Treatments				
	T₀	T1	T2	Тз
Species				
H. molitrix	3.13±0.03ª	2.49±0.09 ^b	1.84±0.04°	1.65±0.04 ^d
C. catla	1.77±0.05 ^b	2.49±0.09ª	1.95±0.05 [⊾]	1.83±0.07⁵
L. rohita	2.87±0.03ª	1.91±0.07 ^b	1.42±0.06°	1.36±0.04°
C. cirrhosus	2.91±0.04ª	2.07±0.23 ^b	1.28±0.11°	1.22±0.09°
C. carpio	2.70±0.03ª	2.25±0.16 ^b	1.73±0.08°	1.65±0.04°
C. idella	2.63±0.26 ^b	3.15±0.06ª	2.51±0.06 ^b	2.48±0.04 ^b
B. gonionotus	2.36±0.32 ^b	3.01±0.06ª	1.59±0.12°	1.59±0.07°
All species	2.62±0.17ª	2.48±0.17ª	1.76±0.15 [♭]	1.68±0.15 [♭]

Table-4.15: Variations in the mean values of SGR (%, bwd⁻¹) of fishes under different treatments at 1st month (April)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.16:	Variations	in th	ne meai	n values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments	at 2 ^r	^{id} month	n (May)			-	-				

Treatments	To	T ₁	T2	T₃
Species				
H. molitrix	1.61±0.01ª	1.47±0.04 ^b	1.29±0.01°	1.38±0.04 ^ь
C. catla	2.07±0.07ª	1.49±0.04 ^b	1.38±0.04 ^b	1.43±0.05 ^b
L. rohita	1.55±0.02ª	1.35±0.06 ^b	1.20±0.04°	1.18±0.01°
C. cirrhosus	1.58±0.01ª	1.37±0.09 ^{ab}	1.19±0.06 ^b	1.18±0.08 ^b
C. carpio	1.52±0.02ª	1.46±0.12ª	1.35±0.05ª	1.39±0.03ª
C. idella	1.56±0.09ª	1.59±0.03ª	1.57±0.04ª	1.55±0.01ª
B. gonionotus	1.57±0.06ª	1.68±0.01ª	1.35±0.03⁵	1.27±0.06 ^b
All species	1.64±0.07a	1.49±0.04b	1.33±0.05c	1.34±0.05c

Treatments	To	T 1	T ₂	T ₃	
Species					
H. molitrix	1.16±0.003ª	1.13±0.02 ^{ab}	1.08±0.01 ^b	1.08±0.02 ^b	
C. catla	1.40±0.01ª	1.15±0.02 ^b	1.04±0.01°	1.03±0.05°	
L. rohita	1.17±0.02ª	1.08±0.03 ^b	0.96±0.03°	1.00±0.03 ^{bc}	
C. cirrhosus	1.20±0.01ª	1.11±0.03 ^b	1.06±0.02 ^b	1.07±0.02 ^b	
C. carpio	1.13±0.003ª	1.10±0.03 ^{ab}	1.06±0.01 ^b	1.05±0.01 ^ь	
C. idella	1.15±0.01ª	1.16±0.02ª	1.12±0.003ª	1.12±0.01ª	
B. gonionotus	1.11±0.003 ^b	1.33±0.01ª	1.05±0.01°	0.97±0.03 ^d	
All species	1.19±0.04ª	1.15±0.03ª	1.05±0.02 ^b	1.05±0.02 ^b	

Table-4.17:	Variations	in the	mean	values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments	at 3rd	month	(June)								

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.18:	Variations in	n the me	an values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments a	at 4 th mor	th (July)								

Treatments Species	T₀	T1	T2	T₃
H. molitrix	0.88±0.01ª	0.82±0.02 ^b	0.74±0.02°	0.76±0.01°
C. catla	1.35±0.02ª	0.85±0.01 ^b	0.71±0.01°	0.74±0.01°
L. rohita	0.83±0.01ª	0.76±0.01 ^b	0.70±0.01°	0.70±0.03°
C. cirrhosus	0.83±0.003ª	0.82±0.02ª	0.81±0.03ª	0.85±0.01ª
C. carpio	0.89±0.01ª	0.82±0.03 ^b	0.75±0.01 ^b	0.75±0.03 ^b
C. idella	0.83±0.02 ^{ab}	0.88±0.03ª	0.79±0.01 ^b	0.79±0.01 ^b
B. gonionotus	0.76±0.02ª	0.81±0.05ª	0.70±0.02ª	0.69±0.08ª
All species	0.91±0.08ª	0.82±0.01 ^b	0.74±0.02°	0.75±0.02°

Treatments Species	Τo	T1	T2	Тз
H. molitrix	0.58±0.01ª	0.54±0.02 ^{ab}	0.50±0.02 ^{bc}	0.47±0.01°
C. catla	0.74±0.01ª	0.51±0.01 ^ь	0.43±0.01°	0.44±0.02°
L. rohita	0.48±0.01ª	0.38±0.02 ^b	0.49±0.03ª	0.49±0.01ª
C. cirrhosus	0.53±0.01ª	0.52±0.01ª	0.57±0.02ª	0.57±0.03ª
C. carpio	0.56±0.01ª	0.52±0.01 ^ь	0.48±0.01°	0.46±0.01°
C. idella	0.46±0.01 ^b	0.56±0.02ª	0.49±0.01 ^b	0.48±0.03 ^b
B. gonionotus	0.35±0.01 ^b	0.50±0.03ª	0.30±0.04 ^b	0.34±0.01 ^b
All species	0.53±0.05ª	0.50±0.02ª	0.47±0.03 ^b	0.46±0.08 ^b

Table-4.19: Variations in the mean values of SGR (%, bwd⁻¹) of fishes under different treatments at 5th month (August)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.20:	Variations in	ו the	mean	values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments a	i t 6 th 1	month	(Septer	nbe	ər)	-	-				

Treatments				
	Τo	T ₁	T ₂	T ₃
Species				
H. molitrix	0.40±0.01ª	0.40±0.01ª	0.32±0.01 ^b	0.30±0.01 ^b
C. catla	0.54±0.01ª	0.37±0.02 ^b	0.28±0.01°	0.28±0.01°
L. rohita	0.35±0.01ª	0.23±0.04°	0.26±0.02 ^{bc}	0.33±0.01 ^{ab}
C. cirrhosus	0.37±0.003ª	0.38±0.01ª	0.38±0.01ª	0.31±0.05ª
C. carpio	0.40±0.01ª	0.38±0.01ª	0.36±0.02ª	0.30±0.003 ^b
C. idella	0.34±0.01 ^b	0.42±0.03ª	0.37±0.01 ^{ab}	0.40±0.01 ^{ab}
B. gonionotus	0.27±0.01 ^b	0.37±0.03ª	0.14±0.01°	0.25±0.02 ^b
All species	0.38±0.03ª	0.36±0.02ª	0.30±0.03ª	0.31±0.02ª

Monthly weight gain (g/month)

The weight gain (g/month) varied with *H. molitrix* from 28.67 ± 0.67 g (T₃ at 6th month) to 109.67 ± 2.03 g (T₀ at 4th month), *C. catla* from 28.33 ± 1.20 g (T₃ at 6th month) to 154.33 ± 3.84 g (T₀ at 4th month), *L. rohita* from 19.33 ± 1.45 g (T₂ at 6th month) to 90.67 ± 1.20 g (T₀ at 3rd month), *C. cirrhosus* from 22.67 ± 3.33 g (T₃ at 6th month) to 90.00 ± 0.58 g (T₀ at 3rd month), *C. carpio* from 29.67 ± 0.33 (T₃ at 6th month) to 96.00 ± 1.53 g (T₀ at 4th month), *C. idella* from 53.33 ± 0.88 (T₂ at 6th month) to 111.33 ± 4.37 g (T₁ at 3rd month) and *B. gonionotus* from 4.67 ± 0.33 (T₂ at 6th month) to 50.00 ± 0.58 g (T₁ at 3rd month) (Table 4.21 to 4.26).

Among the different species, the highest monthly weight gain was observed in *C. catla* as 154.33 ± 3.84 g and the lowest was in *B. gonionotus* as 4.67 ± 0.33 g.

The lowest combined weight gain value was found as 29.10 ± 5.66 (T₂ at 6th month) and the highest combined weight was found as 94.52 ± 13.75 (T₀ at 4th month). The highest weight gain was found in the month of July and the lowest value was found in the month of September for all the species.

Significant variation was found in all months for all the species except the 2nd month (May) and 6th month (September) for *C. idella*.

Treatments				
	Τo	T1	T ₂	T ₃
Species				
H. molitrix	96.33±1.45ª	69.00±3.46 ^b	45.67±1.45°	39.67±1.20℃
C. catla	45.00±1.73 ^b	71.00±3.61ª	51.00±1.73 ^b	47.00±2.31 ^b
L. rohita	78.00±1.15ª	44.00±2.00 ^b	30.33±1.45°	28.67±0.88°
C. cirrhosus	75.33±1.45ª	47.00±7.00 ^b	25.33±2.60°	24.00±2.00°
C. carpio	78.67±1.45ª	61.00±6.08 ^b	43.00±2.65°	40.33±1.20°
C. idella	79.00±11.27 ^b	102.33±2.96ª	73.00±2.65 ^b	71.67±1.67 ^b
B. gonionotus	26.33±5.04 ^b	36.67±1.20ª	15.33±1.45°	15.33±0.88°
All species	68.38±9.07ª	61.57±8.38ªb	40.52±7.18 ^b	38.10±6.93 ^b

Table-4.21: Variations in the mean weight gain (g/month) of fishes under different treatments at 1st month (April)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.22: Variations in the mean weight gain (g/month) of fishes under different treatments at 2nd month (May)

Treatments				
	Τo	T ₁	T ₂	T ₃
Species				
H. molitrix	98.67±0.88ª	73.00±4.36 ^b	50.67±0.33°	52.33±1.33°
C. catla	93.67±2.73ª	76.00±4.36 ^b	59.00±1.15°	59.33±2.33°
L. rohita	80.00±0.58ª	50.67±3.48 ^b	37.67±0.88°	36.33±0.33°
C. cirrhosus	78.67±1.45ª	52.00±7.81 ^b	34.00±1.00°	33.00±1.73°
C. carpio	82.00±1.15ª	69.00±10.82 ^{ab}	53.00±1.00 ^b	53.33±1.20 ^b
C. idella	87.00±13.08ª	102.33±4.48ª	82.67±0.88ª	80.67±0.67ª
B. gonionotus	31.00±3.51 ^b	40.33±0.88ª	20.00±0.58°	18.67±0.67°
All species	78.72±8.42ª	66.19±7.82 ^b	48.14±7.62°	47.67±7.66°

Treatments Species	Τo	T1	T2	T₃
H. molitrix	107.33±1.20ª	83.00±5.29 ^b	60.33±0.33°	59.00±1.15°
C. catla	105.67±1.20ª	87.00±5.20 ^b	64.00±1.15°	61.67±2.33°
L. rohita	90.67±1.20ª	58.33±2.19 ^b	41.67±1.76°	42.67±1.45°
C. cirrhosus	90.00±0.58ª	61.00±7.00 ^ь	42.33±0.33°	42.00±1.00°
C. carpio	90.67±1.20ª	76.00±8.89ª	59.33±0.33 ^b	58.33±0.88 ^b
C. idella	97.00±12.29 ^{ab}	111.33±4.37ª	88.33±0.88°	87.00±1.00°
B. gonionotus	32.33±3.38 ^b	50.00±0.58ª	22.33±0.88°	20.00±0.58°
All species	87.67±9.62ª	75.24±7.91 ^b	54.05±7.93°	52.95±7.89°

Table-4.23: Variations in the mean weight gain (g/month) of fishes under different treatments at 3rd month (June)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.24:	Variations	in	the	mean	weight	gain	(g/month)	of	fishes	under	different
	treatments	at	4 th m	onth (J	July)						

Treatments Species	Τo	Τı	T2	T₃
H. molitrix	109.67±2.03ª	80.33±6.17 ^b	54.00±1.15°	54.33±1.20°
C. catla	154.33±3.84ª	85.67±3.71 ^b	56.33±0.67°	57.33±0.88°
L. rohita	86.67±0.88ª	54.00±2.00 ^b	39.00±0.58°	38.33±1.33°
C. cirrhosus	84.33±1.45ª	59.00±5.13 ^b	43.00±2.08°	44.33±0.33°
C. carpio	96.00±1.53ª	74.00±3.61 ^b	55.33±1.45°	54.33±1.86°
C. idella	99.00±10.58 ^{ab}	107.00±5.13ª	82.00±2.31°	81.67±1.67°
B. gonionotus	31.67±3.53ª	38.67±0.67ª	19.33±1.33 ^b	18.00±2.08 ^b
All species	94.52±13.75ª	71.24±8.57 ^b	49.86±7.27°	49.76±7.38°

Treatments Species	Τo	T₁	T2	Тз
H. molitrix	90.67±1.20ª	65.00±5.51 ^b	44.33±2.33°	40.33±1.20°
C. catla	114.67±3.18ª	63.00±3.61 ^b	40.33±0.88°	40.67±1.67°
L. rohita	60.33±0.88ª	32.67±2.91 ^b	32.33±2.19 ^b	32.00±1.15 ^b
C. cirrhosus	65.67±1.45ª	46.33±5.04 ^b	37.00±1.53 ^b	36.67±2.03 ^b
C. carpio	75.00±2.08ª	58.33±3.93 ^b	42.67±1.20°	40.00±1.15°
C. idella	72.00±1.00 ^{ab}	79.67±10.90ª	61.33±0.67 ^{ab}	60.00±2.52 ^b
B. gonionotus	21.33±0.88ª	24.00±3.21ª	9.67±1.20 ^b	10.33±0.33 ^b
All species	71.38±10.82ª	52.71±7.37⁵	38.24±5.87°	37.14±5.56°

Table-4.25: Variations in the mean weight gain (g/month) of fishes under different treatments at 5th month (August)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.26:	Variations	in	the	mean	weight	gain	(g/month)	of	fishes	under	different
	treatments	at 6	ծ th m	onth (S	Septemb	oer)					

Treatments				
	Τo	T₁	T ₂	Тз
Species				
H. molitrix	72.00±1.15ª	55.33±5.49 ^b	32.00±1.00°	28.67±0.67°
C. catla	101.33±3.53ª	52.33±5.90 ^b	29.67±1.67°	28.33±1.20°
L. rohita	49.33±1.20ª	21.33±4.67 ^b	19.33±1.45⁵	24.67±0.67 ^b
C. cirrhosus	52.00±1.15ª	39.33±5.24 ^b	28.00±1.53°	22.67±3.33°
C. carpio	61.33±1.86ª	48.67±4.41 ^b	36.67±2.19°	29.67±0.33°
C. idella	60.33±0.88ª	68.33±11.22ª	53.33±0.88ª	56.00±2.08ª
B. gonionotus	17.67±0.67ª	20.00±3.21ª	4.67±0.33 ^b	8.33±0.88 ^b
All species	59.14±9.54ª	43.62±6.77 ^b	29.10±5.66°	28.33±5.37°

4.4.2.2: Mean variations

Variations in the mean values of growth parameters of fishes under different treatments are shown in table 4.27 to 4.31, fig. 4.2 to 4.4 and plate 4.4.

SGR (%, bwd⁻¹)

The specific growth rate (SGR, %, bwd⁻¹) varied with *H. molitrix* from 0.94 ± 0.21 (T₃) to 1.29 ± 0.41 (T₀), *C. catla* from 0.96 ± 0.24 g (T₃) to 1.31 ± 0.24 (T₀), *L. rohita* from 0.84 ± 0.16 (T₃) to 1.21 ± 0.38 (T₀), *C. cirrhosus* from 0.87 ± 0.15 (T₃) to 1.24 ± 0.38 (T₀), *C. carpio* from 0.93 ± 0.22 (T₃) to 1.20 ± 0.34 (T₀), *C. idella* from 1.03 ± 0.23 (T₀) to 1.31 ± 0.31 (T₁) and *B. gonionotus* from 0.85 ± 0.21 (T₃) to 1.05 ± 0.25 (T₁).

Among the different species the highest SGR was observed as 1.31 ± 0.31 %, bwd⁻¹ in *C. idella* and the lowest was in *L. rohita* as 0.84 ± 0.16 %, bwd⁻¹.

The lowest combined SGR (%, bwd⁻¹) value was found as 0.93 ± 0.03 (T₃) and the highest combined SGR (%, bwd⁻¹) found as 1.17 ± 0.05 (T₀).

Significant difference was found among the treatments for the mean values of SGR for all the species.

Weight gain (g/month)

Mean weight gain (g/month) varied with *H. molitrix* from 45.72 ± 4.66 g (T₃) to 95.78 ± 5.56 g (T₀), *C. catla* from 49.06 ± 5.28 g (T₃) to 102.45 ± 14.41 g (T₀), *L. rohita* from 33.39 ± 3.30 g (T₂) to 74.17 ± 6.55 g (T₀), *C. cirrhosus* from 33.78 ± 3.68 g (T₃) to 74.33 ± 5.59 g (T₀), *C. carpio* from 46.00 ± 4.51 g (T₃) to 80.61 ± 4.99 g (T₀), *C. idella* from 72.84 ± 5.13 g (T₃) to 95.17 ± 6.99 g (T₁) and *B. gonionotus* from 15.11 ± 1.95 g (T₃) to 34.95 ± 4.53 g (T₁).

Among the different species the highest weight gain was observed as 102.45 ± 14.41 g in *C. catla* and the lowest was in *B. gonionotus* as 15.11 ± 1.95 g.

Considering all the species the lowest mean weight gain was found as 42.33 ± 6.71 g (T₃) and the highest weight gain was found as 76.64 ± 9.24 g (T₀).

Significant difference was found among the treatments for the mean values of weight gain for all the species.

Final weight (g)

The mean final weight (g) varied with *H. molitrix* from 336.33 ± 2.03 (T₃) to 636.67 ± 5.61 g (T₀), *C. catla* from 358.33 ± 4.41 (T₃) to 678.67 ± 12.57 g (T₀), *L. rohita* from 257.33 ± 5.04 (T₂) to 502.00 ± 2.08 g (T₀), *C. cirrhosus* from 256.67 ± 1.76 (T₃) to 500.00 ± 7.23 g (T₀), *C. carpio* from 339.00 ± 3.79 (T₃) to 546.67 ± 8.69 g (T₀), *C. idella* from 502.00 ± 4.16 (T₃) to 620.33 ± 18.02 g (T₁) and *B. gonionotus* from 115.67 ± 2.60 (T₃) to 229.67 ± 2.67 g (T₁).

Among the different species the highest final weight was observed as 678.67±12.57 g in *C. catla* and the lowest was in *B. gonionotus* as 115.67±2.60 g.

Considering all the species the lowest mean final weight was found as 2167.67 \pm 17.31 g (T₃) and the highest final weight was 3629.34 \pm 19.52 g (T₀) for all the species.

Significant difference was found among the treatments for the mean values of final weight for all the species.

Survival rate (%)

The survival rate (%) varied with *H. molitrix* from 81.33 ± 1.20 (T₂) to $84.33\pm2.03\%$ (T₀), *C. catla* from 75.33 ± 1.59 (T₂) to $81.00\pm1.73\%$ (T₀), *L. rohita* from 77.67 ± 1.45 (T₃) to $81.33\pm0.88\%$ (T₀), *C. cirrhosus* from 67.50 ± 1.44 (T₂) to $80.33\pm2.91\%$ (T₀), *C. carpio* from $78.00\pm1.73\%$ (T₂) to $86.33\pm2.40\%$ (T₀), *C. idella* from 73.00 ± 1.15 (T₀) to $83.00\pm1.15\%$ (T₁) and *B. gonionotus* from 70.33 ± 1.45 (T₀) to $78.67\pm0.67\%$ (T₁).

Among the species the highest survival rate was observed as $86.33\pm2.40\%$ in *C. carpio* and the lowest was in *C. cirrhosus* as $67.50\pm1.44\%$.

The lowest mean survival rate was found as $76.33\pm1.74\%$ g (T₂) and the highest final weight as $79.52\pm2.20\%$ (T₀) for all the species.

Significant difference was found among the treatments for the mean values of survival rate of all the species except *H. molitrix* and *L. rohita*.

Yield (Kg/ha/6 months)

The yield (Kg/ha/6 months) varied with *H. molitrix* from 636.00 ± 3.79 (T₃) to 1250.00 ± 14.43 (T₀), *C. catla* from 186.67 ± 2.03 (T₃) to 426.33 ± 8.41 (T₀), *L. rohita* from 479.33 ± 924 (T₂) to 992.00 ± 4.16 (T₀), *C. cirrhosus* from 269.67 ± 1.76 (T₃) to 555.00 ± 7.81 (T₀), *C. carpio* from 226.00 ± 2.65 (T₃) to 385.00 ± 6.03 (T₀), *C. idella* from 527.00 ± 4.16 (T₃) to 603.67 ± 71.64 (T₁) and *B. gonionotus* from 214.33 ± 4.63 (T₃) to 375.67 ± 41.34 (T₁).

Among the different species the highest yield was observed as 1250.00 ± 14.43 kg/ha/6 months in *H. molitrix* and the lowest was in *C. catla* as 186.67 ± 2.03 kg/ha/6 months.

The lowest total yield (Kg/ha/6 months) was found as 2541.00 ± 0.67 (T₃) and the highest total yield was found as 4403.51 ± 0.88 (T₀). Significant difference was found among the treatments.

Significant difference was found among the treatments for the mean values of yield for all the species.

Treatments Species	Το	T ₁	T ₂	Тз
H. molitrix	1.29±0.41ª	1.14±0.31ª	0.96±0.23 ^b	0.94±0.21 ^b
C. catla	1.31±0.24ª	1.14±0.32ª	0.97±0.26 ^b	0.96±0.24 ^b
L. rohita	1.21±0.38ª	0.95±0.26 ^b	0.84±0.18 ^b	0.84±0.16 ^b
C. cirrhosus	1.24±0.38ª	1.05±0.25 ^{ab}	0.88±0.15 ^b	0.87±0.15 ^b
C. carpio	1.20±0.34ª	1.09±0.28 ^{ab}	0.96±0.22 ^b	0.93±0.22 ^b
C. idella	1.03±0.23 ^b	1.31±0.31ª	1.14±0.33 ^b	1.10±0.32 ^b
B. gonionotus	0.93±0.21 ^{ab}	1.05±0.25ª	0.86±0.28 ^b	0.85±0.21 ^b
All species	1.17±0.05ª	1.10±0.04 ^{ab}	0.94±0.04 ^b	0.93±0.03 ^b

Table-4.27: Variations in the mean SGR (%, bwd⁻¹) of fishes under different treatments during the study period

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-4.28: Variations in the mean weight gain (g/month) of fishes under different treatments during the study period.

Treatments Species	Τo	T ₁	T2	T₃
H. molitrix	95.78±5.56ª	70.94±4.17 ^b	47.83±3.96°	45.72±4.66°
C. catla	102.45±14.41ª	72.50±5.46 ^b	50.06±5.24 ^b	49.06±5.28 ^b
L. rohita	74.17±6.55ª	43.50±5.75 ^b	33.39±3.30 ^b	33.78±2.70 ^b
C. cirrhosus	74.33±5.59ª	50.78±3.36 ^b	34.94±2.97°	33.78±3.68°
C. carpio	80.61±4.99ª	64.50±4.26 ^b	48.33±3.60°	46.00±4.51°
C. idella	82.39±6.10 ^{ab}	95.17±6.99ª	73.44±5.57 ^b	72.84±5.13 ^b
B. gonionotus	26.72±2.49ª	34.95±4.53ª	15.22±2.78 ^b	15.11±1.95 ^b
All species	76.64±9.24ª	61.76±7.71 ^b	43.32±6.83°	42.33±6.71°

Treatments Species	Τo	Τı	T2	Тз
H. molitrix	636.67±5.61ª	487.67±30.21 ^b	349.00±4.04°	336.33±2.03°
C. catla	678.67±12.57ª	499.00±26.10 ^b	364.33±2.91°	358.33±4.41°
L. rohita	502.00±2.08ª	318.00±16.29 ^b	257.33±5.04°	259.67±3.48°
C. cirrhosus	500.00±7.23ª	358.67±36.86 ^b	263.67±6.06°	256.67±1.76°
C. carpio	546.67±8.69ª	450.00±37.63 ^b	353.00±4.73°	339.00±3.79°
C. idella	575.00±68.46 ^b	620.33±18.02ª	505.67±5.78 ^b	502.00±4.16 ^b
B. gonionotus	190.33±20.96 ^b	229.67±2.67ª	116.33±4.33°	115.67±2.60°
All species	3629.34±19.52ª	2963.34±21.20 ^b	2209.33±14.25°	2167.67±17.31°

Table-4.29: Variations in the mean final weight (g) of fishes under different treatments during the study period.

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-4.30: Variations in the mean survival rate (%) of fishes under different treatments during the study period.

Treatments				
	Τo	T₁	T ₂	T₃
Species				
H. molitrix	84.33±2.03ª	82.67±2.60ª	81.33±1.20ª	82.67±1.20ª
C. catla	81.00±1.73ª	79.33±2.33ª	75.33±1.59 ^b	75.33±2.03 ^{ab}
L. rohita	81.33±0.88ª	79.33±2.33ª	78.00±1.73ª	77.67±1.45ª
C. cirrhosus	80.33±2.91ª	74.50±3.91 ^{ab}	67.50±1.44 ^b	68.50±0.76 ^b
C. carpio	86.33±2.40ª	79.00±3.06 ^b	78.00±1.73 ^b	78.33±0.88 ^b
C. idella	73.00±1.15 ^b	83.00±1.15ª	80.00±6.56ª	81.33±1.76ª
B. gonionotus	70.33±1.45 ^b	78.67±0.67ª	74.17±3.63ª	76.33±1.45ª
All species	79.52±2.20ª	79.50±1.07ª	76.33±1.74ª	77.17±1.75ª

Treatments Species	T₀	Τı	T ₂	T₃
H. molitrix	1250.00±14.43ª	953.33±59.27 ^b	663.67±6.12°	636.00±3.79°
	(28.39%)	(25.94%)	(25.59%)	(25.03%)
C. catla	426.33±8.41ª	295.67±15.38 ^b	190.00±2.31°	186.67±2.03°
	(9.68%)	(10.86%)	(7.33%)	(7.35%)
L. rohita	992.00±4.16ª	628.33±32.31 ^b	479.33±9.24°	481.33±6.39°
	(20.26%)	(17.10%)	(18.48%)	(18.94%)
C. cirrhosus	555.00±7.81ª	398.33±40.92⁵	278.67±4.98°	269.67±1.76°
	(12.60%)	(10.84%)	(10.74%)	(10.61%)
C. carpio	385.00±6.03ª	317.00±26.66 ^b	235.33±2.96°	226.00±2.65°
	(8.74%)	(8.63%)	(9.07%)	(8.89%)
C. idella	566.10±23.36 ^b	603.67±71.64ª	531.00±6.11 ^b	527.00±4.16 ^b
	(12.86%)	(16.42%)	(20.47%)	(20.74%)
B. gonionotus	329.08±5.37 ^b	375.67±41.34ª	215.67±8.09°	214.33±4.63°
	(7.47%)	(10.22%)	(8.32%)	(8.43%)
Total yield (Kg/ha/6 months)	4403.51±0.88ª	3675.33±0.58 ^b	2593.67±0.11°	2541.00±0.67 ^d
Total yield (Kg/ha/yr)	8807.02±0.69ª	7350.66±0.46 ^b	5187.34±0.23°	5082.00±0.41 ^d

Table-4.31: Variations in the mean yield (kg/ha) of fishes under different treatments during the study period.

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Values in the parentheses means % contribution by individual species to total fish yield under respective treatment.



Fig. 4.2: Variations in the mean values of different growth parameters under different treatments during the study period: a; SGR (%, bwd⁻¹), b; weight gain (g/month), c; final weight (g) and d; survival rate (%)



Fig. 4.3: Mean values of individual fish yield under different treatments



Fig. 4.4: Mean values of total fish yield under different treatments





Plate 4.4: Harvesting of fishes.
4.3.3: Economics

The economics of fish farming under different treatments are presented in table 4.32 and fig. 4.5.

Total cost significantly varied from 123430.50 ± 0.00 (T₁, T₂, T₃) to 235930.50 ± 0.00 BDT/ha/6 months (T₀). Gross benefit significantly varied from 235068.40 ± 1965.31 (T₃) to 418376.85 ± 5125.59 Tk/ha/6 months (T₀). Net benefit significantly varied from 111639.90 ± 2056.87 (T₃) to 206744.85 ± 3221.73 Tk/ha/6 months (T₁). The Net profit margin significantly varied from 77.21 ± 2.40 (T₀) to $167.20\pm18.77\%$ (T₁). The CBR significantly varied from 0.77 ± 0.02 (T₀) to 1.67 ± 0.18 (T₁).

Treatments				
	Τo	T ₁	T ₂	T₃
Components				
Lease value (Tk.)	40000.00±0.00ª (16.95%)	40000.00±0.00ª (32.41%)	40000.00±0.00ª (32.41%)	40000.00±0.00ª (32.41%)
Pond preparation (Tk.)	10550.00±0.00ª (4.47%)	10550.00±0.00ª (8.55%)	10550.00±0.00ª (8.55%)	10550.00±0.00ª (8.55%)
Fertilizer (Tk.)	25400.00±0.00ª (10.77%)	25400.00±0.00ª (20.58%)	25400.00±0.00ª (20.58%)	25400.00±0.00ª (20.58%)
Fish seed (Tk.)	32480.50±0.00ª (13.77%)	32480.50±0.00ª (26.31%)	32480.50±0.00ª (26.31%)	32480.50±0.00ª (26.31%)
Feed (Tk.)	112500.00±0.00ª (47.68%)	0.00±0.00 ^b (0.00%)	0.00±0.00 ^b (0.00%)	0.00±0.00 ^b (0.00%)
Harvesting cost (Tk.)	15000.00±0.00ª (6.36%)	15000.00±0.00ª (12.15%)	15000.00±0.00ª (12.15%)	15000.00±0.00ª (12.15%)
Total cost (Tk.)	235930.50±0.00ª	123430.50±0.00 ^b	123430.50±0.00 ^b	123430.50±0.00 ^b
Gross benefit (Tk.)	418376.85±5125.59ª	330175.35±2155.32 ^b	239526.95±2335.84°	235068.40±1965.31°
Net benefit (Tk.)	182446.35±3265.00 ^b	206744.85±3221.73ª	116096.45±3554.84°	111639.90±2056.87°
Net profit margin (%)	77.21±2.40 ^b	167.20±18.77ª	94.38±2.91 ^{ab}	90.25±1.73 ^{ab}
CBR	0.77±0.02 ^b	1.67±0.18ª	0.94±0.03 ^{ab}	0.90±0.02 ^{ab}

Table 4.32: Economics of fish farming under different treatments during study period
--

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05) % of total cost in parentheses







Fig. 4.5: Economics of fish farming under different treatments: a; total cost (BDT), b; net benefit (BDT) and c; CBR

4.4: Discussion

4.4.1: Water quality parameters

Water quality parameters play an important role on the growth and production of fish and other aquatic organisms. The suitable water quality parameters are prerequisites for a healthy aquatic environment and for the production of sufficient fish food organisms. The primary productivity of a water body depends on the physical, chemical and other factors of the environment (Rahman, 1992).

Fortnightly variations

Water temperature

The temperature of water has extremely important ecological consequences. Temperature exerts a major influence on aquatic organisms with respect to selection and level of activity of the organisms. In general, increasing water temperature results in greater biological activity and more rapid growth. All aquatic organisms have preferred temperature in which they can survive and reproduce optimally. Temperature is a regulator of the solubility of gases and minerals (solids). The solubility of important gases, such as oxygen and carbon dioxide increases as temperature decreases (Jhingran, 1975). In the present study water temperature varied from 29.58±0.03 to 32.23±0.08°c. Lower temperature (29.58±0.03) with treatment T_0 at 1st fortnight (April 2011) might be due to spring season when air temperature was relatively low (Appendix-1). Higher water temperature (32.23±0.08°C) with treatment T₀ at 6th fortnight (June, 2011) might be due to bright sunlight during study period (Appendix-1). The water temperature of the studied ponds increased towards first three months due to summer season and gradually decreased at the end of the study which was influenced by advent of winter season. Ferdoushi et al. (2008) recorded temperature range 23.6°C to 35.5°C, while working on carp polyculture for a period of five months (July to November) in Azolla fed pond. No significant difference was found among the treatments for the fortnightly values of temperature. Rahman et al. (1982) found pond water temperature varied from 26.06°C to 31.97°C that was almost similar to those obtained in the present study. Wahab *et al.* (1995) found that the suitable temperature range in BAU campus, Mymensingh was 27.2°C to 32.4°C. Islam *et al.* (1997) and Sharmin and Jewel (2008) also reported similar findings while working on carp polyculture.

Water transparency

Transparency is a measure of how clear the water is. It is important, because aquatic plants need sunlight for photosynthesis. The clearer the water, the deeper sunlight will penetrate. The water transparency acts as an index of productivity of a water body. The water transparency showed variation with sampling dates and it ranged from 30.00 ± 0.58 cm with treatment T₂ at 7th fortnight (July) to 35.00 ± 0.00 cm with treatment T₁ at 1st fortnight (April). This higher value might be due to the available nutrient concentration resulting from basal dose of both organic and inorganic fertilization with the treatment.

In the present study, the lower value of the transparency in the month of July, 2011 might be due to higher turbidity causes by heavy rain fall in that month (Appendix-2). Significant variation was found at 1st, 10th and 12th fortnights among the treatments, these variations might be due to rainfall and / or nutrient concentrations. This observation is similar to the findings of Chowdhury and Mamun (2006) while working on the water quality analysis in two ponds in Khulna. According to Boyd (1990) secchi disc visibility about 30 to 45 cm means the water body productive, if it is not newly constructed or turbid due to rainfall or burrowing by fish or other organisms.

Dissolved oxygen (DO)

Dissolved oxygen (DO) content is probably the only most important water quality parameters in aquaculture. Prolonged exposure to low concentration of DO can be harmful for aquatic life (BAFRU, 1990). In the present study, DO content of water was observed between 4.90 ± 0.06 mg/l with treatment T₂ at 6th fortnight (June) to 6.17 ± 0.12 mg/l with the treatment T₁ at 10th fortnight (August). The lower value during month of June was probably due to high temperature, respiration and decomposition of organic matter. The maximum DO value was 6.17 ± 0.12 mg/l (T₁ at 10th fortnight) due to advent of winter which agreed with Saha *et al.* (1971) while working on seasonal variations in water quality parameters in fish pond. Wahab *et al.* (1995) worked with carp polyculture in ponds and reported that DO concentration ranging from 5.0-7.0 mg/l is within the good productive range. DoF (1998) reported that the suitable DO of water body for fish culture would be 5-8 mg/l. So the result of the present study is more or less similar to the findings of the above authors. Significant variations were found at 6th, 9th, 11th and 12th fortnights among the treatments. Fortnightly variations might be due to fluctuation of temperature (Appendix-2) and treatments variation in the same fortnight might be decomposition of supplied feeds and weeds. These arguments are supported by Majhi *et al.* (2006) who worked with growth performance and production of organically cultured grass carp.

Hydrogen ion concentration (pH)

pH is defined as the relative degree of acidity or alkalinity in an environment. pH of water is one of the most important factor which has intense effect on the productivity of water body. There were no wide variations in pH values in the investigated ponds. The values of pH in the water ranged from 7.03±0.03 (T₃ at 6th forthright) to 7.87±0.09 (T₀ at 12th forthright). During the experimental period pH of the most ponds were slightly alkaline, which indicated a good pH condition for fish culture. Similar findings were found by Islam *et al.* (1997). Significant variation was found at 12th fortnight among the treatments. According to Bergins (1949) in alkaline water pH value increases with the increase of carbonate because in alkaline water by the increased consumption of CO₂, cause the breaking of bicarbonate into free CO₂ and carbonate (2HCO₃ = CO₂ + CO₃-² + H₂O). However the pH value in alkaline condition in pond water was supposed to be helpful for proper growth and development of fishes and aquatic organisms (Nikolsky, 1963; Huet, 1972, Jhingran, 1975).

Alkalinity

Alkalinity refers to the sum of the carbonate and bi-carbonate alkalinities. In case of aquatic productivity, the total alkalinity is also an important factor. Total alkalinity was found to range from 109.22 ± 1.33 (T₃ at 1st forthright) to 120.00 ± 1.15 mg/l (T₀ at 9th forthright). Welch (1952) and Huchinson (1957)

reported that decrease in alkalinity might probably be caused by the break down of bicarbonate due to the photosynthetic activity of plants. The high value during the spring was possibly due to the low CO₂, low rainfall and evaporation of water, which causes an increase in alkalinity. Significant variations were found at 8th and 9th fortnights (July and August), among the treatments. It might be due to heavy rainfall (Appendix-1). Production of free CO₂ enhanced by high temperature agreed with Michael (1968) and Verma (1969). Alikhuni (1957) reported that total alkalinity more than 100 mg/l should be present in high productive water bodies. Haque *et al.* (2005) found the average total alkalinity values above 100 mg/l in their experiments while working on production and economic return of carp polyculture in farmers ponds.

Free carbon dioxide (CO₂)

The mean values of free carbon dioxide varied from 2.49 ± 0.07 (T₁ at 4th forthright) to 3.46 ± 0.03 mg/l (T₃ at 12th forthright). The lower value of free CO₂ content was obtained at the starting period of the experiment (May, 2011) when temperature and rainfall were relatively low (Appendix-1). Higher values of free CO₂ accelerated the rate of decomposition of organic matter by microbes, decrease of photosynthetic activity and high rate of respiration by benthic biota and microorganisms (Chowdhury *et al.*, 1992). Significant variation was found at 10th fortnight among the treatments. Higher value of CO₂ with treatment T₀ might be due to comparatively low intake of feed by fishes in rainy season. As a result wastage of supplied feed increased and decomposition took place at higher rate which ultimately increased the free CO₂ level with that treatment. The fluctuation of CO₂ value might be due to alteration in the rate of photosynthesis in ponds and oxygen consumption by fish and other decomposer microorganisms (Khan *et al.*, 2009).

Mean variation

Water temperature

In the present study, the mean value of water temperature varied from 31.19 ± 0.20 (T₃) to 31.29 ± 0.24 °C (T₀). The water temperature measured in different treatments throughout the experimental period were within the

acceptable range for fish culture. PRISM-Bangladesh made a study on weedbased carp polyculture and recorded water temperature as 28.1 to 32°C (DWRP, 1996 and 1997). Ferdoushi *et al.* (2008) reported water temperature as 29.30±2.63°C to 29.60±2.47°C while working on aquatic floating macrophytes (Lemna and Azolla) in fish pond. Dewan *et al.* (1991) reported a temperature range of 30.20 to 34.0°C, while Wahab *et al.* (1996) recorded from 28.5 to 31.3°C in their experiment with carp polyculture. Water temperature of 25 to 32⁰C is considered suitable for fish culture (Boyd and Zimmermann, 2000).

Water transparency

The mean value of water transparency in this study varied from 32.28 ± 0.32 (T₂) to 32.83 ± 0.40 cm (T₁). Banforth (1958) reported that transparency indicated the presence and absence of food particles and productivity of a water body, which was influenced by the suspended materials, silt and microorganisms. Rahman *et al.* (2006) found mean values of transparency ranging from 20.71 to 26.37° C while working with carp polyculture in seasonal ponds. Present finding has similarity with the findings of Azim *et al.* (1995) who found the transparency value as 36.2cm in weed based carp polyculture pond. Boyd (1982) also recommended the transparency between 30 to 45 cm as appropriate for fish culture.

Dissolved oxygen (DO)

The mean value of dissolved oxygen varied from 5.33 ± 0.09 (T₃) to 5.51 ± 0.11 (T₁) mg/l. More or less similar finding (4.15 mg/l) was made by Thy *et al.* (2004) in effluent plus water spinach treated pond. Chowdhury *et al.* (2008) reported the mean values of DO ranging from 6.29 to 6.64, while working with duckweed fed Nile tilapia culture in earthen ponds. Azim and Wahab (2003) recorded similar DO value that ranging from 4.1 to 5.25 mg/l in duckweed based carp polyculture pond. The suitable range of dissolved oxygen is 5-8 mg/l for fish culture (Boyd, 1998).

Hydrogen ion concentration (pH)

The mean value of pH varied from 7.18 ± 0.03 (T₃) to 7.38 ± 0.05 (T₁). Ferdoushi *et al.* (2008) recorded pH value as 7.53 ± 0.034 in Azolla fed carp polyculture

pond. Dewan *et al.* (1991) recorded the mean value of water pH ranged from 6.60-8.60 which is similar to the present study. Majhi *et al.* (2006) reported that the variation in pH was 7.5-8.2 of a Azolla fed pond in Meghalaya of North Eastern India. Hossain *et al.* (1997) also obtained a pH range of 6.7 to 8.3 in fish ponds. According to Swingle (1967), pH of 6.5 to 9 is suitable for fish culture. Large changes in pH can cause stress, poor growth and even death of the farmed animals (Wurts and Durborow, 1992). Kohinoor *et al.* (1998) also recorded the mean values of pH as 7.18±2.40 in ponds. In the present study the alkaline pH range in all treatments indicated good pH condition for biological production and fish culture.

Alkalinity

The recorded mean total alkalinity varied from 113.28 ± 0.85 (T₁) to 114.36 ± 0.72 (T₀) mg/l. Total alkalinity values depending upon the location, season, plankton population and nature of bottom deposits (Jhingran, 1991). Mitra *et al.* (1978) found alkalinity values results ranged from 100.00 to 162.00 mg/l in carp polyculture ponds which was relatively higher than the present study. It might be due to low dissolved oxygen content and more production of free CO₂ that enhanced by increasing of fish biomass which strongly supported by Michael (1968) and Verma (1969) while working with water quality parameters in fish ponds and tropical impoundments. Boyd (1982) advocated that the total alkalinity should be more than 20 ppm in fertilized ponds as production increases with the increase in total alkalinity. The suitable range of alkalinity in fish culture is found as 50 to 300 mg/l (Buttner, 1993). Mairs (1996) also stated that the total alkalinity of 40.0 mg/l or more to be productive than water bodies with lower alkalinity. So, the alkalinity values of the study ponds were found within the suitable range for fish farming.

Free carbon-dioxide

Results from the study indicated that the value of free CO₂ varied from 2.82±0.07 (T₁) to 3.03 ± 0.07 mg/l (T₀). Comparatively higher value of CO₂ recorded in ponds used with conventional fish feed pond (T₀) might be due to higher fish biomass and decomposition of unused fish feed. Lower value of CO₂ recorded in Azolla fed pond (T₁) might be due to frequent supply of oxygen

by aquatic macrophytes. This argument is strongly supported by Chowdhury *et al.* (2008) when they used duckweed as supplementary feed. The suitable range of free CO_2 for fish culture is ranged from 1.0 to 10.0 mg/l (Boyd, 1998). Wurts and Durbrow (1992) reported that the desirable range of CO_2 1-2 ppm for pond aquaculture.

4.4.2: Growth of fishes

Monthly variations

Specific growth rate (%, bwd⁻¹)

The specific growth rate (%, bwd⁻¹) significantly varied with *H. molitrix* from 0.30±0.01 (T₃ at 6th month) to 3.13±0.03 (T₀ at 1st month), *C. catla* from 0.28±0.01 (T₂ and T₃ at 6th month) to 2.49±0.09 (T₁ 1st month), *L. rohita* from 0.23±0.04 (T₁ at 6th month) to 2.87±0.03 (T₀ at 1st month), *C. cirrhosus* from 0.31±0.05 (T₃ at 6th month) to 2.91±0.04 (T₀ 1st month), *C. carpio* from 0.30±0.03 (T₃ at 6th month) to 2.70±0.03 (T₃ at 3rd month), *C. idella* from 0.34±0.01 (T₀ 6th month) to 3.15±0.06 (T₁ at 1st month) and *B. gonionotus* from 0.14±0.01 (T₂ at 6th month) to 3.01±0.06 (T₁ at 1st month). Among the species, the highest SGR was observed in *C. idella* as 3.15±0.06 % bwd⁻¹ (T₁) and the lowest was in *B. gonionotus* as 0.14±0.01 %, bwd⁻¹ (T₂). The result indicates that *C. idella* grows very fast at early stage as compared to other carp species if fed with Azolla. Majhi *et al.* (2006) found that grass carp fed with Azolla showed a higher growth performance at early stage. Results indicated that *B. gonionotus* grows slowly at winter, fed with grass.

Significant variation in SGR values with the treatments was found in all months except the 2nd month (May) for *C. carpio*, 2nd month (May) and 3rd month (June) for *C. idella*, 4th month (July) 5th month (August) and 6th month (September) for *C. cirrhosus* and 4th month (July) for *B. gonionotus*.

The lowest combined SGR value was found as 0.30 ± 0.03 (T₂ at 6th month) and the highest combined SGR was found as 2.62 ± 0.17 (T₀ at 1st month). The highest SGR value was found in the month of April and the lowest value was found in the month of September for all the species. The lowest SGR value in the month of September might be due to the slow growth rate at mature stage of fishes. Slow growth due to slow metabolic activity of fish is found in lower water temperature (Boyd, 1998).

Monthly weight gain (g/month)

The weight gain (g/month) varied with *H. molitrix* from 28.67±0.67 g (T₃ at 6th month) to 109.67±2.03 g (T₀ at 4th month), C. catla from 28.33±1.20g (T₃ at 6th month) to 154.33±3.84 g (T₀ at 4th month), *L. rohita* from 19.33±1.45 g (T₂ at 6th month) to 90.67 \pm 1.20 g (T₀ at 3rd month), *C. cirrhosus* from 22.67 \pm 3.33 g (T₃ at 6^{th} month) to 90.00±0.58 g (T₀ at 3rd month), *C. carpio* from 29.67±0.33 (T₃ at 6th month) to 96.00±1.53 g (T₀ at 4th month), *C. idella* from 53.33±0.88 (T₂ at 6th month) to 111.33±4.37 g (T₁ at 3rd month) and *B. gonionotus* from 4.67±0.33 (T₂ at 6th month) to 50.00±0.58 g (T₁ at 3rd month). Among the different species, the highest monthly weight gain observed in *C. catla* as 154.33±3.84 g (T₀ at 4th month) might be due to higher nutrient concentration resulting from regular supply of feed and fertilizer. Jena et al. (2002b) mentioned that C. catla showed a better growth performance than other carp species. Lowest value in B. gonionotus as 4.67±0.33 g (T₂ at 6th month) was due to slow growth pattern of this species. Significant variation was found in all months for all the species except the 2nd month (May) and 3rd month (June) for *C. idella*. The lowest combined weight gain value was found as 29.10 \pm 5.66 (T₂ at 6th month) and the highest combined weight gain was found as 94.52±13.75 (T₀ at 4th month). The highest weight gain was found in the month of July and the lowest value was found in the month of September for all the species. It might be due to higher metabolic rate at comparatively higher temperature. Comparatively higher mean monthly weight gain was observed at the middle period of the experiment, this might be due to influence of air temperature on water temperature resulting fast metabolic activity (Boyd, 1998).

Mean variations

Specific Growth Rate (%, bwd⁻¹)

The mean specific growth rate (SGR, %, bwd⁻¹) varied with *H. molitrix* from 0.94 \pm 0.21 (T₃) to 1.29 \pm 0.41 (T₀), *C. catla* from 0.96 \pm 0.24 g (T₃) to 1.31 \pm 0.24 (T₀), *L. rohita* from 0.84 \pm 0.16 (T₃) to 1.21 \pm 0.38 (T₀), *C. cirrhosus* from 0.87 \pm 0.15 (T₃) to 1.24 \pm 0.38 (T₀), *C. carpio* from 0.93 \pm 0.22 (T₃) to 1.20 \pm 0.34

Chapter Four Experiment-2

(T₀), *C. idella* from 1.03±0.23 (T₀) to 1.31±0.31 (T₁) and *B. gonionotus* from 0.85±0.21 (T₃) to 1.05±0.25 (T₁). Jasmine, et al. (2011) found the mean SGR of silver as 1.26 to 1.40 %, bwd⁻¹ and catla as 1.19 to 1.21 %, bwd⁻¹ in polyculture pond. The mean SGR (%, bwd⁻¹) of catla, rohu and mrigal in different treatments under polyculture system were recorded as 1.09 to 1.12, 1.13 to 1.14 and 1.10 to 1.12 (Rahman et al. 2007). Kohinoor et al. (1999) observed the SGR value of Thai sharpunti to be between 1.33 and 1.35 %, bwd⁻¹ in polyculture with carps using low-cost feed. However, among the species comparively highest SGR (1.31 \pm 0.31% bwd⁻¹) was obtained for *C. idella* in treatment T₁, where Azolla was supplied as feed and the lowest SGR (0.84 \pm 0.16) for *L. rohita* in treatment T₃ where banana leaf was supplied. It is clear that grass carp grows well with Azolla fed pond. Majhi et al. (2006) recorded SGR value of *C. idella* as 1.65 %, bwd⁻¹ in Azolla fed fish pond. The lowest SGR value found as 0.93±0.03 (T₃) might be due to intake of banana leaf only by the macrophytophagus species and the highest combined SGR as 1.17±0.05 (T₀) might be due to combined effect of conventional feed and natural feed. Ferdoushi et al. (2008) reported that supply of Azolla with conventional feed showed better growth performance than that of only used conventional feed in carp polyculture system.

Weight gain (g/month)

Mean weight gain (g/month) significantly varied with *H. molitrix* from 45.72 ± 4.66 g (T₃) to 95.78 ± 5.56 g (T₀), *C. catla* from 49.06 ± 5.28 g (T₃) to 102.45 ± 14.41 g (T₀), *L. rohita* from 33.39 ± 3.30 g (T₂) to 74.17 ± 6.55 g (T₀), *C. cirrhosus* from 33.78 ± 3.68 g (T₃) to 74.33 ± 5.59 g (T₀), *C. carpio* from 46.00 ± 4.51 g (T₃) to 80.61 ± 4.99 g (T₀), *C. idella* from 72.84 ± 5.13 g (T₃) to 95.17 ± 6.99 g (T₁) and *B. gonionotus* from 15.11 ± 1.95 g (T₃) to 34.95 ± 4.53 g (T₁). Azad *et al.* (2004) reported the weight gain of *H. molitrix* as 72.87 g and *C. cirrhosus* as 70.42 g while working with carp polyculture using low cost inputs. Among the species the highest weight gain was observed as 102.45 ± 14.41 g in *C. catla*, which might be due to stocking of comparatively larger sized fingerlings. The lowest weight gain was found in *B. gonionotus* as 15.11 ± 1.95 g which might be due to slow growth rate. Azim *et al.* (2004) worked on feeding relation in a carp

polyculture system and they found that weight of *C. catla* was moderate and weight gain *B. gonionotus* was lowest. The lowest combined mean weight gain was found as 42.33 ± 6.71 g (T₃) and the highest combined weight gain as 76.64 ± 9.24 g (T₀) for all the species. The lowest combined weight gain (T₃) might be due to apply only non-conventional feed (banana leaf) which is consumed only by the macrophytophagous species. On the other hand, highest combined weight gain (T₀) might be due to the use of conventional feed and naturally produced feed, which is favourable for better growth of all species.

Final Weight (g)

The mean final weight (g) varied with *H. molitrix* from 336.33 ± 2.03 (T₃) to 636.67±5.61 g (T₀), C. catla from 358.33±4.41 (T₃) to 678.67±12.57 g (T₀), L. rohita from 257.33±5.04 (T₂) to 502.00±2.08 g (T₀), C. cirrhosus from 256.67±1.76 (T₃) to 500.00±7.23 g (T₀), C. carpio from 339.00±3.79 (T₃) to 546.67±8.69 g (T₀), *C. idella* from 502.00±4.16 (T₃) to 620.33±18.02 g (T₁) and B. gonionotus from 115.67±2.60 (T₃) to 229.67±2.67 g (T₁). Rahman et al. (2006) reported the final weight of rui, catla, mrigal, silver carp and thai sharputi as 144.71g, 189.44g, 89.68g, 302.70g and 79.21g respectively in polyculture system over 90 days of rearing. The growth rate of *C. carpio* recorded by Das (2000) was 170.0 g and was 97.78 g by Azad et al. (2004) which were less than the final weight obtained in the present study. It might be due to the good water quality, regular supply of feed and Azolla, larger stocking size of fingerlings. Majhi, et al. (2006) reported the final weight of 270.34g in C. idella fed with Azolla in 150 days culture period at a stocking density of 0.75 fish/m² and initial stocking size of 23.50 \pm 5.0g. Among the species the highest final weight was observed as 678.67±12.57 g in C. catla and the lowest was in B. gonionotus as 115.67±2.60 g. The lowest mean final weight was found as 2167.67 \pm 17.31 g (T₃) and the highest final weight as 3629.34 \pm 19.52 g (T₀) for all the species. The reason behind the highest final weight might be due to proper utilization of both natural and supplementary feed.

Survival rate (%)

The survival rate (%) varied with *H. molitrix* from 81.33 ± 1.20 (T₂) to $84.33\pm2.03\%$ (T₀), *C. catla* from 75.33 ± 1.59 (T₂) to $81.00\pm1.73\%$ (T₀), *L. rohita* from 77.67 ± 1.45 (T₃) to $81.33\pm0.88\%$ (T₀), *C. cirrhosus* from 67.50 ± 1.44 (T₂) to $80.33\pm2.91\%$ (T₀), *C. carpio* from $78.00\pm1.73\%$ (T₂) to $86.33\pm2.40\%$ (T₀), *C. idella* from 73.00 ± 1.15 (T₀) to $83.00\pm1.15\%$ (T₁) and *B. gonionotus* from 70.33 ± 1.45 (T₀) to $78.67\pm0.67\%$ (T₁). Das *et al.* (1982) reported that the survival of rohu, catla, mrigal and silver carp were 88.64 - 90.18, 85.81 - 90.06, 84.93 - 91.36 and 92.35 - 96.96%, respectively in polyculture of indigenous and exotic fish species. Roy *et al.* (2002) reported the survival rate of grass carp, rohu, catla and mrigal as 76.6%, 87.8%, 84.0% and 88.6%, respectively in carp polyculture with SIS. Mostaque (1995) recorded the survival rates as 86% to 95% in a polyculture system in ponds. Khan *et al.* (2009) also recorded 83.2% survival rate for silver carp while working in carp polyculture system.

Among the species the highest survival rate was observed as $86.33\pm2.40\%$ in *C. carpio* and the lowest in *C. cirrhosus* as $67.50\pm1.44\%$. This might be due to high tolerance capability of *C. carpio* than other carp species. Azad *et al.* (2004) found the lowest survival rate for *C. cirrhosus* among carp species.

The lowest combined mean survival rate was found as $76.33\pm1.74\%$ g (T₂) and the highest survival rate as $79.52\pm2.20\%$ (T₀) for all the species which might be due to available supply of food for all species. Wahab *et al.* (1995) reported that the combined survival rate was 80% in polyculture system.

Yield (Kg/ha/6 months)

The yield (Kg/ha/6 months) varied with *H. molitrix* from 636.00 ± 3.79 (T₃) to 1250.00 ± 14.43 (T₀), *C. catla* from 186.67 ± 2.03 (T₃) to 426.33 ± 8.41 (T₀), *L. rohita* from 479.33 ± 924 (T₂) to 992.00 ± 4.16 (T₀), *C. cirrhosus* from 269.67 ± 1.76 (T₃) to 555.00 ± 7.81 (T₀), *C. carpio* from 226.00 ± 2.65 (T₃) to 385.00 ± 6.03 (T₀), *C. idella* from 527.00 ± 4.16 (T₃) to 603.67 ± 71.64 (T₁) and *B. gonionotus* from 214.33 ± 4.63 (T₃) to 375.67 ± 41.34 (T₁). Majhi *et al.* (2006) worked on effect of Azolla feeding on growth performance of grass carp and obtained total production of grass carp (*C. idella*) was 185.76 kg/1000m² in 150 days culture period. Roy *et al.* (2002) recorded the yield of catla as 698.4, followed by

L. rohita as 754.1, *C. cirrhosus* as 761.4 and *C. idella* as 346.5 kg/ha/7 months at a stocking density of 3167, 3167, 3166 and 500 individual/ha, respectively supplied with only rice brain.

Among the species the highest yield was observed as 1250.00 ± 14.43 kg/ha/6 months in *H. molitrix* and the lowest was in *C. catla* as 186.67 ± 2.03 kg/ha/6 months. This might be the result of higher stocking density of *H. molitrix* and competition for food with *C. catla*. A severe competition for food between planktivorous native carps and exotic carps has been observed by Dewan *et al.* (1991). The maximum total fish production as 4403.51 kg was obtained in T₀, where conventional feeds were supplied and lower production as 2541.00 kg/ha/6 months in T₃, where only banana leaf was supplied as feed. Moderate production was obtained in T₁ (Azolla fed) and comparatively lower production was obtained in T₂ (grass fed). However, significant difference in total fish production was found among the treatments. It is clear that use of conventional feed, Azolla, grass and banana leaf has direct effect on the total production in polyculture system.

Uddin et al. (1994) found a gross production of 3,415 kg/ha/yr from polyculture of carps with rajpunti. Islam et al. (1997) obtained a net yield of 2,966 kg/ha/7 months by culturing silver carp, common carp and tilapia in seasonal ponds through fertilization and supplementary feeding. Mahmud (1998) reported a total gross production of 1,713.4 kg/ha/120 days in a four species composite culture of major carps including Thai sharpunti through the application of fertilizer and supplementary feed comprising of rice bran (60%) and mustard oil cake (40%) daily at the rate of 3% of fish biomass and duckweed (Lemna minor) at the rate of 10% of the body weight of that sharpunti. Miaje (1999) reported the total production of fish from 2,934 to 3,318 kg/ha/4 months in polyculture of Indian major carps with Thai sharpunti fed supplementary diet containing 20.31% crude protein. Das et al. (1982) reported a total gross production of 2,102.50-4,361.43 kg/ha/year on polyculture of major carps with the application of fertilizer and supplementary feed consisting of rice bran (75%) and mustard oil cake (25%). The total productions of fish obtained from the present study are similar to those reported by Das et al. (1982) and Mahmud (1998), higher than those of Uddin et al. (1994), Islam et al. (1997) and nearly closer to Miaje (1999). Roy et al. (2003) obtained total yield as 2560

103

kg/ha/7 months in carp polyculture system which was found lower as compared to the present study. Azim and Wahab (2003) also recorded total yield as 2020 kg/ha/4 months in duckweed based system.

4.4.3: Economics

Economic viability is an important criterion for sustainability of any system. Economic analysis provides a basis not only for the decision making of the individual fish farmer, but also for the formulation of aquaculture policies (Shang, 1981). During study period total cost, gross benefit, net benefit and CBR varied from 123430.50 ± 0.00 BDT/ha/6 months (T₁, T₂, T₃) to 235930.50±0.00 BDT/ha/6 months (T₀), 235068.40±1965.31 (T₃) to 418376.85±5125.59 BDT/ha/6 months (T₀), 111639.90±2056.87 (T₃) to 206744.85±3221.73 BDT/ha/6 months (T₁) and 0.77±0.02 (T₀) to 1.67±0.18 (T₁), respectively.

Total cost was higher in Treatment T_0 which might be due to the feed cost (47.68% of total cost). Abdelghany and Ahmad (2002) worked on cost-benefit analysis in polyculture system and they reported total cost was BDT/ha 2,15,202.00 which was closer to the present study.

Gross benefit in Treatment T₁ was 330175.35 ± 2155.32 BDT/ha which was more or less similar to the findings of Abdelghany and Ahmad (2002) who worked on cost-benefit analysis in carp polyculture pond reported gross benefit from fish sale was BDT/ha 309524.3.0.

In case of net benefit, Grover *et al.* (2000) described the net benefit of the Azolla based carp polyculture in ponds was 310990 Tk/ha/year. Shamugasundaram and Balusamy (1993) stated benefit cost ratio as 1.88 which was similar to the present study. Khan *et al.* (2009) also mentioned BCR value of 1.22 which was lower than the findings of the present study, this might be due to high initial biomass of all the species and the higher survival rate of fishes in the present study.

Data on economics indicated that the treatment T_1 (Azolla fed pond) was more profitable than that of others.

Chapter Four Experiment-2

Overall findings indicated that in treatment T₁ (Azolla fed pond), the water quality parameters were found in suitable range for fish growth, fish production was moderately high and cost benefit ratio was found highest. Azolla, which grows in association with the blue green algae Anabaena azollae, is perhaps the most promising from the point of view of ease of cultivation productivity and nutritive value (Lumpkin and Plucknett 1982; Van-Hove and Lopez 1983). Azolla-fed fish pond provides with a complete and balanced diet for those fish that consume it directly, while the faces of Azolla feeding species are consumed directly by detritus feeders, or indirectly used as fertilizer, enhancing plankton and other food organisms, which can be utilized by remaining surface and column feeding fish species . Providing Azolla as supplementary feed in this system could minimize the food competition between native carps and macrophytes (Journey et al., 1990). Therefore, the present finding regarding the potentials of aquaculture with non-conventional feed specially with Azolla as a low cost farming are proven and agreed well by the aforementioned scientists.

4.5: Conclusion

Considering the water quality, fish production and economic viability it can be concluded that Azolla based carp polyculture is more profitable as well as low cost culture technology and potentially be used for poor fish farmer.

4.6: Recommendation

This study identified Azolla based fish farming as a potential low cost aquaculture. On the other hand the major limitation of this study was with using only one stocking density for all treatments. Therefore, possibilities of finding maximum fish production under Azolla based fish farming was not explored in terms of suitable stocking density. So, it is recommended that optimization of stocking density for Azolla based carp polyculture in ponds should be explored as further research step.

CHAPTER FIVE

Optimization of stocking density for Azolla based carp polyculture in pond

5.1: Introduction

Polyculture or composite culture is the system in which fast growing compatible species of different feeding habits are stocked in different proportions in the same ponds (Jhingran, 1975). The basic principles of the polyculture, species of different feeding habits are culture in the same pond to avoid food competition and best utilization of natural foods of different habits without any harm to each other. It is a fact that, polyculture may produce an expected result if the fish with different feeding habits are stocked in proper ratios and combinations (Halver, 1984). Selection of species plays an important role for any cultural practices. For better utilization of different strata and zones of a pond three or more species must be stocked. In our country, suitable or common combinations of fish for composite culture are rui (*Labeo rohita*), catla (*Catla catla*) and mrigal (*Cirrhinus cirrhosus*) (Miah *et al.*, 1997).

In order to obtain more production several exotic carps cultured combined with the local carps. These include silver carp (*Hypophthalmichthys molitrix*), grass carp (*Ctenopharyngodon idella*), common carp (*Cyprinus carpio*), raj punti (*Barbonymus gonionotus*) etc. In search of compatible fish species for polyculture, Wahab *et al.* (2002) observed that common carp (*Cyprinus carpio*) grew better than its counter-part native species mrigal and did not exert significant antagonistic impact on other native carp species in polyculture. Thai silver barb (*Barbodes gonionotus*) has become increasingly popular owing its bright silvery appearance and good taste. This fish is an omnivorous and reportedly grows well on macrophytes (Phaohorm, 1980; Rothuis *et al.*, 1998). It has been proven to be a suitable species for culturing in the seasonal ponds (Islam *et al.*, 1998; Kohinoor *et al.*, 1999) and rice fields (Haroon and Pittman, 1997; Vromant *et al.*, 2002).

Freshwater fishes dominate global aquaculture production (56.4%, 33.7 million tonnes), followed by molluscs (23.6%, 14.2 million tonnes), crustaceans (9.6%, 5.7 million tonnes), diadromous fishes (6.0%, 3.6 million tonnes), marine fishes (3.1%, 1.8 million tonnes) and other aquatic animals (1.4%, 814 300 tonnes). While feed is generally perceived to be a major constraint to aquaculture development, one-third of all farmed food fish production (20 million tonnes) is currently achieved without artificial feeding, as is the case for bivalves and filter-feeding carps. However, the percentage of non-fed species in world production has declined gradually from more than 50 percent in 1980 to the present level of 33.3 percent, reflecting the relatively faster body-growth rates achieved in the culture of fed species and increasing consumer demand for higher trophic-level species of fishes and crustaceans (FAO, 2012).

According to FAO (2008) estimates, about 31.7 million tonnes (46.1% of total global aquaculture production including aquatic plants) of fish and crustaceans are feed-dependent, either as farm-made aqua feeds or as industrially manufactured compound aquafeeds. In 2008, fed aquaculture contributed to 81.2% of global farmed fish and crustacean production of 38.8 million tonnes and 60.0 percent of global farmed aquatic animal production.

While more than 200 species of fish and crustaceans are currently believed to be fed on externally supplied feeds, just 8 species or species groups account for 62.2% of the total feed used. These are: grass carp, common carp, Nile tilapia, Indian major carps (catla and rohu), white leg shrimp, crucian carp, Atlantic salmon, and pangasiid catfishes. More than 67.7% of farmed fish production is contributed by freshwater fishes, including carps and other cyprinids, tilapias, catfishes and miscellaneous freshwater fishes (FAO, 2012).

Farm-made aqua feeds play an important role in the production of low-value freshwater fish species. More than 97% of carp feeds used by Indian farmers are farm-made aqua feeds (7.5 million tonnes in 2006/07), and they are the mainstay of feed inputs for low-value freshwater fishes in many other Asian and sub-Saharan countries (FAO, 2012). The main production systems for freshwater aquaculture in Bangladesh are extensive and semi-intensive pond

polyculture of Indian and Chinese major carps which account for 80% of the total freshwater production. The remaining 20% are mainly from catfish, tilapia, small indigenous species (SIS) of fish and rice fish farming (ADB, 2005).

The traditional model of carp polyculture is conceptually elegant, and a great deal is known about the nutritional value of supplementary inputs. However, to achieve the highest productivity from a carp pond still involves a high degree of art. High production with current techniques requires a delicate and precarious balancing act between fish density, feed, fertilizer inputs, potential competitors and the amount of dissolved oxygen in the pond (FAO, 2008).

Stocking density is an important parameter in fish culture operations, since it has direct effects on the growth and survival and hence on production (Backiel and Le Cren, 1978). It is an established fact that the growth rate progressively increases as the stocking density decreases and vice-versa. This is because of relatively less number of fish in a pond of similar size could get more space food and dissolved oxygen at the same time. The growth of fishes is dependent on population density (LeCren, 1965; Backiel and LeCren, 1978), Generally direct relationship exists between food abundance and growth rate as well as between population density of the species and its growth rate, where as population density of the species and its growth rate tend to be inversely related (LeCren, 1965). However, there may be no relationship between food abundance and growth rate, when a space limiting effect operates on the population. Higher stocking density may cause crowding effects and reduction of growth rate. Size hierarchies within the fish population as a result, smaller fish 'is inhibited from feeding satisfactorily because of the presence of the larger ones. In many fish culture practices where the fish are confined in a restricted space, this size dominance in feeding is often of considerable significance (Brett, 1979).

One of the major problems faced by rapidly growing aquaculture is the availability of fish feed, since feed is the largest operating cost of semi-intensive fish farming. Feeding cost often constitutes more than 50% of the total cost of production in intensified culture system (Sehagal and Toor, 1991 and De Silva,

Chapter Five Experiment-3

1992). The common supplementary feeds in Bangladesh are rice bran, wheat bran, oil cake and some other agricultural wastes. As a resource constrained country, there is a severe competition for these agricultural wastes especially between raising livestock and fuel. Besides, the increase in cost and demand of feed protein from these conventional resources necessitates fish culturists of the developing countries to incorporate cheap and locally available ingredients in fish feeds. Therefore, the agua farmers have been exploring the possibilities of utilizing alternative sources having high food value like plant leaves and aquatic vascular plants for producing the much required animal protein at low cost (Lakshmanan et al., 1967). Azolla, which grows in association with the blue green algae Anabaena azollae, is perhaps the most promising from the point of view of ease of cultivation, productivity and nutritive value (Lumpkin and Plucknett, 1982; Van-Hove and Lopez, 1983). Fish require diets relatively higher in protein than those of commercially cultured animals. As protein represents the most expensive component in a formulated diet, it is of considerable practical importance to determine the optimum level that will support maximum growth and survival. Azolla contains 20- 25.5% protein, 3.1% fat, 34.9% carbohydrate, 8.5-11.7% cellulose and essential aminoacids. Biochemical and physiological changes in organisms influence the activities of some enzymes and metabolic levels of some tissues (Murty, 1986). It appears to be suitable for a wide range of fish species and may also be applied in the polyculture systems. It can be fed fresh as a sole feed or as an ingredient in combination with other components in artificial feed. A Azolla-fed fish pond provides with a complete and balanced diet for those fish that consume it directly, while the faeces of Azolla feeding species are consumed directly by detritus feeders, or indirectly used as fertilizer, enhancing plankton and other food organisms, which can be utilized by remaining surface and column feeding fish species . Providing Azolla as supplementary feed in this system could minimize the food competition between native carps and macrophytes (Journey et al., 1990).

It would be one of the cheapest ways in organic fish farming to increase the fish production from the aquatic habitat. This would also pave way for producing organic fish and narrow the wide gap between the production and demand of organic fish in the country. Above all, organic fish production would also increase, the farm income and living standard of low income farmers of the region.

Quantitative data on this issue are thus still very scarce to make reliable predictions of optimization of stocking density for Azolla based carp polyculture in pond. So far little or no experimental evidence exists for optimization of stocking density for Azolla based carp polyculture in pond polyculture system in Bangladesh or elsewhere in the region. Therefore, the present study aimed at evaluating the production and economics of Azolla based carp polyculture under different treatments of stocking densities.

5.2: Specific objectives

The specific objectives of this study were as follows-

- to monitor the water quality parameters (in terms of temperature, transparency, dissolved oxygen, pH, alkalinity and CO₂) in Azolla fed carp polyculture ponds under different treatments of stocking densities;
- to evaluate the growth performance of fishes under different treatments of stocking densities;
- to evaluate the economics of fish production under different treatments of stocking densities; and
- finally, to recommend suitable stocking density for Azolla based carp polyculture in ponds.

5.3: Materials and methods

5.3.1: Duration and location of the study

The study was conducted for a period of six months from April 2012 to September 2012 at Alampur village of Sadar Upazila under Kushtia district, Bangladesh (Map 5.1).





5.3.2: Description of the study ponds

A total of 9 ponds (average water area of 0.18ha and depth of 1.9 m) were selected for the present study. All the ponds were rain-fed and well exposed to sunlight (Plate 5.1).

5.3.3: Experimental design

The experiment was designed under Randomized Completely Block Design (RCBD) with three treatments (T_1 , T_2 and T_3) of carp stocking densities each with three replications. The treatment assignment was as follows-

- T₁ : 40 fish/decimal (9880 fish/ha)
- T₂ : 45 fish/decimal (11115 fish/ha)
- T₃ : 50 fish/decimal (12350 fish/ha)

Stocking ratio/combination of fish species were same for all the treatments (Table 5.1).

5.3.4: Pond management

Aquatic weeds were removed from all the ponds manually. Predatory fish and other unwanted species were removed through repeated netting. All the ponds were facilitated well with Azolla bank.

Liming was done at a rate of 250 kg/ha before 7 days of fertilization. All the ponds were fertilized with cowdung (1500 kg/ha), urea (60 kg/ha) and Triple Super Phosphate (TSP) (60 kg/ha) as basal dose.

One tenth area of the research ponds were used as Azolla bank according to Grover *et al.* (2000). Azolla bank was prepared by bamboo fencing (locally called 'Bana'). In Azolla bank, compost manure was deposited (2470 kg/ha). Azolla seeds were introduced in Azolla bank at the rate of 1000 kg/ha for available supply of Azolla at the culture period.

Fishes were stocked in all ponds after five days of basal fertilization. All the ponds were stocked with seven species of fishes (mean initial weight of *Hypophthalmichthyes molitrix, Catla catla, Labeo rohita, Cirrhinus cirrhosus, Cyprinus carpio, Ctenopharyngodon idella* and *Barbonymus gonionotus* were 60, 65, 58, 52, 61, 70 and 22 g, respectively).

Fish seeds were collected from local government fish farm. Stocking of fish seeds were done at early morning. Stocking density and stocking ratio/combination are shown in Table 5.1 (Grover *et. al.*, 2000).

	Treatments→	Т	1	Т	2		T ₃
Pond layer	Fish species	No. of species/ dec	No. of species/ ha	No. of species/ dec	No. of species / ha	No. of specie s/ dec	No. of species/h a
Surface	1. H. molitrix	8	1976	9	2223	10	2470
layer	2. C. catla	2	494	3	741	4	988
Middle layer	3. <i>L. rohita</i>	9	2223	10	2470	11	2717
Bottom	4. C. cirrhosus	4	988	5	1235	6	1482
layer	5. C. carpio	2	494	3	741	4	988
All Javer	6. C. idella	5	1235	5	1235	5	1235
	7. B. gonionotus	10	2470	10	2470	10	2470
	Total	40	9880	45	11115	50	12350

Table 5.1: Stocking density and stocking ratio/combination of fish species.

After stocking, TSP was applied (2.5 kg/ha/day) for ponds under all treatments as periodic fertilization.

In ponds under all treatments, Azolla (100%) was supplied as supplementary feed daily at the rate of 100% of the body weight of herbivorous fishes (*C. idella* and *B. gonionotus*). Azolla was supplied from Azolla bank and made available 24 hours per day (Plate 5.2).

5.3.5: Water quality monitoring

Some important physico-chemical parameters of water such as water temperature, transparency, dissolved oxygen (DO), pH, alkalinity and carbondioxide (CO₂) were monitored fortnightly between 10:00 AM to 11:00 AM for the present study (Plate 5.3).

5.3.5.1: Water temperature

During the study period, water temperature was recorded with the help of a celsius thermometer at 20-30 cm depth of water. The data were expressed as °C.

5.3.5.2: Water transparency

Measurement of limit of visibility i.e. penetration of light in water was done by a Secchi disc. The data thus obtained, were expressed as secchi disc depth in centimeter.

5.3.5.3: Dissolved oxygen (DO)

The dissolved oxygen concentration of water was determined by the aid of a water quality test kit (HACH kit FF-2, USA). Alkaline iodide-azide powder pillows, manganese sulfate powder pillows, sodium thiosulfate titration cartridge (0.2000 N), starch indicator solution and sulfamic acid powder pillows were used for determination of dissolved oxygen. The concentration of dissolved oxygen thus estimated was expressed in milligram per litter (mg/l) of water.

5.3.5.4: Hydrogen ion concentration (pH)

pH was measured by using HACH kit (FF-2, USA). Wide range 4 pH indicator solution (1919-00) were used for determination of water pH. A colour comparator disc ranging from 1-14 was also used for this purpose.

5.3.5.5: Alkalinity

Alkalinity was determined through digital titration by the help of a HACH kit (FF-2, USA). Bromcresol green-methyl red powder pillows, phenolphthalein powder pillows and sulfuric acid titration cartridge (0.1600 N) were used for total alkalinity determination. It was also expressed as mg/l of water.

5.3.5.6: Free carbon dioxide (CO₂)

Free carbon dioxide was determined through digital titration by the help of a HACH kit (FF-2, USA). Phenolphthalein powder pillows and sodium hydroxide titration cartridge (0.3636 N) were used for determination of carbon dioxide. It was also expressed as mg/l of water.

5.3.6: Growth monitoring of fishes

At least 10% (by number) of the fish in each pond were randomly sampled on a monthly basis with a cast net. On each sampling day, individual fish from each pond were weighed and measured. The purpose was to determine fish growth in weight and to adjust the ration. Following growth parameters were used for the present study (Plate 5.4).

Weight gain (g) = Mean final weight (g) - Mean initial weight (g)

Final weight (g) = Weight of fish at harvest (g)

Specific Growth Rate, SGR (% bwd⁻¹) = [L_n (final weight) – L_n (initial weight)] / culture period (days) \times 100

(Brown, 1957)

Survival rate (%) = $\frac{\text{Number of fish harvested}}{\text{Number of fish stocked}} \times 100$

Fish yield (kg/ha) = Fish biomass at harvest – Fish biomass at stock

5.3.7: Economics

Simple economical analysis of the different treatments was performed during the study period. On the basis of the fixed and variable expenditure, the total cost (BDT/ha) was estimated. At the end of the experiment, fishes were sold locally and the gross benefit (BDT/ha) was estimated. Net benefit (BDT/ha) was estimated by deducting the total cost from gross benefit. The following parameters were used to explore the economics of different treatments:

Net profit margin (%) = $\frac{\text{Net benefit}}{\text{Total investment}} \times 100$

CBR (Cost-Benefit Ratio) = $\frac{\text{Net benefit}}{\text{Total investment}}$

5.3.8: Statistical analysis

Data on water quality parameters, fish production and economics under different treatments were subjected to one way ANOVA (Analysis of Variance) using computer software SPSS (Statistical Package for Social Science, version-11). The mean values were also compared to see the significant difference from the DMRT (Duncan Multiple Range Test) after Gomez and Gomez (1984).





Plate 5.1: Experimental ponds



Plate 5.2: Supply of Azolla in experimental pond.



Plate 5.3: Monitoring of water quality parameters





Plate 5.4: Measurement of growth of fishes.

5.4: Results

5.4.1: Water quality

5.4.1.1: Fortnightly variations

The variations in the mean values of water quality parameters in different treatments at different fortnights are presented in table 5.2 to table 5.13.

Water temperature

During the study period, water temperature varied from 30.13 ± 0.15 to 33.47 ± 0.18 °C. The minimum value was recorded in treatment T₃ at 12th fortnight whereas the maximum value was recorded in treatment T₃ at 6th fortnight.

Water transparency

Water transparency value was found to range from 30.33 ± 0.88 to 34.67 ± 0.33 cm. The minimum value was recorded in treatment T₃ at 8th fortnight whereas the maximum value was recorded in treatment T₁ at 2nd and 4th fortnight.

Dissolved Oxygen (DO)

Dissolved oxygen of water varied from 5.00 ± 0.06 to 6.15 ± 0.10 mg/l. The minimum value was recorded in treatment T₃ at 7th and 8th fortnight whereas the maximum value was recorded in treatment T₁ at 4th fortnight.

Hydrogen ion concentration (pH)

pH value of water varied from 6.93 ± 0.07 to 8.10 ± 0.06 . The minimum value was recorded in treatment T₃ at 7th fortnight whereas the maximum value was recorded in treatment T₃ at 3rd fortnight.

Alkalinity

Alkalinity of water was found to range from 106.33 ± 1.76 to 123.00 ± 1.53 mg/l. The minimum value was recorded in treatment T₂ at 10th fortnight whereas the maximum value was recorded in treatment T₃ at 3rd fortnight.

Free carbon dioxide (CO₂)

Free CO₂ value was found to range from 2.03 ± 0.03 to 3.18 ± 0.07 mg/l. The minimum value was recorded in treatment T₁ at 6th fortnight whereas the maximum value was recorded in treatment T₃ at 12th fortnight.

Among the water quality parameters the values of transparency (at 2nd, 7th and 11th fortnight), DO (at 1st, 2nd, 3rd, 5th and 6th fortnight), pH (at 3rd fortnight), alkalinity (at 3rd, 4th, 7th, 9th and 10th fortnight) and CO₂ (at 3rd, 4th, 8th and 10th fortnight) differed significantly among the treatments.

Treatments Parameters	Τ ₁	T ₂	T ₃
Water temperature (°C)	32.47±0.20ª	32.30±0.17ª	32.30±0.10ª
Transparency (cm)	32.33±1.45ª	31.33±0.88ª	32.33±0.88ª
DO (mg/l)	5.63±0.07ª	5.13±0.09 ^b	5.20±0.06 ^b
рН	7.57±0.15ª	7.48±0.07ª	7.52±0.09ª
Alkalinity (mg/l)	109.33±0.33ª	113.00±3.61ª	116.00±2.31ª
Free CO ₂ (mg/l)	2.73±0.05ª	3.05±0.03ª	2.96±0.01ª

Table-5.2: Variations in the mean values of water quality parameters under different treatments at 1st fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-5.3: Variations in the mean values of water quality parameters under different treatments at 2nd fortnight

Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	32.30±0.12ª	32.47±0.20ª	32.10±0.15ª
Transparency (cm)	34.67±0.33ª	32.00±0.58 ^b	33.67±0.33ª
DO (mg/l)	5.80±0.06ª	5.27±0.12 ^b	5.17±0.15 ^b
рН	7.43±0.09ª	7.20±0.12ª	7.43±0.09ª
Alkalinity (mg/l)	115.00±1.53ª	116.00±4.04ª	109.67±0.88ª
Free CO ₂ (mg/l)	2.88±0.04ª	2.95±0.05ª	3.01±0.05ª

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Treatments Parameters	Τ ₁	T ₂	T ₃
Water temperature (°C)	31.43±0.23ª	31.30±0.06ª	31.30±0.06ª
Transparency (cm)	33.00±0.58ª	33.67±0.88ª	33.33±0.88ª
DO (mg/l)	5.93±0.09ª	5.40±0.25 ^{ab}	5.37±0.03 ^b
рН	7.83±0.09 ^{ab}	7.70±0.10 ^b	8.10±0.06ª
Alkalinity (mg/l)	115.33±4.26ªb	123.00±1.53ª	110.00±1.15 ^b
Free CO ₂ (mg/l)	2.65±0.07 ^b	3.01±0.04ª	3.05±0.10ª

Table-5.4: Variations in the mean values of water quality parameters under different treatments at 3rd fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.5: Variations in the mean values of water quality parameters under different treatments at 4th fortnight

Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	31.17±0.03ª	31.23±0.09ª	31.43±0.09ª
Transparency (cm)	34.67±0.33ª	32.00±1.15ª	32.33±0.67ª
DO (mg/l)	6.15±0.10ª	5.98±0.06ª	5.82±0.08ª
рН	7.83±0.12ª	7.73±0.19ª	7.63±0.19ª
Alkalinity (mg/l)	118.00±1.73ª	108.67±1.76 ^b	112.33±3.53ªb
Free CO ₂ (mg/l)	2.81±0.05 ^b	2.95±0.07 ^{ab}	3.03±0.03ª

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	33.46±0.22ª	33.43±0.19ª	33.17±0.12ª
Transparency (cm)	32.97±0.03ª	31.00±0.58 ^b	33.00±0.58ª
DO (mg/l)	5.38±0.06ª	5.20±0.09 ^{ab}	5.03±0.09 ^b
рН	7.23±0.12ª	7.03±0.03ª	7.27±0.18ª
Alkalinity (mg/l)	113.33±1.20ª	113.67±2.40ª	111.00±3.21ª
Free CO ₂ (mg/l)	2.67±0.09ª	2.73±0.09ª	2.90±0.06ª

Table-5.6: Variations in the mean values of water quality parameters under different treatments at 5th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-5.7: Variations in the mean values of water quality parameters under different treatments at 6th fortnight

Treatments Parameters	T1	T ₂	T ₃
Water temperature (°C)	33.27±0.12ª	33.43±0.18ª	33.47±0.18ª
Transparency (cm)	34.33±0.33ª	32.33±0.88ª	33.00±0.58ª
DO (mg/l)	5.67±0.09ª	5.50±0.06ª	5.20±0.06 ^b
рН	7.12±0.06ª	7.20±0.06ª	7.05±0.03ª
Alkalinity (mg/l)	109.00±1.53ª	107.67±1.45ª	107.67±1.33ª
Free CO ₂ (mg/l)	2.03±0.03ª	2.10±0.06ª	2.17±0.07ª

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	32.13±0.09ª	32.47±0.20ª	32.30±0.12ª
Transparency (cm)	32.00±0.58 ^b	34.67±0.33ª	34.00±0.58ª
DO (mg/l)	5.17±0.12ª	5.10±0.06ª	5.00±0.06ª
рН	6.97±0.09ª	7.00±0.06ª	6.93±0.07ª
Alkalinity (mg/l)	112.67±0.88ª	107.00±1.53 ^b	113.00±1.15ª
Free CO ₂ (mg/l)	2.63±0.07ª	2.68±0.02ª	2.80±0.06ª

Table-5.8: Variations in the mean values of water quality parameters under different treatments at 7th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.9: Variations in the mean values of water quality parameters under different treatments at 8th fortnight

Treatments Parameters	Τ ₁	T ₂	T ₃
Water temperature (°C)	32.47±0.20ª	32.20±0.15ª	32.27±0.09ª
Transparency (cm)	31.00±0.58ª	31.00±0.58ª	30.33±0.88ª
DO (mg/l)	5.10±0.06ª	5.07±0.12ª	5.00±0.06ª
рН	7.27±0.09ª	7.03±0.12ª	7.07±0.12ª
Alkalinity (mg/l)	108.67±1.86ª	111.33±1.20ª	108.67±1.76ª
Free CO ₂ (mg/l)	2.47±0.12 ^b	3.01±0.07ª	2.90±0.06ª

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	31.39±0.09ª	31.33±0.18ª	31.23±0.12ª
Transparency (cm)	33.00±0.58ª	31.67±0.33ª	33.00±1.15ª
DO (ppm)	5.10±0.06ª	5.07±0.12ª	5.03±0.03ª
рН	7.20±0.06ª	7.03±0.03ª	7.10±0.15ª
Alkalinity (ppm)	115.33±0.88ª	112.67±2.03ª	107.67±0.88 ^b
Free CO ₂ (ppm)	2.37±0.09ª	2.46±0.09ª	2.52±0.03ª

Table-5.10: Variations in the mean values of water quality parameters under different treatments at 9th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.11: Variations in the mean values of water quality parameters under different treatments at 10th fortnight

Treatments Parameters	Τ ₁	T ₂	T₃
Water temperature (°C)	31.47±0.09ª	31.33±0.03ª	31.30±0.17ª
Transparency (cm)	32.33±0.88ª	31.33±0.88ª	32.33±1.45ª
DO (mg/l)	5.33±0.12ª	5.17±0.09ª	5.13±0.09ª
рН	7.17±0.12ª	7.33±0.12ª	7.13±0.09ª
Alkalinity (mg/l)	115.00±2.31ª	106.33±1.76 ^b	110.67±0.88 ^{ab}
Free CO ₂ (mg/l)	2.70±0.06 ^b	2.95±0.03 ^{ab}	3.13±0.07ª

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)
Treatments Parameters	T ₁	T ₂	T ₃
Water temperature (°C)	30.53±0.09ª	30.37±0.09ª	30.17±0.27ª
Transparency (cm)	34.00±0.58ª	32.33±0.67ªb	31.33±0.33 ^b
DO (mg/l)	5.20±0.12ª	5.10±0.06ª	5.07±0.07ª
рН	7.00±0.06ª	7.07±0.03ª	7.07±0.07ª
Alkalinity (mg/l)	110.33±3.38ª	110.33±0.33ª	115.67±1.76ª
Free CO ₂ (mg/l)	2.67±0.09ª	2.73±0.07ª	2.81±0.09ª

Table-5.12: Variations in the mean values of water quality parameters under different treatments at 11th fortnight

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-5.13: Variations in the mean values of water quality parameters under different treatments at 12th fortnight

Treatments Parameters	Τ ₁	T ₂	T ₃
Water temperature (°C)	30.73±0.12ª	30.17±0.27ª	30.13±0.15ª
Transparency (cm)	33.00±1.53ª	33.67±0.88ª	32.67±0.33ª
DO (mg/l)	5.70±0.06ª	5.67±0.03ª	5.57±0.03ª
рН	7.57±0.07ª	7.40±0.06ª	7.57±0.03ª
Alkalinity (mg/l)	108.67±1.20ª	111.67±1.45ª	107.33±1.20ª
Free CO ₂ (mg/l)	3.05±0.06ª	3.13±0.09ª	3.18±0.07ª

5.4.1.2 Mean variations

The variations in the mean vales of different water quality parameters in different treatments by the total of all fortnights are presented in table 5.14 and fig. 5.1.

Water temperature

During the study period the mean values of water temperature ranged from 31.76 ± 0.30 to 31.93 ± 0.27 °C. The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₁.

Water transparency

The mean value of water transparency varied from 32.25 ± 0.34 to 33.11 ± 0.33 cm. The minimum value was recorded in treatment T₂ whereas the maximum value was recorded in treatment T₁.

Dissolved Oxygen (DO)

The mean value of DO varied from 5.32 ± 0.09 to 5.41 ± 0.09 mg/l. The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₁.

Hydrogen ion concentration (pH)

The mean value of water pH varied from 7.25 ± 0.07 to 7.35 ± 0.09 . The minimum value was recorded in treatment T₂ whereas the maximum value was recorded in treatment T₁.

Alkalinity

The mean value of total alkalinity varied from 110.81 ± 0.85 to 112.56 ± 0.94 mg/l. The minimum value was recorded in treatment T₃ whereas the maximum value was recorded in treatment T₁.

Free carbon-dioxide (CO₂)

The mean value of free CO₂ varied from 2.64 ± 0.07 to 2.87 ± 0.08 . The minimum value was recorded in treatment T₁ whereas the maximum value was recorded in treatment T₃.

No significant difference was found among the treatments for mean values of water quality parameters.

Treatments Parameters	Τ ₁	T ₂	T ₃
Water temperature (°C)	31.93±0.27ª	31.84±0.31ª	31.76±0.30ª
Transparency (cm)	33.11±0.33ª	32.25±0.34ª	32.61±0.29ª
DO (mg/l)	5.41±0.09ª	5.35±0.11ª	5.32±0.09ª
рН	7.35±0.09ª	7.25±0.07ª	7.29±0.10ª
Alkalinity (mg/l)	112.56±0.94ª	111.78±1.32ª	110.81±0.85ª
Free CO ₂ (mg/l)	2.64±0.07ª	2.81±0.09ª	2.87±0.08ª

Table 5.14: Variations in the mean values of water quality parameters under different treatments during study period







b

d









Fig. 5.1: Variations in the mean values of water quality parameters (a, Water temperature; b, transparency; c, DO; d, pH; e, alkalinity; f, free CO₂) under different treatments during study period.

5.4.2: Growth of fishes

5.4.2.1: Monthly variations

Specific growth rate (SGR, %, bwd⁻¹)

The specific growth rate (%, bwd⁻¹) varied with *H. molitrix* from 0.53 ± 0.01 (T₂ at 6th month) to 2.68±0.02 (T₁ at 1st month), *C. catla* from 0.48 ± 0.003 (T₃ at 6th month) to 2.63±0.003 (T₁ at 1st month), *L. rohita* from 0.39 ± 0.01 (T₂ at 6th month) to 2.04±0.07 (T₁ at 1st month), *C. cirrhosus* from 0.39 ± 0.02 (T₂ at 6th month) to 2.30±0.11 (T₂ at 1st month), *C. carpio* from 0.50 ± 0.03 (T₃ at 6th month) to 2.28±0.16 (T₁ at 1st month), *C. idella* from 0.49 ± 0.01 (T₂ at 6th month) to 2.98±0.04 (T₁ at 1st month) and *B. gonionotus* from 0.46 ± 0.02 (T₃ at 6th month) to 3.42±0.07 (T₁ at 1st month) (Table 5.15 to 5.20).

Among the species, the highest SGR was observed in *B. gonionotus* as 3.42 ± 0.07 %, bwd⁻¹ and the lowest was in *L. rohita* as 0.39 ± 0.01 %, bwd⁻¹.

The lowest combined SGR value was found as 0.47 ± 0.022 (T₂ at 6th month) and the highest combined SGR was found as 2.60 ± 0.18 (T₁ at 1st month). The highest SGR value was found in the month of April and the lowest value was found in the month of September for all the species.

Significant variation was found in 1st month (April) for all species.

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	2.68±0.02ª	2.57±0.02 ^b	2.48±0.02°
C. catla	2.63±0.03ª	2.50±0.03 ^b	2.40±0.02°
L. rohita	2.04±0.07ª	1.86±0.04 ^b	1.76±0.03 ^b
C. cirrhosus	2.20±0.15ª	2.30±0.11ª	1.93±0.12ª
C. carpio	2.28±0.16ª	2.02±0.06ª	2.12±0.03ª
C. idella	2.98±0.04ª	2.81±0.06 ^{ab}	2.74±0.07 ^b
B. gonionotus	3.42±0.07ª	3.21±0.04 ^{ab}	3.07±0.09 ^b
All species	2.60±0.18ª	2.47±0.17 ^{ab}	2.36±0.17 ^b

Table-5.15: Variations in the mean values of SGR (%, bwd⁻¹) of fishes under different treatments at 1st month (April)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.16:	Variations	in th	e mean	values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments	at 2nd	month	(May)								

Treatments Species	Τ ₁	T ₂	T ₃
H. molitrix	1.53±0.02ª	1.48±0.01 ^b	1.46±0.01 ^b
C. catla	1.50±0.01ª	1.43±0.02 ^b	1.43±0.01 ^b
L. rohita	1.31±0.01ª	1.23±0.06ª	1.29±0.01ª
C. cirrhosus	1.42±0.05ª	1.24±0.04 ^b	1.30±0.03 ^{ab}
C. carpio	1.70±0.11ª	1.74±0.08ª	1.59±0.07ª
C. idella	1.59±0.05ª	1.59±0.02ª	1.53±0.01ª
B. gonionotus	1.70±0.05ª	1.73±0.02ª	1.64±0.04ª
All species	1.54±0.05ª	1.49±0.08ª	1.46±0.05ª

Treatments Species	Τ ₁	T ₂	T ₃
H. molitrix	1.11±0.01ª	1.07±0.003 [♭]	1.05±0.01 ^b
C. catla	1.17±0.02ª	1.17±0.03ª	1.18±0.01ª
L. rohita	1.01±0.03ª	1.04±0.02ª	1.01±0.01ª
C. cirrhosus	1.12±0.04ª	0.98±0.08 ^b	0.97±0.03 ^b
C. carpio	1.25±0.03ª	1.23±0.04ª	1.09±0.03 ^b
C. idella	1.17±0.02ª	1.11±0.02ª	1.02±0.03 ^b
B. gonionotus	1.36±0.05ª	1.31±0.05ª	1.31±0.11ª
All species	1.17±0.04ª	1.13±0.02ª	1.09±0.32ª

Table-5.17: Variations in the mean values of SGR (%, bwd⁻¹) of fishes under different treatments at 3rd month (June)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.18:	Variations	in th	e mean	values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments	at 4th	month	(July)								

Treatments Species	Τ ₁	T ₂	T ₃
H. molitrix	0.83±0.003ª	0.81±0.01 ^b	0.80±0.003 ^b
C. catla	0.85±0.02ª	0.88±0.02ª	0.87±0.01ª
L. rohita	0.82±0.02ª	0.80±0.01ª	0.81±0.02ª
C. cirrhosus	0.89±0.02ª	0.80±0.02 ^b	0.74±0.01°
C. carpio	0.89±0.03ª	0.89±0.05ª	0.92±0.02ª
C. idella	0.84±0.04ª	0.83±0.02ª	0.83±0.02ª
B. gonionotus	0.99±0.01ª	1.00±0.01ª	1.05±0.03ª
All species	0.87±0.02ª	0.86±0.03ª	0.86±0.04ª

Treatments Species	T₁	T ₂	Τ₃
H. molitrix	0.66±0.003ª	0.65±0.01ª	0.64±0.01ª
C. catla	0.59±0.003ª	0.59±0.01ª	0.58±0.003ª
L. rohita	0.47±0.02ª	0.50±0.02ª	0.48±0.01ª
C. cirrhosus	0.54±0.01 ^{ab}	0.56±0.01ª	0.51±0.02 ^b
C. carpio	0.64±0.02ª	0.63±0.01ª	0.60±0.01ª
C. idella	0.61±0.01ª	0.60±0.01ª	0.61±0.003ª
B. gonionotus	0.59±0.01ª	0.59±0.02ª	0.59±0.02ª
All species	0.59±0.024ª	0.59±0.018ª	0.57±0.022ª

Table-5.19: Variations in the mean values of SGR (%, bwd⁻¹) of fishes under different treatments at 5th month (August)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.20:	Variations	in t	the	mean	values	of	SGR	(%,	bwd ⁻¹)	of	fishes	under	different
	treatments	at 6	3 th m	nonth	(Septen	nbe	ər)						

Treatments Species	Τ ₁	T ₂	T ₃
H. molitrix	0.54±0.003ª	0.53±0.01ª	0.55±0.01ª
C. catla	0.51±0.003ª	0.50±0.02ª	0.48±0.003ª
L. rohita	0.43±0.01ª	0.39±0.01ª	0.43±0.01ª
C. cirrhosus	0.44±0.03ª	0.39±0.02ª	0.42±0.01ª
C. carpio	0.50±0.03ª	0.52±0.02ª	0.50±0.03ª
C. idella	0.50±0.01ª	0.49±0.01ª	0.53±0.02ª
B. gonionotus	0.50±0.01ª	0.47±0.01ª	0.46±0.02ª
All species	0.49±0.015ª	0.47±0.022ª	0.48±0.018ª

Monthly weight gain (g/month)

The weight gain (g/month) varied with *H. molitrix* from 66.33 ± 0.88 g (T₃ at 1st month) to 84.00 ± 1.53 g (T₁ at 4th month), *C. catla* from 68.33 ± 0.88 g (T₃ at 1st month) to 94.33 ± 1.76 g (T₁ at 3rd month), *L. rohita* from 38.67 ± 1.33 g (T₃ at 5th month) to 59.67 ± 0.88 g (T₁ at 4th month), *C. cirrhosus* from 42.67 ± 2.33 g (T₃ at 5th month) to 73.33 ± 2.40 g (T₁ at 4th month), *C. carpio* from 51.00 ± 2.08 (T₂ at 1st month) to 91.67 ± 3.28 g (T₁ at 3rd month), *C. idella* from 89.00 ± 3.21 (T₃ at 5th month) to 115.67 ± 1.76 g (T₁ at 3rd month) and *B. gonionotus* from 32.33 ± 1.33 (T₃ at 6th month) to 53.33 ± 0.33 g (T₁ at 4th month) (Table 5.21 - 5.26).

Among the species, the highest monthly weight gain was observed in *C. idella* as 115.67 ± 1.76 g and the lowest was in *B. gonionotus* as 32.33 ± 1.33 g.

The lowest combined weight gain value was found as 56.81 ± 7.29 g (T₃ at 1st month) and the highest combined weight was found as 80.62 ± 7.64 g (T₁ at 4th month). The highest weight gain was found in the month of July and the lowest value was found in the month of April for all the species.

Significant variation in monthly weight gain was found under different treatments in all the months for all species.

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	74.33±0.67ª	69.67±0.88 ^b	66.33±0.88°
C. catla	78.33±1.20ª	72.67±1.20 ^b	68.33±0.88°
L. rohita	49.00±2.31ª	43.33±1.20 ^b	40.33±0.88 ^b
C. cirrhosus	54.33±4.81ª	57.67±3.84ª	45.67±3.84ª
C. carpio	60.00±6.08ª	51.00±2.08ª	54.33±1.20ª
C. idella	101.00±2.08ª	92.67±2.96ªb	89.33±3.48 ^b
B. gonionotus	39.33±1.20ª	35.67±0.67 ^{ab}	33.33±1.45⁵
All species	65.19±7.89ª	60.38±7.36 ^b	56.81±7.29°

Table-5.21: Variations in the mean weight gain (g/month) of fishes under different treatments at 1st month (April)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.22: Variations in the mean weight gain (g/month) of fishes under different treatments at 2nd month (May)

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	78.33±1.76ª	72.33±1.20 ^b	69.33±0.67 ^b
C. catla	81.67±1.20ª	73.67±1.76 ^b	71.67±0.88 ^b
L. rohita	51.33±1.45ª	45.33±2.03 ^b	46.67±0.33ªb
C. cirrhosus	59.67±0.88ª	52.00±1.73 ^b	49.33±1.20 ^b
C. carpio	80.00±3.21ª	77.00±4.04ª	70.67±3.38ª
C. idella	104.33±3.18ª	99.67±0.67 ^{ab}	92.67±2.73 ^b
B. gonionotus	40.67±0.88ª	39.33±0.33 ^{ab}	35.33±2.03 ^b
All species	70.86±8.16ª	65.62±8.00 ^b	62.24±7.35°

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	83.67±1.86ª	76.67±0.88 ^b	72.67±0.88 ^b
C. catla	94.33±1.76ª	88.67±2.40 ^{ab}	87.33±0.88 ^b
L. rohita	56.33±1.76ª	53.67±1.33ª	51.67±0.88ª
C. cirrhosus	69.00±4.36ª	57.33±3.48 ^b	51.67±1.45 ^b
C. carpio	91.67±3.28ª	84.67±2.33ª	72.00±2.08 ^b
C. idella	115.67±1.76ª	104.33±2.19 ^b	91.00±4.51°
B. gonionotus	51.33±2.03ª	46.67±2.03 ^b	43.67±3.53 ^b
All species	80.29±8.63ª	73.14±8.00 ^b	67.14±7.00°

Table-5.23: Variations in the mean weight gain (g/month) of fishes under different treatments at 3rd month (June)

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.24: Variations in the mean weight gain (g/month) of fishes under different treatments at 4th month (July)

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	84.00±1.53ª	76.67±0.33 ^b	73.33±0.67 ^b
C. catla	92.33±1. 86ª	91.33±1.76ª	87.33±1.45ª
L. rohita	59.67±0.88ª	54.67±0.88 ^b	54.33±1.76 ^b
C. cirrhosus	73.33±2.40ª	61.33±1.86 ^b	51.33±1.86°
C. carpio	89.67±1.45ª	83.67±4.84ª	82.00±1.53ª
C. idella	112.00±4.73ª	103.67±3.18 ^{ab}	96.33±3.71 ^b
B. gonionotus	53.33±0.33ª	50.33±0.88 ^b	49.33±0.67 ^b
All species	80.62±7.64ª	74.52±7.52 ^b	70.57±7.18°

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	83.00±2.00ª	76.33±0.33 ^b	73.00±0.58 ^b
C. catla	80.00±1.15ª	76.00±2.31ªb	71.67±1.20 ^b
L. rohita	41.67±1.20ª	41.00±1.53ª	38.67±1.33ª
C. cirrhosus	55.67±2.40ª	52.00±1.00ª	42.67±2.33 ^b
C. carpio	80.67±1.20ª	74.00±3.21 ^{ab}	67.00±1.73 ^b
C. idella	101.00±2.08ª	93.00±2.52 ^{ab}	89.00±3.21 ^b
B. gonionotus	40.00±0.58ª	37.67±1.45 ^{ab}	35.67±0.67 ^b
All species	68.86±8.79ª	64.29±7.87 ^b	59.67±7.78°

Table-5.25:	Variations	in	the	mean	weight	gain	(g/month)	of	fishes	under	different
	treatments	at {	5 th m	onth (A	August)						

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05)

Table-5.26:	Variations	in	the	mean	weight	gain	(g/month)	of	fishes	under	different
	treatments	at	6 th m	onth (S	Septemb	er)					

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	81.67±2.03ª	74.67±0.88 ^b	73.33±2.03 ^b
C. catla	81.67±0.33ª	75.67±1.76 ^b	69.67±1.45°
L. rohita	43.33±0.88ª	37.00±1.53 ^b	39.67±0.88ªb
C. cirrhosus	52.33±2.19ª	42.67±3.18 ^b	40.00±1.15 ^b
C. carpio	75.67±3.53ª	72.67±2.19ª	65.33±3.18ª
C. idella	97.67±2.33ª	89.33±1.20 ^b	91.33±1.33 ^b
B. gonionotus	39.67±0.88ª	35.33±0.88 ^b	32.33±1.33 ^b
All species	67.43±8.41ª	61.05±8.33 ^b	58.81±8.24°

5.4.2.2: Mean variations

Variations in the mean values of growth parameters of fishes under different treatments are shown in table 5.27 to 5.31, fig. 5.2 to 5.4 and plate 5.5.

SGR (%, bwd⁻¹)

The specific growth rate (SGR, %, bwd⁻¹) varied with *H. molitrix* from 1.16±0.30 (T₃) to 1.23±0.32 (T₁), *C. catla* from 1.16±0.29 g (T₃) to 1.21±0.32 (T₁), *L. rohita* from 0.96±0.21 (T₃) to 1.01±0.25 (T₁), *C. cirrhosus* from 0.98±0.23 (T₃) to 1.10±0.26 (T₁), *C. carpio* from 1.14±0.25 (T₃) to 1.21±0.28 (T₁), *C. idella* from 1.21±0.34 (T₃) to 1.28±0.38 (T₁) and *B. gonionotus* from 1.35±0.39 (T₃) to 1.43±0.44 (T₁).

Among the species the highest SGR was observed as 1.43 ± 0.44 %, bwd⁻¹ in *B*. *gonionotus* and the lowest was in *L. rohita* as 0.96 ± 0.21 %, bwd⁻¹.

The lowest combined SGR value was found as 1.14 ± 0.05 (T₃) and the highest combined SGR as 1.21 ± 0.05 (T₁).

Significant difference was found among the treatments for the mean values of SGR of all the species.

Weight gain (g/month)

Mean weight gain (g/month) varied with *H. molitrix* from 71.33 \pm 1.18 g (T₃) to 80.83 \pm 1.55 g (T₁), *C. catla* from 76.00 \pm 3.62 g (T₃) to 84.72 \pm 2.78g (T₁), *L. rohita* from 45.22 \pm 2.73 g (T₃) to 50.22 \pm 2.89g (T₁), *C. cirrhosus* from 46.78 \pm 1.96 g (T₃) to 60.72 \pm 3.49 g (T₁), *C. carpio* from 68.56 \pm 3.71 g (T₃) to 79.61 \pm 4.65 g (T₁), *C. idella* from 91.61 \pm 1.09 g (T₃) to 105.28 \pm 2.88 g (T₁) and *B. gonionotus* from 38.28 \pm 2.75 g (T₃) to 44.06 \pm 2.64 g (T₁).

Among the species the highest weight gain was observed as 105.28 ± 2.88 g in *C. idella* and the lowest was in *B. gonionotus* as 38.28 ± 2.75 g.

The lowest mean weight gain was found as 62.54 ± 7.36 g (T₃) and the highest weight gain as 72.21 ± 8.15 g (T₁) for all the species.

Significant difference was found among the treatments for the mean values of weight gain of all the species.

Final Weight (g)

The mean final weight (g) varied with *H. molitrix* from 468.00 ± 24.01 (T₃) to 545.00 ± 9.61 g (T₁), *C. catla* from 521.00 ± 6.25 (T₃) to 573.33 ± 3.71 g (T₁), *L. rohita* from 329.33 ± 3.71 (T₃) to 359.33 ± 3.18 g (T₁), *C. cirrhosus* from 338.67 ± 10.17 (T₃) to 422.33 ± 11.61 g (T₁), *C. carpio* from 472.33 ± 3.71 (T₃) to 538.67 ± 3.84 g (T₁), *C. idella* from 619.67 ± 16.48 (T₃) to 701.67 ± 7.22 g (T₁) and *B. gonionotus* from 251.33 ± 4.06 (T₃) to 287.67 ± 1.67 g (T₁).

Among the species the highest final weight was observed as 701.67 ± 7.22 g in *C. idella* and the lowest was in *B. gonionotus* as 251.33 ± 4.06 g.

The lowest mean final weight was found as 3000.33 ± 31.25 g (T₃) and the highest final weight as 3428.00 ± 25.35 g (T₁) for all the species.

Significant difference was found among the treatments for the mean values of final weight of all the species.

Survival rate (%)

The survival rate (%) varied with *H. molitrix* from 85.33 ± 1.17 (T₃) to $90.33\pm2.91\%$ (T₁), *C. catla* from 80.87 ± 0.58 (T₃) to $83.63\pm0.75\%$ (T₁), *L. rohita* from 83.10 ± 2.18 (T₁) to $84.13\pm1.21\%$ (T₂), *C. cirrhosus* from 85.50 ± 0.58 (T₂) to $91.53\pm1.13\%$ (T₁), *C. carpio* from $93.07\pm0.81\%$ (T₂) to $94.53\pm0.61\%$ (T₁), *C. idella* from 85.07 ± 0.96 (T₂) to $86.37\pm0.52\%$ (T₁) and *B. gonionotus* from 80.50 ± 1.15 (T₁) to $81.73\pm0.79\%$ (T₃).

Among the species the highest survival rate was observed as $94.53\pm0.61\%$ in *C. carpio* and the lowest was in *B. gonionotus* as $80.50\pm1.15\%$.

The lowest mean survival rate was found as $85.20\pm1.55\%$ g (T₃) and the highest survival rate as $87.14\pm1.94\%$ (T₁) for all the species.

No significant difference was found among the treatments for the mean values of survival rate of all the species except *C. cirrhosus*.

Yield (Kg/ha/6 months)

The individual fish yield (Kg/ha/6 months) varied with *H. molitrix* from 969.00 \pm 17.16 (T₁) to 982.67 \pm 50.35 (T₃), *C. catla* from 241.00 \pm 1.53 (T₁) to 411.67 \pm 4.81 (T₃), *L. rohita* from 603.67 \pm 5.21 (T₁) to 781.00 \pm 8.62 (T₃), *C. cirrhosus* from 375.67 \pm 10.17 (T₁) to 426.33 \pm 12.84 (T₃), *C. carpio* from 252.67 \pm 1.86 (T₁) to 419.67 \pm 3.38 (T₃), *C. idella* from 520.33 \pm 13.87 (T₃) to 884.00 \pm 9.07 (T₁) and *B. gonionotus* from 496.67 \pm 8.11 (T₃) to 568.33 \pm 3.33 (T₁).

Among the species the highest yield was observed as 982.67 ± 50.35 (T₃) kg/ha/6 months in *H. molitrix* and the lowest was in *C. catla* as 241.00 ± 1.53 (T₁) kg/ha/6 months.

The lowest total fish yield (Kg/ha/6 months) was found as 3894.33 ± 18.00 (T₁) and the highest total yield was found as 4038.33 ± 84.41 (T₃). Significant difference in total yield was found among the treatments.

Significant difference was found among the treatments for the mean values of yield of all the species.

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	1.23±0.32ª	1.19±0.31⁵	1.16±0.30 ^b
C. catla	1.21±0.32ª	1.18±0.30ª	1.16±0.29ª
L. rohita	1.01±0.25ª	0.97±0.22ª	0.96±0.21ª
C. cirrhosus	1.10±0.26ª	1.01±0.28 ^b	0.98±0.23 ^b
C. carpio	1.21±0.28ª	1.17±0.25ª	1.14±0.25ª
C. idella	1.28±0.38ª	1.22±0.35 ^b	1.21±0.34 ^b
B. gonionotus	1.43±0.44ª	1.39±0.41ª	1.35±0.39ª
All species	1.21±0.05ª	1.16±0.05 [♭]	1.14±0.05 ^b

Table-5.27: Variations in the mean SGR (%, bwd⁻¹) of fishes under different treatments during the study period

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.28: Variations in the mean weight gain (g/6 month) of fishes under different treatments during the study period.

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	80.83±1.55ª	74.39±1.17⁵	71.33±1.18 ^b
C. catla	84.72±2.78ª	79.67±3.32ª	76.00±3.62ª
L. rohita	50.22±2.89ª	45.83±2.87ª	45.22±2.73ª
C. cirrhosus	60.72±3.49ª	53.83±2.67 ^{ab}	46.78±1.96 ^b
C. carpio	79.61±4.65ª	73.84±4.99ª	68.56±3.71ª
C. idella	105.28±2.88ª	97.11±2.57⁵	91.61±1.09 ^b
B. gonionotus	44.06±2.64ª	40.83±2.54ª	38.28±2.75ª
All species	72.21±8.15ª	66.50±7.67 ^{ab}	62.54±7.36 ^b

Treatments Species	Τ ₁	T ₂	T ₃
H. molitrix	545.00±9.61ª	506.33±2.73 ^{ab}	468.00±24.01 ^b
C. catla	573.33±3.71ª	543.00±5.57 ^b	521.00±6.25°
L. rohita	359.33±3.18ª	333.00±4.00 ^b	329.33±3.71⁵
C. cirrhosus	422.33±11.61ª	381.00±7.00 ^b	338.67±10.17°
C. carpio	538.67±3.84ª	504.00±11.15 ^b	472.33±3.71°
C. idella	701.67±7.22ª	652.67±10.27⁵	619.67±16.48 ^b
B. gonionotus	287.67±1.67ª	267.00±2.52 ^b	251.33±4.06°
All species	3428.00±11.21ª	3187.00±14.05 ^b	3000.33±13.25 ^b

Table-5.29: Variations in the mean final weight (g) of fishes under different treatments during the study period.

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Table-5.30: Variations in the mean survival rate (%) of fishes under different treatments during the study period.

Treatments Species	T ₁	T ₂	T ₃
H. molitrix	90.33±2.91ª	85.50±0.76ª	85.33±1.17ª
C. catla	83.63±0.75ª	83.00±1.26ª	80.87±0.58ª
L. rohita	83.10±2.18ª	84.13±1.21ª	83.87±1.02ª
C. cirrhosus	91.53±1.13ª	85.50±0.58 ^b	85.73±0.62 ^b
C. carpio	94.53±0.61ª	93.07±0.81ª	93.47±1.55ª
C. idella	86.37±0.52ª	85.07±0.96ª	85.43±1.16ª
B. gonionotus	80.50±1.15ª	80.73±1.30ª	81.73±0.79ª
All species	87.14±1.94ª	85.29±1.45ª	85.20±1.55ª

Treatments			
	T ₁	T ₂	T ₃
Species			
H. molitrix	969.00±17.16ª	979.00±5.13ª	982.67±50.35 ^a
	(24.88%)	(24.74%)	(24.33%)
C. catla	241.00±1.53°	330.33±3.38 ^b	411.67±4.81ª
	(6.19%)	(8.35%)	(10.19%)
L. rohita	603.67±5.21°	699.00±8.50 ^b	781.00±8.62ª
	(15.50%)	(17.67%)	(19.34%)
C. cirrhosus	375.67±10.17 ^b	400.00±7.57 ^{ab}	426.33±12.84 ^a
	(9.65%)	(10.11%)	(10.56%)
C. carpio	252.67±1.86°	336.00±7.37 ^b	419.67±3.38ª
	(6.49%)	(8.49%)	(10.39%)
C. idella	884.00±9.07ª	684.67±10.81 ^b	520.33±13.87°
	(22.70%)	(17.30%)	(12.88%)
B. gonionotus	568.33±3.33ª	527.67±4.70 ^b	496.67±8.11°
	(14.59%)	(13.34%)	(12.30%)
Total yield	0004 00 40 000	0050 07 40 70h	4000 00 04 443
(Kg/ha/6 months)	3894.33±18.00°	3956.67±43.72°	4038.33±84.41ª
Total yield	7700 00 00 000	7040.04.04.00h	0070 00 00 003
(Kg/ha/yr)	7788.66±38.69°	7913.34±84.23°	8076.66±96.35ª

Table-5.31: Variations in the mean yield (kg/ha) of fishes under different treatments during the study period.

Figures bearing common letter(s) in a row as superscript do not differ significantly (P < 0.05)

Values in the parentheses means % contribution by individual species to total fish yield under respective treatment.



Fig. 5.2. Variations in the mean values of different growth parameters under different treatments during the study period: a; SGR (%, bwd⁻¹), b; weight gain (g), c; final weight (g) and d; survival rate (%)



Fig. 5.3: Variations in the mean values of fish yield under different treatments



Fig. 5.4: Variations in the mean values of total fish yield under different treatments





Plate 5.5: Harvesting of fishes.

5.4.3: Economics

The economics of fish farming under different treatments are presented in table 5.32 and fig. 5.5.

Total cost significantly varied from 136862.00 \pm 606.45 (T₁) to 145754.00 \pm 57.35 Tk/ha/6 months (T₃). Gross benefit significantly varied from 384562.35 \pm 1890.32 (T₁) to 402667 \pm 8862.88 Tk/ha/6 months (T₃). Net benefit significantly varied from 247700.35 \pm 1890.32 Tk/ha/6 months (T₁) to 256913.35 \pm 8862.88 Tk/ha/6 months (T₃). The Net profit margin significantly varied from 176.31 \pm 5.55 (T₃) to 181.00 \pm 1.53% (T₁). The CBR significantly varied from 1.76 \pm 0.06 (T₃) to 1.81 \pm 0.02 (T₁).

Treatments	T.	Ta	Ta
Components	• 1	12	13
Lease value (Tk.)	45000.00±0.00 ^a	45000.00±0.00 ^a	45000.00±0.00 ^a
Pond preparation (Tk.)	11480.00±57.74°	11480.00±98.56 ^b	(30.87 %) 11480.00±59.65ª
Fertilizer (Tk.)	(8.39%) 26790.00±0.00ª	(8.12%) 26790.00±0.00ª	(7.88%) 26790.00±0.00ª
Fish seed (Tk.)	(19.57%) 33592.00±0.00ª (24.54%)	(18.96%) 38038.00±0.00ª (36.02%)	(18.38%) 42484.00±0.00 ^a
Feed (Tk.)	(24.34%) 00.00±0.00 ^a (0.00%)	(20.92%) 00.00±0.00ª (0.00%)	(29.15%) 00.00±0.00ª (0.00%)
Harvesting cost (Tk.)	(0.00%) 20000.00±0.00ª (14.61%)	(0.00%) 20000.00±0.00ª (14.15%)	(0.00%) 20000.00±0.00ª (13.72%)
Total cost (Tk.)	136862.00±606.45°	141308.00±57.74 ^b	145754.00±57.35ª
Gross benefit (Tk.)	384562.35±1890.32 ^b	392651.95±4590.61 ^b	402667.35±8862.88ª
Net benefit (Tk.)	247700.35±1890.32 ^b	251343.95±4590.61 ^{ab}	256913.35±8862.88ª
Net profit margin (%)	181.00±1.53ª	178.33±2.96 ^{ab}	176.31±5.55 ^b
CBR	1.81±0.02ª	1.78±0.03 ^{ab}	1.76±0.06 ^b

Figures bearing common letter(s) in a row as superscript do not differ significantly (P <0.05) % of total cost in parentheses







Fig. 5.5: Economics of fish farming under different treatments: a; total cost (BDT.), b; net benefit (BDT) and c; CBR

5.4: Discussion

5.4.1: Water quality parameters

Fortnightly variation

Water temperature

Water temperature varied from 30.13 ± 0.15 (T₃ at 12^{th} fortnight) to 33.47 ± 0.18 °C (T₃ at 6^{th} fortnight). Lower temperature (30.13 ± 0.15) with treatment T₃ at 12^{th} fortnight (September 2012) might be due to start of winter season when air temperature was decreased gradually (Appendix-2). Higher water temperature (33.47 ± 0.18 °C) with treatment T₃ at 6^{th} fortnight (June, 2012) might be due to summer season (Appendix-2). The change of water temperature might be due to the change of weather condition from summer to winter season. No significant difference was found among the treatments for the fortnightly values of temperature. Wahab *et al.* (1996) recorded water temperature as 28.5 to 31.3° C in fish pond which was closer to the findings of the present study. The fortnightly fluctuation in water temperature appeared to be due to heavy rainfall resulting in reduced temperature condition as well as bright sunlight leading to warming up of the pond water. Vass and Sachlan (1955) stated that temperature.

Water transparency

In the present study, water transparency was found to range from 30.33 ± 0.88 (T₃ at 8th fortnight) to 34.67 ± 0.33 cm (T₁ at 2nd and 4th fortnight). The lower value of the water transparency of the treatment might be due to the direct effect of basal and monthly fertilization. The values of transparency also varied with sampling date, which could be due to differences in abundance of plankton and suspended particles. Azim *et al.* (1995) found similar water transparency (36.2 cm) in weed based system. Significant variation in transparency values was found at 2nd, 7th and 11th fortnights among the treatments, these variations might be due to rainfall and / or nutrient concentrations. Thy *et al.* (2004) reported that mean value of transparency (cm) was 30.7 in effluent plus

duckweed pond. Boyd (1982) recommended a transparency between 30 to 45 cm as appropriate for fish culture.

Dissolved Oxygen (DO)

Dissolved Oxygen (DO) content is probably the only most important water quality parameters in aquaculture. The DO concentrations varied from 5.00 ± 0.06 (T₃ at 7th and 8th fortnight) to 6.15 ± 0.10 mg/l (T₁ at 4th fortnight). Openheimer *et al.* (1978) and Wahab *et al.* (1995) recorded dissolved oxygen values ranged from 3.18 to 7.58 and 2.2 to 7.1 mg/l, respectively. Significant variations in DO values were found at 1st, 2nd, 3rd, 5th and 6th fortnights among the treatments. Fortnightly variations might be due to fluctuation of temperature (Appendix-2). The concentration of DO showed a decreasing trend with the progress of culture duration, obviously due to the increase in biomass with the progress of culture (Jena *et al.*, 1998).

Hydrogen ion concentration (pH)

The pH in the natural water has great importance as it regulates the productivity of water body. The values of pH in the water ranged from 6.93 ± 0.07 (T₃ at 7th fortnight) to 8.10 ± 0.06 (T₃ at 3rd fortnight). The neutral to slightly alkaline pH in the study pond were possibly due to local soil condition and natural waters. Wurts and Durbrow (1992) stated that pH of a waterbody depends both soil and water condition. Moreover, the initial lime treatment during pond preparation possibly helped in maintaining carbon buffer system in the pond water (Khan *et al.*, 2009). Ferdoushi *et al.* (2008) found the range of pH was 7.53±0.34 to 8.09±0.46 in weed based carp polyculture pond.

Total alkalinity

Total alkalinity was found to range from 106.33 ± 1.76 (T₂ at 10^{th} fortnight) to 123.00 ± 1.53 mg/l (T₂ at 4^{th} fortnight). It might be due to low dissolved oxygen content and more production of free CO₂ that enhanced by increasing of fish biomass which strongly supported by Michael (1968) while working with water quality parameters in fish ponds and tropical impoundment. Boyd (1998) stated that the acceptable range of alkalinity for freshwater fish culture is between 40

to 200 mg/l. Significant variations in alkaline values were found at 3^{rd} , 4^{th} , 7^{th} , 9^{th} and 10^{th} fortnights among the treatments. It might be due to variation in rainfall (Appendix-2). Total alkalinity values depending upon the location, season, plankton population, nature of bottom deposits etc. (Jhingran, 1991). In the present study, the alkalinity levels in water were within the productive range as stated by Rahman *et al.* (1982).

Free carbon dioxide (CO₂)

The mean values of free carbon dioxide varied from 2.03 ± 0.03 (T₁ at 6th fortnight) to 3.18 ± 0.07 mg/l (T₃ at 12th fortnight). The higher value of free CO₂ content was obtained during the last month of the experiment (September, 2012) due to higher fish production. The suitable range of free CO₂ for fish culture is ranged from 1.0 to 10.0 mg/l (Boyd, 1998). Significant variation was found at 3rd, 4th, 8th and 10th fortnights among the treatments. It might be due to variation in rainfall (Appendix-2). The fluctuation in CO₂ values might be due to alteration in the rate of photosynthesis in the pond and due to rate of DO consumption by fish through respiration. Talukdar *et al.* (2012) recorded the value of free CO₂ as 2.85 mg/l while working on suitability of aquatic macrophytes as fish feed in carp polyculture system.

Mean variation

Water temperature

In the present study, the mean value of water temperature varied from 31.76 ± 0.30 (T₃) to 31.93 ± 0.27 °C (T₁). The findings of the present study is quite similar to the findings recorded by Kabir *et al.* (2009) who worked on the duckweed fed polyculture in pond and reported the water temperature ranged between 29.50-31.65°C. Azim and Wahab (2003) found water temperature of 18.2 to 27.8°C in the duckweed fed polyculture pond. Water temperature of 25 to 32°C is considered suitable for fish culture (Boyd and Zimmermann, 2000). In another study, Effiong *et al.* (2009) found water temperature as 27.72±0.01°C in weed based system.

Water transparency

The mean value of water transparency in this study varied from 32.25 ± 0.34 (T₂) to 33.11 ± 0.33 cm (T₁). Azim and Wahab (2003) reported Secchi disc reading as 9.0-42.0 cm in duck weed fed polyculture pond. Chowdhury *et al.* (2008) found the transparency value ranged from 28.66 to 34.50 cm in pond where duckweed was used as supplementary feed. Boyd (1982) recommended a transparency between 30 to 45 cm as appropriate for fish culture. Wahab *et al.* (1994) found transparency depth ranging from 15.0 to 74.0 cm in polyculture ponds. Rahman (1992) concluded that the transparency of productive water bodies should be 40.00 cm or less.

Dissolved Oxygen (DO)

The mean value of dissolved oxygen varied from 5.32 ± 0.09 (T₃) to 5.41 ± 0.09 mg/l (T₁). The higher value of dissolved oxygen was found in T₁ due to comparatively lower stocking density. Talukdar *et al.* (2012) recorded mean value of DO ranging from 6.23 mg/l to 6.63 mg/l while working on suitability of duckweed as feed for fish in polyculture system. Fluctuation of dissolved oxygen concentration might be attributed to photosynthetic activity and variation in the rate of oxygen consumption by fish and other aquatic organisms (Boyd, 1982). The suitable range of dissolved oxygen is 5-8 mg/l for fish culture (Boyd, 1998).

Hydrogen ion concentration (pH)

The mean values of pH varied from 7.25 ± 0.07 (T₂) to 7.35 ± 0.09 (T₁). Slight alkaline condition (7.35) was good for fish culture. The range of pH obtained in the resent study is suitable for fish culture which is agreed with Boyd (1998). Swingle (1967) considered pH values of 6.5 to 9.0 as satisfactory level for fish culture. The findings of the present study also agree well with the observation of Talukdar *et al.* (2012) who recorded the pH value as 7.45 ± 0.21 while working on suitability of duckweed as fish feed in polyculture system. Majhi *et al.* (2006) reported pH 7.5 – 8.2 in two ponds were fed with organically produced Azolla.

Total alkalinity

The recorded mean total alkalinity varied from 110.81 ± 0.85 (T₃) to 112.56 ± 0.94 mg/l (T₁). It is desirable to have an alkalinity of above 20 mg/l for optimal fish production (BAFRU, 1990). Alkalinity has direct effect on primary productivity. In low alkalinity aquatic environments certain nutrients are unavailable to aquatic life (Wurts and Durbrow, 1992). Mairs (1996) considered a total alkalinity of 40.0 mg/l or more to be productive than water bodies with lower alkalinity. Ferdoushi *et al.* (2008) recorded total alkalinity as 133.87 ± 18.95 mg/l in Azolla fed fish pond. In another study, Kabir *et al.* (2009) recorded alkalinity value ranged from 61 to 97.5 mg/l in duckweed fed polyculture pond.

Free Carbon-dioxide (CO₂)

Results from the study indicated that the value of CO_2 varied from 2.64±0.07 (T₁) to 2.87±0.08 mg/l (T₃) was not harmful for fish culture. This assumption was supported by Boyd (1998). According to Moriarty (1997), biochemical pathway of organic matter decomposition requires oxygen. Carbon-dioxide is produced by this process makes the water more acidic, which resulted in lower pH and lower alkalinity value. Higher photosynthesis might be resulted in higher DO and lower free CO_2 concentration in the water column. Talukdar *et al.* (2012) in their study on suitability of duckweed as fish feed recorded free CO_2 value of 2.85±0.30 mg/l which was also strongly agreed with the findings of the present study.

5.4.2: Growth of fishes

5.4.2.1: Monthly variations

Specific growth rate (SGR, %, bwd⁻¹)

The specific growth rate (%, bwd⁻¹) varied with *H. molitrix* from 0.53 ± 0.01 (T₂ at 6th month) to 2.68 ± 0.02 (T₁ at 1st month), *C. catla* from 0.48 ± 0.003 (T₃ at 6th month) to 2.63 ± 0.003 (T₁ at 1st month), *L. rohita* from 0.39 ± 0.01 (T₂ at 6th month) to 2.04 ± 0.07 (T₁ at 1st month), *C. cirrhosus* from 0.39 ± 0.02 (T₂ at 6th month) to 2.30 ± 0.11 (T₂ at 1st month), *C. carpio* from 0.50 ± 0.03 (T₃ at 6th month) to 2.28 ± 0.16 (T₁ at 1st month), *C. idella* from 0.49 ± 0.01 (T₂ at 6th month)

to 2.98±0.04 (T₁ at 1st month) and *B. gonionotus* from 0.46±0.02 (T₃ at 6th month) to 3.42±0.07 (T₁ at 1st month) . Among the species, the highest SGR was observed in *B. gonionotus* as 3.42 ± 0.07 %, bwd⁻¹ and the lowest was in *L. rohita* as 0.39 ± 0.01 %, bwd⁻¹. *B. gonionotus* showed highest SGR, possibly due to the availability of macrophytes. Talukdar *et al.* (2012) reported the highest SGR for *B. gonionotus* among the carp species in duckweed fed ponds. Lowest SGR in *L. rohita* might be due to the fact that indigenous rohu (*Labeo rohita*) is not affected by duckweed (Azim and Wahab, 2003). Grass carp showed moderately higher SGR, possibly due to the availability of macrophytes (Roy *et al.* 2002).

The lowest combined SGR value was found as 0.47 ± 0.022 (T₂ at 6th month) and the highest combined SGR was found as 2.60 ± 0.18 (T₁ at 1st month). The highest SGR value was found in the month of April and the lowest value was found in the month of September for all the species. The lowest SGR value in the month of September might be due to the slow growth rate at the starting of winter season. Slow growth due to slow metabolic activity of fish is found in lower water temperature (Boyd, 1998).

Monthly weight gain (g/month)

The weight gain (g/month) varied with *H. molitrix* from 66.33 ± 0.88 g (T₃ at 1st month) to 84.00 ± 1.53 g (T₁ at 4th month), *C. catla* from 68.33 ± 0.88 g (T₃ at 1st month) to 94.33 ± 1.76 g (T₁ at 3rd month), *L. rohita* from 38.67 ± 1.33 g (T₃ at 5th month) to 59.67 ± 0.88 g (T₁ at 4th month), *C. cirrhosus* from 42.67 ± 2.33 g (T₃ at 5th month) to 73.33 ± 2.40 g (T₁ at 4th month), *C. carpio* from 51.00 ± 2.08 (T₂ at 1st month) to 91.67 ± 3.28 g (T₁ at 3rd month), *C. idella* from 89.00 ± 3.21 (T₃ at 5th month) to 115.67 ± 1.76 g (T₁ at 3rd month) and *B. gonionotus* from 32.33 ± 1.33 (T₃ at 6th month) to 53.33 ± 0.33 g (T₁ at 4th month). Among the species, the highest monthly weight gain was observed in *C. idella* as 115.67 ± 1.76 g due to herbivorous nature and the lowest was in *B. gonionotus* as 32.33 ± 1.33 g which might be due to slow growth pattern. Azad *et al.* (2004) reported the weight gain for *H. molitrix* as 91.78 and *C. cirrhosus* as 85.75g in carp polyculture system which were more or less closer to the present findings.

The lowest combined weight gain value was found as 56.81 ± 7.29 (T₃ at 1st month) and the highest combined weight was found as 80.62 ± 7.64 (T₁ at 4th month). The highest weight gain was found in the month of July and the lowest value was found in the month of April for all the species. Most of the species gains highest weight gain in 3rd or 4th month which might be due to the favourable temperature during this study (Appendix-2). The low growth rate of fishes in treatment T₃ appeared to be related with higher densities and increased competition for food and space (Haque *et al.*, 1994; Islam, 2002).

Mean variations

Specific growth rate (%bwd⁻¹)

The mean specific growth rate varied with *H. molitrix* from 1.16 ± 0.30 (T₃) to 1.23 ± 0.32 (T₁), *C. catla* from 1.16 ± 0.29 g (T₃) to 1.21 ± 0.32 (T₁), *L. rohita* from 0.96 ± 0.21 (T₃) to 1.01 ± 0.25 (T₁), *C. cirrhosus* from 0.98 ± 0.23 (T₃) to 1.10 ± 0.26 (T₁), *C. carpio* from 1.14 ± 0.25 (T₃) to 1.21 ± 0.28 (T₁), *C. idella* from 1.21 ± 0.34 (T₃) to 1.28 ± 0.38 (T₁) and *B. gonionotus* from 1.35 ± 0.39 (T₃) to 1.43 ± 0.44 (T₁). Kabir *et al.* (2009) reported the SGR value of *H. molitrix*, *C. cirrhosus*, *B. gonionotus* and *C. carpio* as 1.38, 1.25, 1.24 and 1.34%, respectively in duckweed fed fish culture ponds.

Specific growth rate of exotic carps, viz., grass carp, silver carp, common carp and Thai sharpunti showed higher values than those of Indian major carps, viz., catla, rohu and mrigal. It is due to the fact that silver carp (*H. molitrix*) is recognized for its rapid growth rate, common carp (*C. carpio*) shows better growth rte over mrigal (*C. cirrhosus*) and Thai punti (*B. gonionotus*) is primarily macrophyte feeder (Morrice, 1998). Further, among the exotic species, *B. gonionotus* showed the highest specific growth rate $(1.35\pm0.39-1.43\pm0.44\%)$ than those of the other two, attributed to its inherent higher growth potential and taking Azolla as feed. Higher growth rates of all the exotic species over Indian major carps in the present study are in agreement with the observations by earlier workers (Lakshmanan *et al.*, 1971; Chaudhuri *et al.*, 1974, 1975, 1978; Sinha and Gupta, 1975; Mahaboob and Sheri, 1997; Tripathi *et al.*, 2000). Combined specific growth rate (SGR, %bwd⁻¹) of all fishes was significantly highest in treatment T₁ where the stocking density was lowest as compared to those of treatments T₂ and T₃ although the same food (Azolla) was supplied in all the treatments. Low growth rate of fishes in treatments T₂ and T₃ appeared to be related with higher densities and increased competition for food and space. A similar experience with regard to growth, i.e. retardation of fish growth at higher densities, was observed by Lakshmanan *et al.* (1971) and Jena *et al.* (1998) while working with carps and other fish species. Better growth in lower density than higher density also noted by Papoutsoqlou *et al.* (1987).

Weight gain (g/month)

The mean weight gain (g) varied with *H. molitrix* from 71.33±1.18 g (T₃) to 80.83 ± 1.55 g (T₁), *C. catla* from 76.00±3.62 g (T₃) to 84.72 ± 2.78 g (T₁), *L. rohita* from 45.22±2.73 g (T₃) to 50.22 ± 2.89 g (T₁), *C. cirrhosus* from 46.78±1.96 g (T₃) to 60.72 ± 3.49 g (T₁), *C. carpio* from 68.56 ± 3.71 g (T₃) to 79.61 ± 4.65 g (T₁), *C. idella* from 91.61±1.09 g (T₃) to 105.28 ± 2.88 g (T₁) and *B. gonionotus* from 38.28 ± 2.75 g (T₃) to 44.06 ± 2.64 g (T₁). Talukdar *et al.* (2012) found similar weight gain in duckweed fed fish culture ponds as 65.86 g (*C. cirrhosus*) and 35.34 g (*B. gonionotus*). The individual weight gain of all the species were found lower in ponds with high stocking density (T₃), which might be due to high competition for feed and space that retard the weight gain.

Among the three Indian major carp species catla showed higher weight gain over, mrigal and rohu in all the treatments, which might be due to the fast growth rate of catla (*C. catla*) over mrigal (*C. cirrhosus*) and rohu (*L. rohita*). Morrice (1998) observed similar trend in weight gain for catla. Among the exotic species the highest weight gain was observed as 105.28 ± 2.88 g in *C. idella* and the lowest was in *B. gonionotus* as 38.28 ± 2.75 g.

The combined mean weight gain (g) of all the species were found to be higher in treatment T_1 over T_2 and T_3 , due to low biomass contained at lower stocking densities of T_1 and other associated favourable environmental parameters compared to the treatments with higher densities. A similar experience with regard to growth, i.e. retardation of fish growth at higher densities, was observed by Jhingran (1991) while working with carps and other fish species.

Final Weight (g)

The final weight (g) after 6 months growing period with *H. molitrix* varied from 468.00±24.01 (T₃) to 545.00±9.61 g (T₁), *C. catla* from 521.00±6.25 (T₃) to 573.33±3.71 g (T₁), *L. rohita* from 329.33±3.71 (T₃) to 359.33±3.18 g (T₁), *C. cirrhosus* from 338.67±10.17 (T₃) to 422.33±11.61 g (T₁), *C carpio* from 472.33±3.71 (T₃) to 538.67±3.84 g (T₁), *C. idella* from 619.67±16.48 (T₃) to 701.67±7.22 g (T₁) and *B. gonionotus* from 251.33±4.06 (T₃) to 287.67±1.67 g (T₁). Kabir *et al.* (2009) worked on use of duckweed as feed for fishes in polyculture system and found final weight of *H. molitrix*, *C. cirrhosus*, *B. gonionotus* and *C. carpio* as 300, 210, 162.50 and 211.20 g/6 months, respectively which were lower than the present findings, due to stocking small sized fingerlings and shorter culture period. *C carpio* grows better than *C. cirrhosus*. Wahab *et al.* (2002) observed that common carp (*C. carpio*) grows better than its counter-part native species mrigal and does not exert significant antagonistic impact on other native carp species in polyculture.

Among the species the highest final weight was observed in as 701.67 ± 7.22 g (T₁) in *C. idella* and the lowest was in *B. gonionotus* as 251.33 ± 4.06 (T₃). Azolla is preferred by grass carp because of its low content of fibre and fat (Mandal *et al.* (2010). Talukdar *et al.* (2012), Morrice (1998) and Jena *et al.* (2002b) reported the highest final weight of grass carp (*C. idella*) among carp species, fed with duckweed. Jena *et al.* (1998) mentioned that Thai punti (*B. gonionotus*) is a medium sized carp species and reported to have moderate growth rate compared to the major carps. Chaudhuri *et al.* (1974, 1975), Sinha *et al.* (1973) and Sinha and Saha (1980) recorded similar higher growth rates of grass carp over other species.

The lowest mean final weight was found as 2167.67 ± 17.31 g (T₃) and the highest final weight as 3629.34 ± 19.52 g (T₁) for all the species. This might be due to direct impact of stocking density. Similar argument is supported by Jena

et al. (1998) while working with the culture of Indian major carps under different stocking density.

Survival rate (%)

The mean survival rate with *H. molitrix* varied from 85.33 ± 1.17 (T₃) to 90.33±2.91% (T₁), C. catla from 80.87±0.58 (T₃) to 83.63±0.75% (T₁), L. rohita from 83.10±2.18 (T1) to 84.13±1.21% (T2), C. cirrhosus from 85.50±0.58 (T2) to 91.53±1.13% (T1), C. carpio from 93.07±0.81% (T2) to 94.53±0.61% (T1), C. idella from 85.07±0.96 (T₂) to 86.37±0.52% (T₁) and B. gonionotus from 80.50±1.15 (T₁) to 81.73±0.79% (T₃). Talukdar *et al.* (2012) found survival rate as 82% (C. catla), 88% (B. gonionotus) in the weed based carp polyculture system. In a yearlong grow-out carp polyculture trial with olive barb as a component species at different species combinations and ratios, Chakraborty et al. (2005) in Bangladesh observed 75.5-78.6% survival in silver carp and catla, 73.2-82.4% in rohu and 65.2-75.4% in mrigal. Similar survival rate of silver carp followed by rohu, mrigal and catla have also been reported earlier in some grow-out carp polyculture trials (Tripathi et al., 2000; Jena et al., 2002a). In contrast, we recorded higher survivals in all the species, attributed to the relatively larger stocking size and lower stocking density (except B. gonionotus). The lowest mean survival rate was found as 85.20±1.55 % (T₃) and the highest survival rate as 87.14±1.94 % (T₁) for all the species .Variation in stocking density of fish may change growth and survival rates (Miao, 1992).

Yield (Kg/ha/6 months)

The yield (Kg/ha/6 months) with *H. molitrix* varied from 969.00 ± 17.16 (T₁) to 982.67 ± 50.35 (T₃), *C. catla* from 241.00 ± 1.53 (T₁) to 411.67 ± 4.81 (T₃), *L. rohita* from 603.67 ± 5.21 (T₁) to 781.00 ± 8.62 (T₃), *C. cirrhosus* from 375.67 ± 10.17 (T₁) to 426.33 ± 12.84 (T₃), *C. carpio* from 252.67 ± 1.86 (T₁) to 419.67 ± 3.38 (T₃), *C. idella* from 520.33 ± 13.87 (T₃) to 884.00 ± 9.07 (T₁) and *B. goninotus* from 496.67 ± 8.11 (T₃) to 568.33 ± 3.33 (T₁). Azad *et al.* (2004) reported that Azolla for *C. idella* show encouraging result.

In treatment T₁, grass carp (*C. idella*) contributed 22.70% of total fish production and Thai punti (*B. gonionotus*) contributed 14.59% of total fish production. In treatment T₂, grass carp (*C. idella*) contributed 17.30% of total fish production and Thai punti (*B. gonionotus*) contributed 13.34% of total fish production. In treatment T₃, grass carp (*C. idella*) contributed 12.88% of total fish production and Thai punti (*B. gonionotus*) contributed 12.88% of total fish production and Thai punti (*B. gonionotus*) contributed 12.30% of total fish production. Macrophytophagus species (*C. idella* and *B. gonionotus*) contributed 37.29%, 30.64% and 25.18% in treatment T₁, T₂ and T₃, respectively. In addition, it is assumed that the uneaten dead Azolla along with fish faeces settled on to the bottom and consumed directly by common carp and supplied nutrients for phytoplankton.

The highest combined yield (4038.33±84.41 kg/ha/6 months) was obtained from treatment T₃ stocked with high density of fish. The increased growth and production with higher density of fish might have been due to the proper stocking density, synergistic interaction and confounding effects of additional numbers of silver, rui, catla, mrigal and carpio. Significant difference was found among the treatments. The net production of this experiment is comparable to other reported productions in semi-intensive polyculture in Bangladesh. Gopakumar *et al.* (1999) mentioned that the production of weed based carp polyculture was 3-4 tonnes/ha/year which was lower than the production of the present study. It might be due to the good water quality and regular supply of Azolla. Grover *et al.* (2000) mentioned that the possible production of Azolla based carp polyculture was 5575 kg/ha which was more or less closer to the findings of the present study. Jena *et al.* (2002b) also reported the total production as 3472 kg/ha in 6 months in carp polyculture system.

Mazid *et al.* (1997) recorded gross fish production of 2545–3688 kg/ ha during 330-day culture period, the highest production with artificial feed using 53% rice bran, 30% mustard oil cake, 10% fish meal and 6% molasses and the lowest with the traditional 75% rice bran and 25% mustard oil cake. Kohinoor *et al.* (1999) reported a gross fish production of 2056 kg/ha in ponds supplied with

159

only duckweed and 2566 kg/ha in ponds supplied with only rice bran during a 180-day culture period.

5.4.3: Economics

Along with the increase in production, the purpose of aquaculture practices is to earn a profit. Wyban *et al.* (1988) indicated that stocking density, growth rate, survival and market price are the most sensitive factors to increase profit. Total investment cost and total return were directly related to stocking density. The highest density required the highest total investment cost and also provides the highest return.

During study period total cost, gross benefit, net benefit and CBR varied from 136862.00 \pm 606.45 Tk/ha/6 months (T₁) to 145754.00 \pm 57.35 Tk/ha/6 months (T₃), 384562.35 \pm 1890.32 (T₁) to 402667.35 \pm 8862.88 Tk/ha/6 months (T₃), 247700.35 \pm 1890.32 (T₁) to 256913.35 \pm 8862.88 Tk/ha/6 months (T₃) and 1.76 \pm 0.06 (T₃) to 1.81 \pm 0.02 (T₁), respectively.

Total cost was higher in Treatment T_3 might be due to the higher stocking density i.e. seed cost (24.54% of total cost). Roy et al. (2003) studied economics of carp-SIS polyculture and their economic analysis under 3 treatments and they mentioned the operational cost was 32,450.00, 39,950.00 and 42,450.00 Tk./ha/7 months in treatments T_1 , T_2 and T_3 , respectively which was lower than the present study might due to comparatively smaller size of stocked fish seed. In another Grover et al. (2000) described total cost as 85,000.00 BDT/ha which more or less similar to the present study. Gross benefit was varied from 384562.35±1890.32 BDT/ha (T1) to 402667.35±8862.88 BDT/ha (T3) which were higher than the findings of Roy et al. (2003) who reported the gross benefit was 128,000.00, 128,280.00 and 110,720.00 Tk./ha/7 months in treatments T₁, T₂, and T_3 , respectively because of lower production rate than the present study. In case of net benefit, Majhi et al. (2006) reported the net benefit of 227338.00 BDT/ha was obtained from Azolla fed pond which was slightly lower than the present findings. Shanmugasundaram and Balusamy (1993) stated that benefit cost ratio was 1.88 which was similar to the present study. Khan et al. (2009) also mentioned BCR value of 1.22 which was lower than the findings of the

present study was due to high initial biomass of all the species and also the higher survival rate.

Inspite of having lowest total fish production with treatment T_1 best growth performance of macrophytophagus species was found with that treatment. This indicated that treatment T_1 was found most potential for the development of low cost fish farming which was reflected with having significantly lowest total cost and highest CBR with that treatment. The study indicated that Azolla based carp polyculture with stocking density of 9880 fish/ha could be used for low cost weed based fish farming in Bangladesh.
5.5: Conclusion

Considering the water quality, production and economics, it can be concluded that Azolla based carp polyculture with stocking density of 9880 fish/ha be used for low cost Azolla based fish farming in Bangladesh.

5.6: Recommendation

In this study Azolla based carp polyculture with stocking density of 9880 fish/ha was found more profitable. Potentials of Azolla for farming of macrophytophagus fishes other than carps were not included in this study. Studies on Azolla based farming for macrophytophagus fishes other than carps are recommended as future step.

CHAPTER SIX

Summary, Conclusion and Recommendation

Weed based aquaculture has been demonstrated to be one of the easiest method of fish production requiring less capital which improves sustainability, productivity and profitability. The aim of the present study was to explore the potentials of weed based aquaculture as a low cost technology in ponds. A total of 3 experiments were conducted in 3 separate years (2010, 2011 and 2012). Experiment-1 was conducted at the Protein and Enzyme Research Laboratory, Department of Biochemistry and Molecular Biology, Rajshahi University, Rajshahi whereas experiment-2 and experiment-3 were conducted at Alampur village of Sadar Upazila under Kushtia district, Bangladesh. In experiment-1, six different types of conventional and non-conventional fish feed items like rice bran, wheat bran, mustard oilcake, Azolla, grass and banana leaves were tested to determine the nutrient contents (protein, lipid and carbohydrate) under 6 treatments as T₁, T₂, T₃, T₄, T₅ and T₆, respectively. Considering the nutrient content in experiment-1, experiment-2 evaluated the fish production and economics under 4 treatments of feed and weed based system with a similar stocking density of 11115 fish/ha in polyculture ponds as T₀: ponds fed with conventional feed like rice bran, wheat bran and mustard oilcake (no weed was used as fish feed), T1: Azolla (Azolla pinnata) fed ponds, T2: Grass (Cynodon *dactylon*) fed ponds and T₃: Banana (*Musa acuminata*) leaf fed ponds. Based on the performance of weeds and feeds in experiment-2, experiment-3 optimized the stocking density for Azolla based carp polyculture system in ponds under 3 different treatments of stocking densities like T1: 9880 fish/ha, T₂: 11115 fish/ha and T₃: 12350 fish/ha.

Nutrients in weeds and feeds were compared for a period of 6 months (April to September) in experiment-1. Fishes (*Hypophthalmichthyes molitrix, Catla catla, Labeo rohita, Cirrhinus cirrhosus, Cyprinus carpio, Ctenopharyngodon idella* and *Barbonymus gonionotus*) were also grown for a period of 6 months (April to September) in experiment-2 and experiment-3. Mean initial stocking weight of *H. molitrix, C. catla, L. rohita, C. cirrhosus, C. carpio, C. idella* and *B. gonionotus* were 62, 64, 57, 54, 63, 65 and 25 g, respectively for experiment-2

and 60, 65, 58, 52, 61, 70 and 22 g, respectively for experiment-3. There were 3 replications for each treatment under different experiments.

Liming (250 kg/ha) and basal fertilization (cowdung: 1500 kg/ha, urea: 60 kg/ha and TSP: 60 kg/ha) were done for all the treatments under experiment-2 and experiment-3. Urea (2.5 kg/ha/day in all treatments except T₁ under experiment-2 and in no treatment under experiment-3) and TSP (2.5 kg/ha/day in all treatments under experiment-3) were applied as periodic fertilization.

In case of experiment-1, nutrient contents (protein, lipid and carbohydrate) were monitored monthly whereas in case of experiment-2 and experiment-3 water quality parameters (water temperature, transparency, DO, pH, alkalinity and free CO₂) were monitored fortnightly and fish growth parameters (weight gain and SGR) were monitored monthly. Economics (in terms of total cost, gross benefit, net profit margin and CBR) of fish farming were also evaluated for both the experiment-2 and experiment-3.

In experiment-1, significant variations were found in the mean values of nutrient contents with different treatments of feed items but in case of same feed item no significant difference was found in the nutrient content at different months. Among the non-conventional feed items treatment T₄ (Azolla) varied more significantly for the mean values of protein content.

In experiment-2, the mean values of water temperature, transparency, DO, pH, total alkalinity and free CO₂ varied from 31.19 ± 0.20 (T₃) to 31.29 ± 0.24 °C (T₀), 32.28 ± 0.32 (T₂) to 32.83 ± 0.40 cm (T₁), 5.33 ± 0.09 (T₃) to 5.51 ± 0.11 (T₁) mg/l, 7.18 ± 0.03 (T₃) to 7.38 ± 0.05 (T₁), 113.28 ± 0.85 (T₁) to 114.36 ± 0.72 (T₀) mg/l, and 2.82 ± 0.07 (T₁) to 3.03 ± 0.07 (T₀), respectively. Considering all species, mean values of SGR (%, bwd⁻¹), monthly weight gain (g), final weight (g), survival rate (%) and yield (Kg/ha/6 months) of fishes significantly varied from 0.93 ± 0.03 (T₃) to 1.17 ± 0.05 (T₀), 42.33 ± 6.71 (T₃) to 76.64 ± 9.24 (T₀), 2167.67 ± 17.31 (T₃) to 3629.34 ± 19.52 (T₀), 76.33 ± 1.74 (T₂) to 79.52 ± 2.20 (T₀) and 2541.00 ± 0.67 (T₃) to 4403.51 ± 0.88 (T₀), respectively. Total cost (BDT/ha), gross benefit (BDT/ha), net benefit (BDT/ha), net profit margin (%) and CBR

significantly varied from 123430.50 ± 0.00 (T₁, T₂, T₃) to 235930.50 ± 0.00 (T₀), 235068.40 ± 1965.31 (T₃) to 418376.85 ± 5125.59 (T₀). 111639.90 ± 2056.87 (T₃) to 206744.85 ± 3221.73 (T₁), 77.21 ± 2.40 (T₀) to 167.20 ± 18.77 (T₁) and 0.77 ± 0.02 (T₀) to 1.67 ± 0.18 (T₁), respectively. Findings indicated that treatment T₁ (Azolla based carp polyculture) was more profitable than that of other treatments.

In experiment-3, the mean values of water temperature, transparency, DO, pH, total alkalinity and free CO₂ varied from 31.76±0.30 (T₃) to 31.93±0.27°C (T₁), 32.25 ± 0.34 (T₂) to 33.11 ± 0.33 cm (T₁), 5.32 ± 0.09 (T₃) to 5.41 ± 0.09 mg/l (T₁), 7.25 \pm 0.07 (T₂) to 7.35 \pm 0.09 (T₁), 110.81 \pm 0.85 (T₃) to 112.56 \pm 0.94 mg/l (T₁), and 2.64 \pm 0.07 (T₁) to 2.87 \pm 0.08 mg/l (T₃), respectively. Considering all species, mean SGR (%, bwd⁻¹), monthly weight gain (g), final weight (g), survival rate (%) and yield (Kg/ha/6 months) of fishes significantly varied from 1.14 ± 0.05 (T₃) to 1.21±0.05 (T₁), 62.54±7.36 (T₃) to 72.21±8.15 (T₁), 3000.33±31.25 (T₃) to 3428.00±25.35 (T₁), 85.20±1.55 (T₃) to 87.14±1.94 (T₁) and 3894.33±18.00 (T₁) to 4038.33±84.41 (T₃), respectively. Total cost (BDT/ha), gross benefit (BDT/ha), net benefit (BDT/ha), net profit margin (%) and CBR significantly varied from 136862.00±606.45 (T_1) to 145754.00±57.35 (T₃). 384562.35±1890.32 (T1) to 402667±8862.88 (T3), 247700.35±1890.32 (T1) to 256913.35±8862.88 (T₃), 176.31±5.55 (T₃) to 181.00±1.53 (T₁) and 1.76±0.06 (T₃) to 1.81±0.02 (T₁), respectively. Inspite of having lowest total fish production with treatment T₁ best growth performance of macrophytophagus species was found with that treatment. This indicated that treatment T1 was found most potential for the development of low cost fish farming which was reflected with having significantly lowest total cost and highest CBR with that treatment. The study indicated that Azolla based carp polyculture with stocking density of 9880 fish/ha could be used for low cost weed based fish farming in Bangladesh.

In this study Azolla based carp polyculture with a stocking density of 9880 fish/ha was found more profitable. Potentials of Azolla for farming of macrophytophagus fishes other than carps were not included in this study. Studies on Azolla based farming for macrophytophagus fishes other than carps are recommended as future step.

165

REFERENCES

- Abdelghany, A. E. and Ahmad, M. H. 2002. Effects of feeding rates on growth and production of Nile tilapia, common carp and silver carp polycultured in fertilized ponds. *Aquaculture Research*, **33**: 415-423.
- Abou, Y., Fiogbe, E. D. and Micha, J. 2007. Effects of stocking density on growth, profitability of farming Nile tilapia, *Oreochromis niloticus* L., fed *Azolla* diet, in earthen ponds. *Aquaculture Research*, **38**: 595-604.
- ADB, 2005. An evaluation of small-scale freshwater rural aquaculture development for poverty reduction. Operations Evaluation Department, Asian Development Bank (ADB), Asian Development Bank Publications, ISBN 071-561-550-3, Manila, Philippines. 164p.
- Ahmed, F. 2011. Use of *Azolla* as fish feed and poultry feed, an alternative use of urea. *Farmhouse*, p. 32.
- Ahmed, M., Rab, M. A. and Gupta, M. V. 1995. Impact if improved aquaculture technologies: Results of an extension program on the farming systems of Bangladesh. *Asian Fisheries Science*, 8(1): 27-39.
- Alaerts, G. J., Rahman, M. M. and Kelderman, P., 1996. Performance analysis of a full-scale duckweed-covered sewage lagoon. *Water Res.*, **30**: 843– 852.
- Alalade, O. A. and Iyayi, E. A. 2006. Chemical composition and the feeding value of Azolla (*Azolla pinnata*) meal for egg-type chicks. *International Journal of Poultry Science*, 5(2): 137-141.
- Alam, F. M. and Thomson, K. J. 2001. Current constraints and future possibilities for Bangladesh fisheries. *Food Policy*, **26**: 297-313.
- Ali, M. Z., Zaher, M., Parveen, R., Hussain, M. G. and Mazid, M. A. 2006. The use of alternate feeding schedules on the growth performance and reduction of cost of production for thai pungas, *Pangasius hypophthalmus* (Sauvage, 1978) in ponds. *J. Aquacult. Trop.*, **21**: 149-156.

- Alikhuni, K. H. 1957. Fish culture in India. *F.M. Bull. Indian Coun. Agric. Res.,* **20**: 144.
- Ansal, M. D. and Dhawan, A. 2007. Spirodela for low cost carp feed formulation. Abstract In: 8th Asian Fisheries Forum, Kochi, India, November, 20-23, 2007, p. 164.
- Antonie, T., Wery, P., Micha, J. C. and Van-Hove, C. 1987. Comparison of the growth and chemical composition of *Oreochromis* (*Tilapia*) *niloticus* and *Cichlasoma* (*Theraps*) *melanurum* fed with *Azolla*. *Aquaculture*, **66**: 181-196.
- Ayyappan, S., Dash, B., Pani, K. C. and Tripathi, S. D. 1993. Azolla a new aquaculture input, National meet on Aquafarming system practices and potential, Abstract No. OFS-2, 10-11 February, 1993, CIFA, Bhubaneswar, India.
- Azad, M. A. K., Rahman, M. R., Rahman, Z., Kader, M. A., Haque, M. M. and Alam, M. J. 2004. Polyculture of carp, tilapia and pangas using low cost inputs. *Pak. J. Biol. Sci.*, 7(11): 1918-1926.
- Azim, M. E. and Wahab, M. A. 2003. Development of a duckweed-fed carp polyculture system in Bangladesh. *Aquaculture*, **18**(1-4): 425-438.
- Azim, M. E., Talukder, G. S., Wahab, M. A., Haque, M. M. and Haq, M. S. 1995. Effect of liming and maintenance of total hardness levels on fish production in fertilized ponds. *Progress. Agric.*, 6(2): 7-14.
- Azim, M. E., Wahab, M. A., Kamal, A. H. M. and Zoarder, F. A. 2004. Feeding relations of silver barb *Barbodes gonionotus* (Bleeker) with major Indian and common carp and its effect on fish production in a polyculture system. *Journal of the World Aquaculture Society*, **35**(1): 100-108.
- Backiel, T. and LeCren, E. D. 1978. Some density relationship for fish population parameters. In: S. D. Gerking (ed.) *Ecology of freshwater fish production*. Blackwell Scientific publication Oxford. pp. 279-302.
- BAFRU (Bangladesh Aquaculture and Fisheries Resource Unit), 1990. Water quality management for Aquaculture and Fisheries. Institute of Aquaculture, University of Stirling, UK. p. 49.

- Bag, M. P., Mahapatra, S. C., Rao, P. S. and Chakrabarty, D. 2011. Making aquatic weed as potential feed for Nile tilapia (*Oreochromis niloticus* L.) and its impact on fatty acid profile. *International Research of Pharmacy and Pharmacology*, 1(8):194-202.
- Banerjee, A. and Matai, S. 1990. Composition of Indian aquatic plants in relation to utilization as animal forage. *J. Aquat. Plant Manage.*, **28**: 69-73.
- Banforth, S. 1958. Ecological studies on the planktonic protozoa of a small artificial pond. *Limnol. Oceanogr.*, **3**: 398-412.
- Bardach, J. E., Ryther, J. H. and MeLarney, W. O. 1972. Aquaculture: The Farming and Husbandry of Freshwater and Marine Organisms. Wiley-Interscience, New York. pp. 256-289.
- Becking, J. H. 1979. Environmental requirements of *Azolla* for use in tropical rice production. pp. 345-373. In: *Nitrogen and Rice*, Los Banos, Laguna, International Rice Research Institute.
- Bergins, B. 1949. Zur. Limnologe der seen Sudostlentdlend, Schweig. 2. *Hydrobiologica*, **11**: 583-607.
- Bligh, E. G. and Dyer, W. 1989. Total lipid extraction and purification., *Can. Jour. Biochem. Physiol*, **37**: 911.
- Boeck, A. 1996. Feeding your fish. International Centre for Aquaculture and Aquatic Environments. Swingle Hall, Auburn University, Alabama, USA. p.12.
- Boel, E., Huge-Jensen, B., Christensen, M., Thim, L. and Fill, N. P. 1988. *Lipids*: 701.
- Boyd, C. E. 1968. Fresh water plants: a potential source of protein. *Econ. Bot.*, **22**: 359-368.
- Boyd, C. E. 1969. The nutritive value of three species of water weeds. *Econ. Bot.*, **23**: 123-127.
- Boyd, C. E. 1982. *Water quality management of pond fish culture*. Elsevier Sci. Pub. Co. Amsterdam-Oxford, New York, pp. 318.

- Boyd, C. E. 1990. *Water quality for fish pond for aquaculture.* Birmingham Publishing Co., Birmingham, Alabama, USA, p. 477.
- Boyd, C. E. 1998. *Water quality for fish pond*. Aquaculture Research and Development Series No. 43. Aburn University, Alabama, USA, p. 37.
- Boyd, C. E. and Zimmermann, S. 2000. *Grow-out systems-water quality and soil management*. Blackwell Science. Oxford, Uk. pp. 221-238.
- Brett, J. R. 1979. Factors affecting fish growth. In: Haor, W.S., D.J. Randall and J.R. Brett (eds), *Fish physiology*, Vol. 8. Academic press, New York. pp. 599-675.
- Brown, M. E. 1957. Experimental studies on growth. In: *The physiology of fishes*. Brown M.E. (ed.), Vol. 1, Academic Press, New York, pp. 361-400.
- Buttner, J. K. 1993. An introduction to water chemistry in freshwater aquaculture. NRAC Fact Sheet No. 170-1993. p. 4.
- Cagauan, A. G., Branckaert, R. D. and Van-Hove, C. 2000. Integrating fish and *Azolla* into rice-duck farming in Asia. *Naga*, *The ICLARM Quarterly*.
 23(1): 4-10.
- Cassani, J. R. 1981. Feeding behaviour of underyearling hybrids of grass carp, *Ctenopharyngodon idella* and the bighead, *Hypophthalmichthys nobilis* on selected species of aquatic plants. *J. Fish Biol.*, **18**: 127-133.
- Chakraborty, B. K., Miah, M. I., Mirza, M. J. A. and Habib, M. A. B. 2005. Growth, yield and returns to *Puntius sarana* (Hamilton) Sarpunti, in Bangladesh under semi-intensive Aquaculture. *Asian Fish. Sci.*, **18**: 307–322.
- Chaudhuri, H., Chakrabarty, R. D., Rao, N. G. S., Janakiram, K., Chatterjee, D.
 K. and Jena, S. 1974. Record of fish production with intensive culture of Indian and exotic carps. *Curr. Sci.*, **43**: 303–304.
- Chaudhuri, H., Chakrabarty, R. D., Sen, P. R., Rao, N. G. S. and Jena, S. 1975. A new high in fish production in India with record yields by composite fish culture in freshwater ponds. *Aquaculture*, **6**: 343–355.

- Chaudhuri, H., Rao, N. G. S., Rout, M. and Kanaujia, D. R. 1978. Record of fish production through intensive fish culture. *J. Inland Fish. Soc. India*, **10**: 19–27.
- Chen, T. P. 1976. *Aquaculture practices in Taiwan*. Fishing News (Books) Ltd., West Byfleet, Surrey, England. 124p.
- Chowdhury, A. and Mamun, A. A. 2006. Physiochemical conditions and plankton population of two ponds in Khulna. *Univ. j. zool. Rajshahi Univ.*, 25: 41-44.
- Chowdhury, A. N., Begum, S. and Sultana, N. 1992. Physico-chemical and biological parameters affecting the occurrence and seasonal variation of Ostracods in tropical fish ponds. *Bangladesh J. Zool.*, **20**(2): 315-323.
- Chowdhury, M. M. R., Shahjahan, M., Rahman, M. S. and Islam, M. S. 2008. Duckweed (*Lemna minor*) as supplementary feed in monoculture of nile tilapia, Oreochromis niloticus. Journal of Fisheries and Aquatic Sciences, 3: 54-59.
- Culley, D. D. and Epps, E. A. 1973. Use of duckweed for waste treatment and animal feeds. *J. Wat. Pollut. Control. Fed.*, **45**: 337-347.
- Culley, D. D., Rejmankova, E., Koet, J. and Prye, J. B. 1981. Production, chemical quality and use of duckweeds (Lemnaceae) in aquaculture, waste management and animal feeds. *J. World Maricult. Soc.*, **12**: 27-49.
- Das, A. K. 2000. Limno-chemistry of some Andhra Pradesh reservoirs. *J. Inland Fish. Soc. India*, **32**(2): 37-44.
- Das, B. 1997. *Fisheries and Fisheries Resources Management* Vol. III, Bangla Academy, Dhaka, Bangladesh, pp. 202.
- Das, P., Kumar, D., Bhaumik, U., Roy, B. and Chakraboty, D. P. 1982. Observation on four species culture of Indian major carps and exotic silver carps. *J. Indian Fish. Soc. India*, **141**: 8-22.
- De Silva, S. S. 1992. Fish nutrition research for semi-intensive culture systems in Asia. *Asian Fisheries Science*, **5**: 129-144.

- De Silva, S. S. and Anderson, T. A. 1995. *Fish nutrition in aquaculture.* London, Chapman & Hall, pp. 319.
- Dewan, S., Wahab, M. A., Beveridge, M. C. M., Rahman, M. H. and Sarker, B. K. 1991. Food selection selectivity and dietary overlap among planktivorous Chinese and Indian major carps fry and fingerlings grown in extensively managed, rain-fed ponds in Bangladesh. *Aquacult. and Fish. Management.*, **22**: 277-294.
- Dey, M. M., Bose, M. L. and Alam, M. F. 2008. Recommendation Domains for Pond Aquaculture. Country case study: Development and Status of Freshwater Aquaculture in Bangladesh. WorldFish Center Studies and Reviews No. 1872. The WorldFish Center, Penang, Malaysia. pp.73.
- Dhawan, A. and Kaur, S. 2002a. Effect of pig dung on water quality and polyculture of carp species during winter and summer. *Aquaculture International*, **10**: 297-307.
- Dhawan, A. and Kaur, S. 2002b. Pig dung as pond manure: Effects on water quality, pond productivity and growth of carps in polyculture system. *Naga: The ICLARM Quarterly*, **25**: 11-14.
- DoF (Department of Fisheries), 1998. Pukure Uthpadon Jibbiggan (in Bangla) II Fisheries Training Extension Project (FTEP-II). pp.90.
- DoF (Department of Fisheries), 2003. The Future for Fisheries: Economic Performance. Fisheries Sector Review and Future Development Study. Ministry of Fisheries and Livestock, Government of Peoples Republic of Bangladesh, 172pp.
- DoF, 2010. National Fish Week 2010 Compendium (in Bengali), Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh, 120p.
- DoF, 2011. National Fish Week 2011 Compendium (in Bengali), Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh, 136p.
- DoF, 2012. National Fish Week 2012 Compendium (in Bengali), Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh, 144p.
- DoF, 2013. National Fish Week 2013 Compendium (in Bengali), Department of Fisheries, Ministry of Fisheries and Livestock, Bangladesh, 144p.

- DWRP (Duckweed Research Project), 1996. Socio-economic validation of duckweed based fish production as practiced by PRISM joint stock companies. Vol. No.3. Dhaka, Bangladesh. pp.80.
- DWRP (Duckweed Research Project). 1997. Scientific and technical validation of PRISM duckweed activities. Vol. No.2. Dhaka, Bangladesh. pp.40.
- Ebrahim, M. S. M., Zeinhom, M. M. and Abou-Seif, R. A. 2007. Response of Nile tilapia (*Oreochromis niloticus*) fingerlings to diets containing *Azolla* meal as a source of protein. *Journal of Arabian Aquaculture Society*, 2(1): 54-68.
- Edwards, P. 1980. Food potential of aquatic macrophytes. *ICLARM Studies and Reviews,* **5** : 51.
- Edwards, P. 1987. Use of terrestrial and aquatic macrophytes in aquaculture. In: Moriarty, D. and Pullin, R. S. V., Editors, *Detritus and Microbial Ecology in Aquaculture. ICLARM Conf. Proc.* **14**: 311–385.
- Edwards, P. 1990. Use of terrestrial vegetation and aquatic macrophytes in aquaculture. In: *Detritus and microbial ecology in aquaculture. ICLARM. Conf Proc.* 14. *International Aquat Living Resour Manag Cent, Manila.* pp. 311-335.
- Edwards, P. 1993. Environmental issues in integrated agriculture-aquaculture and waste water-fed fish culture systems. In: *Environment and aquaculture in developing countries* (Eds. Pullin, R. S. V., Rosenthal, H., Maclean, J. L.). *ICLARM conference proceedings 31*. International Center for Living Aquatic Resources Management, Manila, Philippines. pp. 139-170.
- Edwards, P., Kamal, M. and Wee, K. L. 1985. Incorporation of composted and dried water hyacinth in pelleted feed for the tilapia *Oreochromis niloticus*. *Aquaculture and Fisheries Management*, **16**: 233–248.
- Effiong, B. N, Sanni, A. and Fakunle, J. O. 2009. Effect of partial replacement of fish meal with duckweed (*Lemna pauciscostata*) meal on the growth performance of *Heterobranchus longifilis* fingerlings. *Report and opinion*, **1**(3): 76-81.

- El-Shafai, S. A., El-Gohary, F. A., Nasr, F. A., Van der Steen, N. P. and Gijzen,H. J. 2004. Suitability of using duckweeds as feed and treated sewage as water source in tilapia aquaculture. Taylor & Francis, Inc. pp. 65-82.
- FAO (Food and Agriculture Organization), 2006. Livelihood adaptation to climate variability and change in drought- prone areas of Bangladesh. Asian Disaster Preparedness Center, Food and Agriculture Organization of the United Nations, Rome, Italy, pp.1-96.
- FAO (Food and Agriculture Organization), 2008. Fisheries Department. Fishery Information. Data and Statistics Unit. Fish stat plus: Universal software for Fishery statistical times series. Version 2.3.
- FAO (Food and Agriculture Organization), 2012. The State of World Aquaculture and Fisheries. Rome. 209pp.
- Fasakin, E. A. 1999 Nutrient quality of leaf protein concentrates produced from water fern (*Azolla africana*) and duckweed (*Spirodella polyrrhiza*). *Bioresource Technology*, **69**(2): 185-187.
- Fasakin, E. A. and Balogun, A. M. 2001. Nutritional and anti-nutritional analyses of Azolla africana Desv. and Spirodela polyrrhiza L. Schleiden as feedstuffs for fish production. In: 14th Annual Conference of the Fisheries Society of Nigeria (FISON), 19-23 January 1998, Ibadan, Nigeria. pp. 31-39.
- Ferdoushi, Z., Haque, F., Khan, S. and Haque, M. 2008. The effects of two aquatic floating macrophytes (*Lemna* and *Azolla*) as biofilters of nitrogen and phosphate in fish ponds. *Turkish Journal of Fisheries and Aquatic Sciences*, 8: 253-258.
- Gaiger, I. G., Porath, D. and Granoth, G. 1984. Evaluation of duckweed (*Lemna gibba*) as feed for tilapia (*Oreochromis nilotieus* cross *Oreochromis aureus*) in a recirculating unit. *Aquacultre*, **41**: 235-244.
- Gavina, L. D. 1994. Pig-Duck-Fish-Azolla integration L. A. Union Philippines. *Naga The ICLARM Quarterly*, pp. 18-20.
- Gomez, K. A. and Gomez, A. A. 1984. *Statistical Procedure for Agricultural Research*. 2nd Ed. John Wiley & Sons. 697 p.

- Gopakumar, K., Ayyappan, S., Jena, J. K., Sahoo, S. K., Sarkar, S. K., Satapathy, B. B. and Nayak, P. K. 1999. National Freshwater Aquaculture Development Plan. Central Institute of Freshwater Aquaculture, Bhubaneswar, India. 75p.
- Grover, J. H., Islam, M. A., Shah, W. A., Rana, M. A. H. and Chowdhury, H. A. 2000. Training manual for extension personnel on low cost environment friendly sustainable aquaculture practices. ICLARM, Dhaka, Bangladesh. p.101.
- Gupta, M. V., Mazid, M. A., Islam, M. S., Rahman, M. and Hussain, M. G. 1999. Integration of aquaculture into the farming system of the floodprone ecosystem of Bangladesh: an evaluation of adoption and impact. ICLARM Tech. Rep. 56. Manila.
- Halver, J. E. 1972. Fish nutrition. Academic Press, New York. 713p.
- Halver, J. E. 1984. *Special methods in pen fish husbandry*. Akademiai Nyomda, Budapest, 146p.
- Haque, M. M., Khan, S. and Sarkar, M. R. U. 2005. Production and economic return in pangasid catfish (*Pangasius hypophthalmichthys*) monoculture and polyculture with silver carp (*Hypophthalmichthys molitrix*) in farmers ponds. *Bangladesh J. Fish. Res.*, **9**(2): 111-120.
- Haque, M. Z., Rahman, M. A., Hossain, M. M. and Rahman, M. A. 1994. Effect of stocking densities on the growth and survival of mirror carp, *Cyprinus carpio* var. *specularis* in rearing ponds. *Bangladesh J. Zool*, **22**: 109–116.
- Haroon, A. K. Y. and Pittman, K. A. 1997. Rice-fish culture: feeding, growth and yield of two size classes of *Puntius gonionotus* Bleeker and *Oreochromis* spp. in Bangladesh. *Aquaculture*, **154**: 261–281.
- Hasan, M. R., Alam, M. G. M. and Islam, M. A. 1989. Evaluation of some indigenous ingredients as dietary protein sources for the catfish (*Clarias batrachus*, Linnaeus) fry. pp. 125–137. In: Huisman, E. A., Zonneveld, N. and Bouwmans, A. H. M. (eds.). *Aquacultural research in Asia: management techniques and nutrition.* AA Wageningen, The Netherlands.

- Hasan, M. R., Ali, M. A., Akand, A. M. and Ali, M. M. 1996. Nutritional requirements of Indian major carp *Labeo rohita* – 2: Effect of dietary lipid on growth, food conversion and body composition. *Bangladesh J. Aquacult.*, **18**: 24-31.
- Hasan, M. R., Roy, P. K. and Akand, A. M. 1994. Evaluation of leucaena leaf meal as a dietary protein source for Indian major carp, *Labeo rohita* fingerlings. pp. 69–76. In: S.S. De Silva, (ed.) *Fish nutrition research in Asia*. Special Publication No. 9, Manila, Asian Fisheries Society.
- Hassan, M. S. and Edwards, P. 1992, Evaluation of duck weed (*Lemna perpusilla and Spirodela polyrhiza*) as feed for Nile Tilapia (*Oreochromis niloticus*). Aquaculture, **104**: 315-326.
- Hepher, B. 1988. *Nutrition of Pond Fishes.* Cambridge University Press, London, UK., 180p.
- Hepher, B. 1990. *Nutrition of pond fishes*. Cambridge, Cambridge University Press, 388p.
- Hepher, B. and Pruginin, Y. 1979. Guide to fish culture in Israel. 4. Fertilization, manuring and feeding. Foreign Training Dept., Israel, 61pp.
- Hepher, B., Milstein, A., Leventer, H., and Teltsch, B. 1989. The effect of fish density and species combination on growth and utilization of natural food in ponds. *Aqua. Fish. Manage.*, **20**: 59-71.
- Hossain, M. A. 1996. Proximate and amino acid composition of some potential Bangladeshi fish feed ingredients. *Bangladesh Journal of Zoology*, 24: 163–168.
- Hossain, M. A., Rahmatullah, S. M., Islam, M. S., Kabir, A. K. M. A., Islam, M. S. and Dewan, S. 1997. Impact of chapila (*Gudusia chapra* Ham.) on growth of carps in polyculture. *Bangladesh Journal of Fisheries Research*, **1**(2): 19-23.
- Huchinson, G. E. 1957. *A treatise on limnology*. Vol. I. John Willey Sons. Inc. London, pp. 34-36.

- Huet, M. 1972. *Text book of fish culture, breeding and cultivation of fish.* Fishing New books Ltd. UK, p. 436.
- IIRR-ICLARM, 1992. Integration of aquaculture into crop-animal systems in Asia. Aquaculture System. *Volume 71, Issues 1-2.* January-February 2002, pp. 127-146.
- Iqbal, S. 1995. Practical report PRISM Shobuj Shona Centre Mirzapur, Bangladesh.
- Iqbal, S. 1999. Duckweed Aquaculture: potentials, possibilities and limitations for combined waste water treatment and animal feed production in developing countries. SUNDEC Report No. 6/99, Switzerland, 91p.
- Islam, M. S. 2002. Evaluation of supplementary feeds for semiintensive pond culture of Mahaseer, *Tor putitora* (Hamilton). *Aquaculture*, **212**: 263– 276.
- Islam, M. S., Ali, M. A. and Mazid, M. A. 1997. Polyculture of three exotic species in seasonal ponds under different fertilization and feeding conditions. *Bangladesh J. Train. Dev.*, **10**:175-179.
- Islam, M. S., Azim, M. E., Rahman, M. M. and Mazid, M. A. 1998. Culture prospects of sharpunti (*Puntius gonionotus* Bleeker) and common carp (*Cyprinus carpio* Linnaeus) in backyard ditches at farmers level. *Progress. Agric.*, 8(1-2): 191–194.
- Jasmine, S., Molina, M., Hossain, M. Y., Jewel, M. A. S., Ahamed, F. and Fulanda, B. 2011. Potential and economic viability of freshwater prawn *Macrobrachium rosenbergii* (de Man, 1879) polyculture with Indian major carps in Northwestern Bangladesh. *Our Nature*, **9**: 61-72
- Jena, J. K., Aravindakshan, P. K., Chandra, S., Muduli, H. K. and Ayyappan, S. 1998. Comparative evaluation of growth and survival of Indian major carps and exotic carps in raising fingerlings. *J. Aqua. Trop.*, **13**(2), 143– 150.
- Jena, J. K., Ayyappan, S. and Aravindakshan, P. K. 2002a. Comparative evaluation of production performance in varied cropping patterns of carppolyculture systems. *Aquaculture*, **207**(1-2): 49-64.

- Jena, J. K., Ayyappan, S., Aravindakshan, P. K., Dash, B., Singh, S. K. and Muduli, H. K. 2002b. Evaluation of production performance in carp polyculture with different stocking densities and species combinations. *Journal of Applied Ichthyology*, **18**(3): 165-171.
- Jhingran, V. G. 1975. *Fish and Fisheries of India*. Hindustan Publishing Corporation, Delhi, India, 453p.
- Jhingran, V. G. 1991. *Fish and Fisheries of India*. Hindustan Publishing Co., New Delhi, 727p.
- Journey, W. K., Skillcorn, P. and Spira, W. 1990. Duckweed aquaculture: a new aquatic farming system in developing countries. Technical Department, Agriculture Division, The World Bank.
- Kabir, A. N. M. A., Hossain, M. A. and Rahman, M. S. 2009. Use of duckweed as feed for fishes in polyculture. *J. Agric. Rural Dev.*, **7**(1 & 2): 157-160.
- Khan, S., Hossain, M. S. and Haque, M. M. 2009. Effects of feeding schedule on growth, production and economics of pangasiid catfish (*Pangasius hypophthalmus*) and silver carp (*Hypophthalmichthys molitrix*) polyculture. *J. Bangladesh Agrill. Univ.* **7**(1): 175-181.
- Kohinoor, A. H. M., Islam, M. L., Wahab, M. A. and Thilsted, S. H. 1998. Effect of mola (*Amblypharyngodon mola* Ham.) on the growth and production of carps in polyculture. *Bangladesh J. Fish. Res.*, **2**(2): 119-126.
- Kohinoor, A. H. M., Islam, M. S., Begum, N. and Hussain, M. G. 1999. Production of Thai sharpunti (*Barbodes gonionotus* Bleeker) in polyculture with carps using low-cost feed. *Bangladesh J. Fish. Res.*, 3(2): 157–164.
- Lakshmanan, M. A. V., Murthy, D. S., Pillai, K. K. and Banerjee, M. 1967. On a new artificial feed for carp. fry. *FAO Fisheries Report* (44): 373–387.
- Lakshmanan, M. A. V., Sukumaran, K. K., Murty, D. S., Chakraborty, D. P. and Philipose, M. T. 1971. Preliminary observations on intensive fish farming in freshwater ponds by the composite culture of Indian and exotic species. *J. Inland Fish. Soc. India*, **3**: 1–21.

- LeCren, E. D. 1965. Some factors regulating the size of population of Freshwater. *Mitt. Int. Verein. Theoro Agew. Limnol,* **13:** 88-105.
- Lumpkin, T. A. and Plucknett, L. 1982. *Azolla* as a green manure: use and management in crop production. Westview press Boulder, Colorado. Westview Tropical Agriculture, series#15, 230p.
- Mahaboob, S. and Sheri, A. N. 1997. Growth performance of major, common and some Chinese carps under composite culture systems with special reference to pond fertilization. *J. Aqua. Trop.*, **12**: 201–207.
- Mahmud, A. A. 1998. Optimization of stocking density density of Thai sarpunti (*Puntius gonionotus*) as an additional component in polyculture.*Bangladesh J. Train. Dev.*, **11**: 111-117.
- Mairs, D. F. 1996. A total alkalinity atlas for marine lakes waters. *Limnol. Oceanogr.*, **11**: 68-72.
- Majhi, S. K., Das, A. and Mandal, B. K. 2006. Growth performance and production of organically cultured grass carp *Ctenopharyngodon idella* (Val.) under mid-hill conditions of Meghalaya; North Eastern India. *Turkish Journal of Fisheries and Aquatic Sciences*, **6**: 105-108.
- Mandal, R. N., Datta, A. K., Sarangi, N. and Mukhopadhyay, P. K. 2010. Diversity of aquatic macrophytes as food and feed components to herbivorous fish – a review. *Indian J. Fish.*, **57** (3): 65-73.
- Mazid, M. A., Zaher, M., Begum, N. N., Ali, M. Z. and Nahar, F. 1997. Formulation of cost-effective feeds from locally available ingredients for carp polyculture system for increased production. *Aquaculture*, **151**: 71–78.
- Miah, M. S., Uddin, M. S. and Shah, M. S. 1993. Effects of artificial feed in carps polyculture system. *Bangladesh J. Agric. Sci.*, **20**: 359-364.
- Miah, M. S., Uddin, M. S. and Shah, M. S. 1997. Effect of stocking ratios on the growth and production of fishes in mixed polyculture system. *Bangladesh Journal of Fisheries*, **20**: 135-138.

- Miajae, M. A. H. 1999. Effect of supplementary feed on the growth of Indian major carps with Thai sarpunti in polyculture. *Bangladesh J. Train. Dev.*, **13**: 117-124.
- Miao, S. 1992. Growth and survival model of red tail shrimp (*Penaens penicillatus*) (Alock) according to manipulating stocking density. *Bull. Inst. Zool., Academia Sinica*, **31**: 1-8.
- Mibagwv, I. G. and Adeniji, H. A. 1988. The nutritional content of duckweed (*Lemna paucicostata*) in the Kainji lake area, Nigeria. *Aquatic Botany*, 29: 357-366.
- Michael, R. G. 1968. Seasonal trends in physico- chemical factors and plankton in a freshwater fish pond and their role in fish culture. *Hydrobiol*, **33**: 144-158.
- Mitra, B., Gupta, A. and Laka, U. K. 1978. Effects of some manures on the growth and production of major carps in village ponds of district Birbhum, West Bengal. *Environ. Ecol.*, **5**: 381-385.
- Mondal, T. K. and Ray, A. K. 1999. The nutritive value of *Acacia auriculiformis* leaf meal in compounded diets for *Labeo rohita* fingerlings. *The Fourth Indian Fisheries Forum Proceedings* 1999, *Kochi*. pp. 295–298.
- Moriarty, D. J. W. 1997. The role of microorganisms in aquaculture ponds. *Aquaculture*, **151**(1): 333-349.
- Morrice, C. P. 1998. A trial to test the suitability of a low-input cost, silver barb (*Puntius gonionotus*)-based, polyculture model for seasonal ponds in northwest Bangladesh. *Asian Fisheries Science*, **11**: 121-132.
- Mostaque, A. 1995. Studies on the mixed culture of fishes in seasonal ponds through fertilization and feeding. *Bangladesh J. Train. Dev.*, **8**: 97-106.
- Mukhopadhyay, P. K. and Jena, J. K. 1999. Use of non-conventional dietary ingredients in fish feed formulation, In: Saksena. D. N. (Ed.), *Recent Research Advances in Ichthyology* (chapter-15): Oxford IBH Publishing House Co. Pvt. Ltd., New Delhi and Calcutta, pp. 225-248.

- Murthy, H. S. 2003. Nutrient requirements of Indian Major carps. *Global Aquaculture Advocate*, pp. 58-59.
- Murty, A. S. 1986. *Toxicity of pesticides to fish*. Vols 1 and 11 C.R .C. Press Inc., 483p.
- New, M. B. and Csavas, I. 1993. Aquafeeds in Asia-a regional overview, p.1-23. In: M. B. New, A. G. J. Tacon and I. Csavas (eds.) *Farm-made aquafeeds*. Proceedings of the FAO/AADCP Regional Expert Consultation on Farm-Made Aquafeeds, 14-18 December 1992, Bangkok, Thailand.
- Nikolsky, G. V. 1963. The Ecology of Fishes. Academic Press, London. 352p.
- Okeyo, O. D. 1989. Herbivory in freshwater fishes: a review. *Israeli J. Aquacult.*, *Bamidgeh*, **41**: 79-97.
- Openheimer, J. R., Ahmed, M. G., Haq, A., Alam, A. K. M. A., Aziz, K. M., Ali, S. and Haque, A. S. M. N. 1978. Limnological studies of three ponds in Dhaka, Bangladesh. *Bangladesh J. Fish.*, **1**: 10-28.
- Oron, G. 1994. Duckweed culture for wastewater renovation and biomass production. *Agric. Water Manag.* **26**: 27–40.
- Papoutsoqlou, S. E., Papaparskeve-Papoutsoglou, E. and Aklexis, M. N. 1987. Effect of density on growth rate and production of rainbow trout (*Salmo gairdneri* Richardson) over a full rearing period. *Aquaculture*, **66**: 9-17.
- Patra, B. C. and Ray, A. K. 1988. A preliminary study on the utilization of the aquatic weed *Hydrilla verticillata* Rayle as feed by the carp, *Labeo rohita* (Hamilton): growth and certain biochemical composition of flesh. *Indian Biology*, **XX**(I): 44–50.
- Peters, G. A., Mayne, B. C., Ray, T. B. and Toia, R. E. 1979. Physiology and biochemistry of the *Azolla-Anabaena* symbiosis. pp. 325-344. In: *Nitrogen and Rice*. Los Baños, Laguna, Phillipines, International Rice Research Institute.
- Phaohorm, S. 1980. The study of biology, feeding and growth of *Puntius gonionotus*. Annual report, Department of Fisheries, Bangkok, Thailand.

- PRISM-Bangladesh, 1990. Shobuj Shona Project. Progress Report No. 3. Dhaka, Bangladesh: Prism-Bangladesh, 51p.
- Pullin, R. S. V. and Almazan, G. 1983. *Azolla* as a fish food. *ICLARM Newsletter*, **6**(1): 6-7.
- Rahman, M. A., Hossain, M. K. and Azad, M. A. K. 2006. Culture potential of Thai sharpunti, *Barbodes gonionotus* (Bleeker) with major carps in seasonal ponds. *Pakistan J. Biological Sci.* 9(10): 1891-1897.
- Rahman, M. A., Rahman, M. R. and Rahman, M. S. 2007. Evaluation of growth and production of the mahseer, *Tor putitora* (Ham.) in polyculture with indigenous major carps. pp. 161-175. In: *Mahseer: The Biology and conservation* (Eds. S.S. Siraj, A. Christianus, N.C. Kiat and S.S. De Silva). Malaysian Fisheries Society Occasional Publication No. 14. Kuala Lumpur, Malaysia.
- Rahman, M. M. 1999. Effects of species combination on pond ecology and growth of fish in carp SIS polyculture systems. *Bangladesh J. Aquacult.*, **2-5**(1): 43-49.
- Rahman, M. M., Varga, I. and Chowdhury, S. N. 1992. Manual on polyculture and integrated fish farming in Bangladesh. Food and Agriculture Organization (FAO), Rome, Italy.
- Rahman, M. S. 1992. *Water quality management in aquaculture*. BRAC Prokashana. 66 Mohakhali, Dhaka-1212 Bangladesh, p. 27.
- Rahman, M. S., Chowdhury, M. Y., Haque, A. K. M. A. and Haq, M. S. 1982. Limnological studies of four ponds. *Bangladesh J. Fish*, **2-5**(1-2): 25-35.
- Rangama, S. 1979. Manual of analysis of fruits and vegetable products, Tata McGraw- Hill Publishing Company Ltd. New Delhi. 179pp.
- Ray, A. K. and Das, I. 1995. Evaluation of dried aquatic weed, *Pistia stratiotes* meal as a feedstuff in pelleted feed for rohu, *Labeo rohita* fingerlings. *Journal of Applied Aquaculture*, **5**: 35–44.

- Robinette, H. R., Brunson, M. W. and Day, E. J. 1980. Use of duck weed in diets of channel catfish. *Proc.* 13th Ann. Conf. SE Assoc. Fish Wildlife Age, 108-114 pp.
- Rothuis, A. J., Duong, L. T., Richter, C. J. J., Ollevier, F. 1998. Polyculture of silver barb, *Puntius gonionotus* (Bleeker), Nile tilapia *Oreochromis niloticus* (L.) and common carp, *Cyprinus carpio* L., in Vietnamese ricefields: feeding ecology and impact on rice and ricefield environment. *Aquac. Res.*, **29**: 661–668.
- Roy, N. C., Kohinoor, A. H. M., Wahab, M. A. and Thilsted, S. H. 2002. Evaluation of performance of carp-SIS polyculture technology in the rural farmers' pond. *Asian Fisheries Science*, **15**: 43-52.
- Roy, N. C., Wahab, M. A., Khatoon, H. and Thisted, S. H. 2003. Economics of Carp-SIS polyculture in rural farmer's pond. *Pakistan Journal of Biological Sciences*, 6(1): 61-64.
- Saha, G. N., Sehgal, K. L., Mitra, E. and Nandy, A. C. 1971. Studies on the seasonal and diurnal variations in physico-chemical conditions of a perennial freshwater pond. *J. Inland Fish. Soc. India*, **3**: 79.
- Santhanam, R., Sukumaran, N. and Natarajan, P. 1990. *A Manual of Fresh Water Aquaculture.* Oxford and IBH Publishing Go. Pvt. Ltd., New Delhi, 193 pp.
- Sehagal , H. S. and Toor, H. S. 1991. Comparison of feeding strategies based on biomass and biomass-pond interactions. In: *Fish Nutrition Research in Asia* (ed. By S.S. De Silva), pp.181-192. Asian Fisheries Society, Manila, Philippines.
- Shang, Y. C. 1981. Aquaculture Economics: Basic concept and methods of analysis. West View Press. Boulder, Colorado. 153p.
- Shanmugasundaram, V. S and Balusamy, M. 1993. Rice-fish-azolla: a sustainable farming system. *NAGA, The ICLARM Quarterly*, **16**(2-3): 23.
- Sharmin, R. and Jewel, M. A. S. 2008. Occurrence and abundance of *Microcystis* species in the urban fish ponds. *Int. J. Ani. And Fish. Sci.* 1(2): 53-59.

- Singh, P. K., Gaur, S. R. and Chari, M. S. 2006. Growth Performance of Labeo rohita (Ham.) Fed on Diet Containing Different Levels of Slaughter House Waste. J. Fish. Aquat. Sci., 1(1): 10-16.
- Singh, S. B., Pillai, K. K., and Chakraborty, P. C. 1967. Observation on the efficacy of grass carp in controlling and utilizing aquatic weeds in ponds in India. *Proc. Indo-Pacific Fish Counc.*, **12**(2): 220-235.
- Sinha, M. and Saha, P. K., 1980. Efficacy of a commercial fish feed for composite fish culture. *J. Inland Fish. Soc. India*, **12**: 51– 55.
- Sinha, V. R. P. and Gupta, M. V. 1975. On the growth of grass carp *Ctenopharyngodon idella* Val. in composite fish culture at Kalyani, West Bengal (India). *Aquaculture*, **5**: 283–290.
- Sinha, V. R. P., Gupta, M. V., Banerjee, M. K. and Kumar, D. 1973. Composite fish culture in Kalyani. *J. Inland Fish. Soc. India*, **5**: 201–208.
- Sithara, K. and Kamalaveni, K. 2008. Formulation of low-cost feed using *Azolla* as a protein supplement and its influence on feed utilization in fishes. *Current Biotica*, **2**(2): 212-219.
- Skillicorn, P., Spira, W. and Journey, W. 1993. *Duckweed Aquaculture- A New Aquatic Farming System for Developing Countries*. The World Bank, Washington DC. 76p.
- Subasinghe, R., Soto, D. and Jia, J. 2009. Global aquaculture and its role in sustainable development. *Review in Aquaculture*, **1**: 2-9.
- Suresh, V. R. and Mandal, B. K. 2000. Growth response and nutritive value of Azolla and Alternanthera incorporated pelleted feeds on fingerlings of Cyprinus carpio var. communis: a preliminary study. Indian J. Fish. 47(3): 225-229.
- Swingle, H. S. 1967. Standardization of chemical analysis for water and pond mud. FAO, *Fish Rep.* **44**(4): 397-421.
- Tacon, A. G. J. 1990. *Standard methods for the nutrition and feeding of farmed fish and shrimp.* Washington DC, Argent Laboratories Press, 454 pp.

- Tacon, A. G. J. and Jackson, A. J. 1985. Utilization of conventional and unconventional protein sources in practical fish feeds. In: *Nutrition and Feeding in Fish,* Cowey, C.B., Mackie, A.M. and Bell, J.G., (eds). Academic Press, London, pp. 119-145.
- Talukdar, M. Z. H., Shahjahan, M. and Rahman, M. S. 2012. Suitability of duckweed (*Lemna minor*) as feed for fish in polyculture system. *Int. J. Agril. Innov. & Tech.*, 2(1): 42-46.
- Tavares, F. A., Roudrigues, J. S. R., Fracalossi, D. M., Esquivel, J. and Roubach, R. 2008. Dried duckweed and commercial feed promote adequate growth performance of tilapia fingerlings. *Biotemas*, **21**(3): 91-97.
- Thy, S., Borin, K., Vanvuth, T., Buntha, P. and Preston, T. R. 2004. Effect of water spinach and duckweed on fish growth performance in polyculture ponds. *Livestock Research for Rural Development*, **20**(1): 1-9.
- Tripathi, S. D., Aravindakshan, P. K., Ayyappan, S., Jena, J. K., Muduli, H. K., Suresh, C. and Pani, K. C. 2000. New high in carp production in India through intensive polyculture. *J. Aqua. Trop.*, **15**(2): 119–128.
- Uddin, M. S., Miah, M. S. and Alam, M. S. 1994. Study on production optimization through polyculture of indigenous and exotic carps. *Bangladesh J. Train. Dev.*, **7**: 67-72.
- Vander Steen, P., Brenner, A. and Oron, G. 1998. An integrated duckweed and algae pond system for nitrogen removal and renovation. *Water Sci. Technol.*, **38**: 335–343.
- Van–Dyke, J. M. and Sutton, D. L. 1977. Digestion of duck weed (*Lemna spp*) by the grass carp (*Ctenopharyngodon idella*). *Journal of Fish Biology*, 11: 273-278.
- Van-Hove, C. and Lopez, Y. 1983. In *Azolla* physiology. Technical Bulletin, National University of Colombia, Faculty of Agricultural Sciences, Palmira. Volume I, Number 1. pp. 43-58.

- Van-Hove, C., de Waha Baillonville, T., Diara, H. F., Godard, P., Mai Kodomi, Y. and Sanginga, N. 1987. *Azolla* collection and selection. *Azolla* Utilization. *In*: Proceedings of the Workshop on *Azolla* Use, Fuzhou, Fujian, China, 31 March±5 April 1985. *Int. Rice Res. Inst., Los Banos, Philippines*, pp. 77-87.
- Vass, K. F. and Sachlan, M. 1955. Limnological studies on diurnal fluctuation on shallow ponds in Indonesia. *Verh. Int. Ver. Limnol.*, **12**: 309-319.
- Verma, N. M. 1969. Hydrobiological study of a tropical impoundment, Gualior, India, with special reference to the breeding of Indian carps. *Hydrobiol.*, 34: 358-368.
- Vromant, N., Nam, C. Q. and Ollevier, F. 2002. Growth performance of Barbodes gonionotus (Bleeker) in intensively cultivated rice fields. Aquaculture, 212: 167–178.
- Wahab, M. A., Ahmed, Z. F., Haq, M. S. and Begum, M. 1994. Compatibility of silver carp in the polyculture of cyprinid fishes. *Prog. Agric.*, 5(2): 221-227.
- Wahab, M. A., Ahmed, Z. F., Islam, A. and Ramatullah, S. M. 1995. Effect of introduction of common carp, *Cyprinus carpio* (L.) on the pond ecology and growth of fish in polyculture. *Aquacult. Res.*, **26**: 619-628.
- Wahab, M. A., Azim, M. E., Haque, M. M. and Ahmed, Z. F. 1996. Effects of frequency of fertilization on water quality and fish yields. *Progress. Agric.*, 7(2): 33-39.
- Wahab, M. A., Rahman, M. M. and Milstein, A. 2002. The effect of common carp, *Cyprinus carpio* (L.) and mrigal, *Cirrhinus mrigala* as bottom feeders in major Indian carp polyculture. *Aquac. Res.*, **33**: 547–556.
- Wee, K. L. and Wang, S. S. 1987. Nutritive value of Leucaena leaf meal in pelleted feed for Nile Tilapia. *Aquaculture*, **62**(2): 97–108.
- Welch, P. S. 1952. *Limnology*. McGraw-Hill Book Company Inc, New York, p. 218.

- Wurts, W. A. and Durbrow, R. M. 1992. Interactions of pH, carbon dioxide, alkalinity and hardness in fish ponds. Southern Regional Aquaculture Center. SRAC Publication No. 464. 4p.
- Wyban, J. A., Sweeney, J. N. and Kanna, R. A. 1988. Shrimp yields and economic potential of intensive round pond systems. *Journal of the World Aquaculture Society*, **19**: 210-217.
- Zaher, M. and Mazid, M. A. 1994. Aquafeeds and feeding strategies in Bangladesh. pp. 161–180. In: New, M. B., Tacon, A. G. J. and Csavas, I. (eds.) *Farm-made aquafeeds*. FAO Fisheries Technical Paper No. 343, Rome, FAO.

APPENDICES

Appendix-1: Meteorological data of Kushtia district in the year of 2010

Months Parameters	April	Мау	June	July	August	September
Air temperature (°C)	32.82	32.95	34.02	32.50	32.00	31.52
Average Rainfall (mm)	84.00	171.00	351.00	330.00	418.00	235.00

Source: Bangladesh Meteorological Department, First class Meteorological Observatory, Chuadanga, Bangladesh

Appendix-2: Meteorological data of Kushtia district in the year of 2011

Months Parameters	April	Мау	June	July	August	September
Air temperature (°C)	32.65	33.25	34.15	33.26	32.64	30.25
Average Rainfall (mm)	85.00	150.00	358.00	321.00	403.00	239.00

Source: Bangladesh Meteorological Department, First class Meteorological Observatory, Chuadanga, Bangladesh

Appendix-3:	Meteorological	data of Kushtia	district in the	vear of 2012
Appendix-0.	meteororogical	uata or Rushila		year or 2012

Months Parameters	April	Мау	June	July	August	September
Air temperature (°C)	33.25	33.80	34.25	33.80	32.20	31.65
Average Rainfall (mm)	88.00	166.00	384.00	352.00	415.00	243.00

Source: Bangladesh Meteorological Department, First class Meteorological Observatory, Chuadanga, Bangladesh