

2015

Effect of Methoxy and Lipophilic Prenyl Substituents on Structure Activity Relationship of Some Synthesized Hydroxychalcones; Determined by Antimicrobial, Antioxidant and Cytotoxic Activity

Khatun, Mst. Marzina

University of Rajshahi

<http://rulrepository.ru.ac.bd/handle/123456789/805>

Copyright to the University of Rajshahi. All rights reserved. Downloaded from RUCL Institutional Repository.

A Thesis for the M. Phil. Degree

**Effect of Methoxy and Lipophilic Prenyl Substituents
on Structure Activity Relationship of Some
Synthesized Hydroxychalcones; Determined by
Antimicrobial, Antioxidant and Cytotoxic Activity**



*Submitted to the Department of Chemistry,
University of Rajshahi, in fulfillment of the requirement for the degree of
M. Phil. in chemistry*

SUBMITTED BY

Mst. Marzina Khatun
Exam Roll No.: 12232
Registration No. : 2830
M. Phil. Session: 2012-2013

December, 2015
Second Science Building

Department of Chemistry,
Faculty of Science

Rajshahi University
Rajshahi - 6205



EXAMINER COPY

*DEDICATED TO
MY
BELOVED PARENTS
& MY FAMILY*

DECLARATION

I, Mst. Marzina Khatun, hereby declare that this thesis has been written independently. It has not been published anywhere before. Due references and acknowledgement have been stated properly where materials of other Authors were used.

Marzina Khatun

Mst. Marzina Khatun

CERTIFICATE

This is to certify that the thesis entitled “Effect of Methoxy and Lipophilic Prenyl Substituents on Structure Activity Relationship of Some Synthesized Hydroxychalcones; Determined by Antimicrobial, Antioxidant and Cytotoxic Activity” submitted to the Department of Chemistry, University of Rajshahi for the degree of Master in Philosophy (M.Phil.) embodied the result of a piece of bona fide research work carried out by Mst. MarzinaKhatun bearing Roll No. 12232, Registration No. 2830 under our supervision and guidance. No Part of the thesis has been submitted for any other degree.

Supervisor



Dr. Bilkis Jahan Lumbiny
Assistant Professor
Department of Chemistry
University of Rajshahi.

Co-Supervisor



Dr. Md. Azizul Islam
Professor
Department of Chemistry
University of Rajshahi.

Abstract

Natural chalcones and their derivatives, plant pigments are abundantly present in fern to higher plants especially in different parts of edible plants. They are belonging to flavonoid family and also considered to be main precursors in the biosynthesis of flavones, isoflavones and other biologically essential heterocycles. Flavonoids are the most common naturally occurring antioxidants and found ubiquitously in plants as pigments for flower coloration, in fruits and vegetables and play an ecological role in nature. A series of 2'- hydroxychalcones and their derivatives are found to have potential therapeutic agents against bacterial diseases. Chalcones are 1, 3-diphenyl-2-propene-1-one, in which two aromatic rings are linked by a three carbon α , β -unsaturated carbonyl system (-CO-CH=CH-). These possess conjugated double bonds and a completely delocalized π -electron system on both benzene rings. Molecules possessing such system have relatively low redox potentials and have a greater probability of undergoing electron transfer reactions. During the last sixty years synthetic as well as isolation works on the chalcones are being done throughout the world. They have been demonstrated to possess unique templates associated with very diverse application mainly biological and pharmacological activities such as antibacterial, antiviral, antioxidant, anti-inflammatory, antimutagenic, antiallergic, antiviral, antineoplastic, anti-thrombotic and vasodilatory activities and inhibitory activities in several enzymes. Structure activity relationship (SAR) is also an important factor which has aroused considerable interest to the chemist. Different functional groups which are attached to the benzene ring of chalcones can be varied to enhance activity. Chalcones and its derivatives are still an object of sustained interest thus the present study has been extended by synthesizing eight substituted chalcones; 2' - hydroxychalcone (**3a**), 2' - hydroxy - 4 - methoxychalcone (**3b**), 2' -

hydroxy - 2, 4, 5 - trimethoxychalcone (**3c**), 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (**3d**), 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (**3e**), 2' - hydroxy - 3' - C - prenylchalcone (**4a**), 2' - hydroxy - 5' - C - prenylchalcone (**4b**), 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (**4c**) based on Claisen-Schmidt condensation method from substituted benzaldehyde and acetophenone in good yield. They have been characterized by spectral data. The potency of the development of new antibiotic or chemotherapeutic agents is highly based on *in vitro* antimicrobial screening. The synthesized chalcones (**3a-3e**, **4a-4c**) were screened *in vitro* for their antibacterial activity against four pathogenic bacteria viz. *Bacillus caerius* (G⁺, B₁), *Staphylococcus aureus* (G⁺, B₂), *Eschericia coli* (G⁻, B₃), *Agrobacterium Species* (G⁻, B₄). The primary assay was performed by disc diffusion technique to classify the microorganism susceptible as well as resistant towards particular compounds. The bioactivity is expressed by the diameter of zone of inhibition in mm. Synthesized chalcones 3d, 3b, 3b, 4c in 250 µg/disc showed very high activity against B₁ (38), B₂ (40), B₃ (30), B₄ (40) respectively comparing to those of the standard drug (ciprofloxacin, C-50). The rest of the chalcones showed moderately good activity in different condition. In addition the synthesized chalcones (**3a-3e**, **4a-4c**) were evaluated for *in vitro* antioxidant activity using diphenylpicrylhydrazyl (DPPH) model. Observation for antioxidant activity is expressed in terms of percent scavenging of DPPH radical and the inhibitory concentration 50% (IC₅₀) is lowest for structure **3d**, 0.923 (µg/mL) at DPPH conc. 0.02% indicating highest activity. The compound **3d** contains two phenolic -OH group and suppose to produce phenoxide free radical easily and stabilized by electronic group methoxy occupied ortho and para position of aromatic ring attached to benzaldehyde part. Other structures showed good antioxidant activity as compared to the standard, ascorbic acid,

IC₅₀ 0.08 (μg/mL). The antioxidant activity was also measured in another method using reducing power capacity. All the chalcones especially prenylated chalcone showed appreciable reducing activity and **3c** showed the highest value. Cytotoxic activities of the same compounds were undertaken *in vivo* by brine shrimp lethality test (BST) and expressed by lethal concentration 50%, (LC₅₀). The compound **3d** showed the least toxicity having LC₅₀ 71.75 (μg/mL) whereas LC₅₀ values less than 30 ppm for pure compounds were considered toxic. Increment the no. of methoxy and hydroxy group causes higher LC₅₀ values among the compounds (**3a-3d**). As steric hindrance effect causes better tolerance by reducing their killing ability. On the contrary prenylation causes cell membrane permeability and enhance toxicity. In **3e** three methoxy groups are occupied ring A and in three adjacent carbon. It shows poor activities in any sort of biological screening undertaken.

Contents

Chapter One: Introduction

1. General Discussion on Flavonoid and related compounds	1
1.1 Chalcone	2
1.1.1 Natural chalcones and biological activity	4
1.1.2 Synthetic chalcones and their importance	11
1.1.3 Prenylated chalcones and their biological importance	12
1.2 Synthesis of chalcones through various methods	13
1.2.1 The Claisen-Schmidt reaction	13
1.2.1.1 Mechanism of chalcone formation	14
1.2.2 Synthesis of chalcone via Suzuki coupling reaction	14
1.2.3 Synthesis of chalcone using Borontrifluoride- etherate	15
1.2.4 Synthesis of chalcone via Friedel-Crafts acylation	16
1.2.5 Synthesis of chalcones by using some modified methods	16
1.3 Aim of the present study	19
1.4 Statement of present study	20

Chapter Two: Experimental

2.1 Section A : Materials and Instrumentals	23
2.1.1 Melting Point Apparatus	23
2.1.2 Spectral techniques (UV, IR, NMR)	23
2.1.3 Thin Layer Chromatography (TLC)	24
2.1.3.1 Development of TLC plates	24
2.1.4 Preparative Thin Layer Chromatography (PTLC)	25
2.1.5 Column Chromatography	26
2.1.6 Reagent and Solvent and their Purification	27
2.1.7 Cleaning and drying of glassware	28

2.2	Section B: Synthesis and characterization of 2'-hydroxychalcones (3a-3e, 4a-4c)	29
2.2.1	Preparation of β -resacetophenone	29
2.2.2	Synthesis of 2' - hydroxychalcone (3a) and its structural determination	30
2.2.3	Synthesis of 2' - hydroxy - 4 - methoxychalcone (3b) and its structural determination	31
2.2.4	Synthesis of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone (3c) and its structural determination	32
2.2.5	Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (3d) and its structural determination	33
2.2.6	Synthesis of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (3e) and its structural determination	35
2.2.7	Synthesis of 2' - hydroxy - 3' - C - prenylchalcone (4a) and 2' - hydroxy - 5' - C - prenylchalcone (4b) and its structural determination	36
2.2.8	Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (4c) and its structural determination	38

Chapter Three: Biological Activity Screening

3.1	Section A: General discussion on Antimicrobial Activity	40
3.1.1	Methods of Antimicrobial activity determination	42
3.1.2	Disk Diffusion Technique	43
3.1.3	Chemical used	45
3.1.4	Test organism	46
3.1.5	Collection of organisms	46
3.1.6	Description of procedure	46

3.1.6.1 Sterilization procedure	46
3.1.6.2 Culture media	47
3.1.6.3 Preparation of nutrient agar	47
3.1.6.4 Preparation of medium	47
3.1.6.5 Preparation of fresh culture	48
3.1.6.6 Preparation of test plates	48
3.1.6.7 Preparation of discs containing sample	48
3.1.6.7.1 Sample discs	48
3.1.6.7.2 Standard discs	49
3.1.6.8 Placement of the discs and incubation	49
3.1.6.9 Precaution	50
3.1.6.10 Determination of antibacterial activity of the test agents by measuring the zone of inhibition	50
3.2 Section B:General Discussion on Antioxidant activity	50
3.2.1 Study of the antioxidant property by some common methods	55
3.2.1.1 DPPH Radical Scavenging Activity	56
3.2.1.2 Ferrous ion Chelating Activity	57
3.2.1.3 Inhibition of β -carotene Bleaching	57
3.2.1.4 Trolox Equivalent Antioxidant Capacity (TEAC)	58
3.3 Radical Scavenging Activity Determination of the Present Study by DPPH Method	58
3.3.1 Principle	59
3.3.2 Materials	60
3.3.3 Experimental procedure	60
3.4 Reducing Power Capacity	62
3.4.1 Reagents	63
3.4.2 Procedure	63
3.5 Cytotoxicity Bioassay	65

Chapter Four: Result and Discussion

4.1 Section A: Synthesis and structure Elucidation	67
4.2 Section B: Results for Antimicrobial activity	72
4.3 Antioxidant Activity Screening by DPPH method	78
4.4 Reducing Power Capacity	81
4.5 Cytotoxic activity	85

Chapter 5: Conclusion

CONCLUTION	88-91
REFERENCES	92-95
APPENDIX	96-97

List of Tables

Table 1: The list of naturally occurring chalcones and their sources	4
Table 2: List of synthesized chalcones used for antibacterial activities	45
Table 3: List of the test microorganisms	46
Table 4: List of some foods containing high levels of antioxidants	53
Table 5: List of some naturally occurring antioxidant	53
Table 6: List of some commonly used antioxidant used as standard	56
Table 7: List of spectral data for some important functionality for compounds (3a-3e, 4a-4c)	69
Table 8: Results of the antibacterial activity of the compounds (3a-3e, 4a-4c) against <i>Bacillus caerius</i> . (G^+ , B_1) at 250 ($\mu\text{g disc}^{-1}$)	72
Table 9: Results of the antibacterial activity of the compounds against <i>Staphylococcus aureus</i> (G^+ , B_2) at 250 ($\mu\text{g disc}^{-1}$)	73
Table 10: Results of the antibacterial activity of the compounds against <i>Escherichia coli</i> . (G^- , B_3) at 250 ($\mu\text{g disc}^{-1}$)	74
Table 11: Results of the antibacterial activity of the compounds against <i>Agrobacterium Species</i> (G^- , B_4) at 250 ($\mu\text{g disc}^{-1}$)	75
Table 12: DPPH radical scavenging data of synthesized chalcones (3a-3e, 4a, 4c) and their corresponding IC_{50} values	78
Table 13: Reducing power of chalcones (3a-3e, 4a-4c) with standard Ascorbic acid	82
Table 14: Brine Shrimp Lethality Assay of Chalcones (3c-3e, 4a-4c)	85

List of Figures

Figure 1: General structure of chalcone	3
Figure 2: A plate for calculation of R_f value	25
Figure 3: Structure of 1, 1-diphenyl-2-picrylhydrazyl (DPPH)	59
Figure 4: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against <i>Bacillus caerius</i> (G^+ , B_1)	76
Figure 5: Photographic representation of the zone of inhibition against <i>Staphylococcus aureus</i> (G^+ , B_2)	76
Figure 6: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against <i>Escherichia coli</i> (G^- , B_3).	77
Figure 7: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against <i>Agrobacterium Species</i> (G^- , B_4)	77
Figure 8: DPPH radical scavenging activity of synthesized chalcones (3a-3e, 4a, 4c) without standard Ascorbic acid.	80
Figure 9: Comparison of IC_{50} value of Ascorbic acid and synthesized chalcones (3a-3e, 4a, 4c).	80
Figure 10: Reducing power of chalcones (3a-3e, 4a-4c) with standard Ascorbic acid	83
Figure 11: Reducing power of chalcones (3a-3e, 4a-4c) without standard	84
Figure 12: Reducing power of chalcones (3a-3e, 4a-4c) compare to standard Ascorbic acid in $10 \mu\text{g/mL}$ concentration	84
Figure 13: Bar diagram of LC_{50}	87

Scheme

Scheme 1: The Claisen-Schmidt condensation reaction scheme.	13
Scheme 2: Mechanism for the base-catalyzed Claisen-Schmidt condensation between acetophenone and benzaldehyde	14
Scheme 3: Synthesis of chalcone via Suzuki coupling reaction	15
Scheme 4: Synthesis of chalcone using Borontrifluoride-etherate	16
Scheme 5: Synthesis of chalcone via Friedel-Crafts Acylation	16
Scheme 6: Base catalyzed condensation for synthesis of chalcone	17
Scheme 7: Synthetic route for five synthesized chalcones (3a-3e) from their corresponding acetophenones (1a-1c) and aldehydes (2a-2e)	21
Scheme 8: Synthesis of 2' - hydroxychalcone (3a) by condensation of 2 - hydroxyacetophenone (1a) with benzaldehyde (2a) under alkaline conditions	30
Scheme 9: Synthesis of 2' - hydroxy - 4 - methoxychalcone (3b) by condensation of 2 -hydroxyacetophenone (1a) with 4 - methoxybenzaldehyde (2b) under alkaline condition	31
Scheme 10: Synthesis of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone (3c) by condensation of 2 - hydroxyacetophenone (1a) with 2, 4, 5 - trimethoxybenzaldehyde (2c) under alkaline condition	32
Scheme 11: Synthesis of 2', 5' - dihydroxy - 2, 4, 6 -trimethoxy chalcone (3d) by condensation of 2, 5 - dihydroxyacetophenone (1b) with 2, 4, 6 - trimethoxy benzaldehyde (2d) under alkaline condition	34

Scheme 12: Synthesis of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (3e) by condensation of 3, 4, 5 - trimethoxyacetophenone (1c) with 4 - hydroxybenzaldehyde (2e) under alkaline condition	35
Scheme 13: Synthesis of 2' - hydroxy - 3' - C - prenylchalcone (4a) and 2' - hydroxy - 5' - C - prenylchalcone (4b) by prenylation of 2' -hydroxychalcone (3a)	36
Scheme 14: Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (4c) by prenylation of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (3d)	38
Scheme 15: Free radical scavenging mechanism of vitamin E	54
Scheme 16 Determination of DPPH radical scavenging activity	62
Scheme 17: Determination Reducing power capacity	64
Scheme 18: Determination of cytotoxicity of chalcone	66

Abbreviations

IR	Infra red
IUPAC	International Union of Pure and Applied Chemistry
IC ₅₀	The Inhibitory Concentration 50%
LC ₅₀	Lethal concentration 50%
NMR	Nuclear Magnetic Rsonance
TLC	Thin Layer Chromatography
TMS	Tetramethylsilane
UV	Ultra Violet

List of Symbols and Units

Symbol	Full Expression
asym	Asymmetric Stretching
bs	Broad Singlet
b.p.	Boiling point
β	Bronsted Coefficient
$^{\circ}\text{C}$	Degree Celsius
cm	Centimeter
d	Doublet
dd	Doublet of doublet
δ	Delta, Chemical Shift in NMR Spectrum

g	Gram
hrs.	Hours
Hz	Hertz
J	Coupling Constant
λ_{\max}	UV absorbance maxima
m	Multiplet
mL	Mili (10^{-3}) litre
mins.	Minutes
m.p	Melting point
MHz	Mega Hertz
μ	Micro (10^{-6})
μL	Microlitre
μg	Microgram
nm	Nano Meter
ppm	Parts per million
s	Singlet
Str	Stretching
Sym	Symmetric Stretching
t	Triplet



CHAPTER ONE
INTRODUCTION

CHAPTER ONE

Introduction

1. General Discussion on Flavonoid and related compounds

A natural product is a chemical compound or substance produced by a living organism, found in nature. In the broadest sense, natural products include any substance produced by life. Natural products can also be prepared by chemical synthesis and have played a central role in the development of the field of organic chemistry by providing challenging synthetic targets. The term natural product has also been extended for commercial purposes to refer to cosmetics, dietary supplements, and foods produced from natural sources without added artificial ingredients.

Within the field of organic chemistry, the definition of natural product is usually restricted to mean purified organic compounds isolated from natural sources that are produced by the pathways of primary or secondary metabolism. Within the field of medicinal chemistry, the definition is often further restricted to secondary metabolites. Secondary metabolites are not essential for survival, but nevertheless provide organisms that produce them an evolutionary advantage. Natural products sometimes have pharmacological or biological activity that can be of therapeutic benefit in treating diseases. As such, natural products are the active components not only of most traditional medicines but also many modern medicines. Furthermore, the structural diversity of natural products can be readily achievable by chemical synthesis and thus synthetic analogues can be prepared with improved potency. Hence natural products are often used as starting points for drug discovery.

The world of nature abounds in organic compounds of every conceivable structural class. The cells of living organisms, plants, fungi, bacteria, insects, other animals are the sites of intricate and complex biosyntheses that result in the fragmentation of many varieties of organic compounds of great practical importance to mankind. The structures of these naturally occurring compounds are often extremely complex and elucidation of their structures being a major challenge to organic chemists and biochemists alike. Structural and synthetic study of natural products constitute one of the most fascinating and fruitful field of study open to organic chemists.

Flavonoids, the most common naturally occurring antioxidants are found ubiquitously in plants as pigments for flower coloration, in fruits, vegetables and beverages.^[1] They are prominent plant secondary metabolites that have been found in dietary components including fruits, vegetables, olive oil, tea, and red wine.^[2] Chalcone is one of distinctive subclass of flavonoid being generated high interest in scientific study all over the world as it is associated with many different biological and pharmacological activities.^[2, 3]

1.1 Chalcone

Benzylideneacetophenones (IUPAC name 1, 3- diphenyl-2-propen-1-one) constitute a class of naturally occurring pigments, which are often referred to as 'Chalcones' [Fig.1, also see Table-1]. The term was first coined by Kostanecki.^[4] An interesting feature of Chalcones (polyhydroxylated) is that they serve as the precursors for the synthesis of another class of naturally occurring and widely distributed flavonoid pigments called 'Flavones'.





Figure 1: General structure of chalcone

The chemistry of chalcones has generated intensive scientific studies throughout the world. Especially interest has been focused on the synthesis and biodynamic activities of chalcones. These compounds are also known as benzalacetophenone or benzylidene acetophenone. In chalcones, two aromatic rings are linked by an aliphatic three carbon chain. Chalcone bears a very good synthon so that variety of novel heterocyclic derivatives with good pharmaceutical profile can be designed. Chalcones are α,β -unsaturated ketone containing the reactive ketoethylenic group. These are colored compounds because of the presence of the chromophore $-\text{CO}-\text{CH}=\text{CH}-$, which deepens in the presence of other auxochromes, also used as non azo dye industrially.^[5] Chalcones, bright yellow colored compounds are found most conspicuously in flowers and are known to play an ecological role in nature in relation to colors of the leaves and flowers of plants. Naturally occurring chalcones are all hydroxylated to a greater or lesser extent. Chemically chalcones are polyphenolic heteronuclear compounds include two aromatic ring bounds by an α,β unsaturated carbonyl group, a unique templates associated with very diverse application as antioxidant and antimicrobial activity.^[2] During last sixty years synthetic as well as isolation works from plant material of chalcone being carried out throughout the world. Structural modification leads structure activity relationship (SAR) is also an important factor which has aroused considerable interest to the chemist.








1.1.1 Natural chalcones and biological activity







Natural Chalcones and their derivatives, plant pigments are abundantly present in fern to higher plants especially in different parts of edible plants. They are belonging to flavonoid family and also considered to be main precursors in the biosynthesis of flavones, isoflavones and other biologically essential heterocycles.^[6] The chalcones have been found useful in elucidating structure of natural products like hemlock tannin, cyanomaclurin, ploreitin, eriodictyol and homo eriodictyol, naringenin etc.^[5] Chalcone and its derivatives have also attracted vast attention due to numerous pharmacological properties. The compounds with chalcone's skeleton have been reported to possess a broad spectrum of biological activities such as anti-inflammatory, chemopreventive, anticancer, antiproliferative, antimalarial, antiparasitary and anti-HIV activity. Some of their derivatives exhibit analgesic, antipyretic, antitumor and cytotoxic activity.^[6] Chalcones are isolated and separated from natural resources or dye yielding plants and are known to be used as a natural dye, a good alternative to nitrogenous toxic azo dye. The list of some naturally occurring chalcones given by Mabry *et al.*^[7] can be supplemented with new chalcones which have been isolated recently as tabulated below.







Table 1: The list of naturally occurring chalcones and their sources







Name	Structure	Source	Reference
2, 4- dihydroxy - 6 - methoxychalcone		<i>Dragon's blood Resin</i>	Cardillo, <i>et al.</i> , (1971), <i>J. Chem. Soc., C</i> , 3967-3970







2, 4- dihydroxy-3- methyl - 6 - methoxychalcone		-do-	-do-
2'-hydroxy - 4', 6', 3, 4 tetramethoxy chalcone		<i>Pongamia</i> <i>Pinata</i>	Tanaka <i>et al.</i> , (1992), <i>Phytochem</i> , 31 (3), 998
2', 4'- dimethoxy - 3, 4 -methylene dioxy chalcone		-do-	-do-
2', 4', 6'- trimethoxy - 3, 4 - methylenedioxy- chalcone		-do-	-do-
2'-methoxyfurono [2'', 3'': 4', 3']- chalcone		-do-	-do-
Triangularin		<i>Pityrogramma</i> <i>triangularis</i>	Mabry <i>et al.</i> , (1978) <i>Phytochemistry</i> , 17, 586
2'- hydroxy - 4', 5', 6', 3, 4 - pentamethoxy chalcone		<i>Chromo-</i> <i>laena odorata</i>	Barua <i>et al.</i> , (1978) <i>Phytochemistry</i> , 17, 1807


Quinochalcone-C		-do-	-do-
2'- hydroxy - 3, 4, 5, 4', 6'- pentamethoxy chalcone		<i>Merrillea caloxylon</i>	Fraser <i>et al.</i> , (1972), <i>Phytochemistry</i> , 11, 868
Flavokawain-C		<i>Piper Methylsticum</i>	Dutta <i>et al.</i> , (1973), <i>Indian J. Chem.</i> 11, 509
2', 4', 6', 4, α - pentahydroxychalcone		<i>Berchemia zeyheria</i>	Volsteedt <i>et al.</i> , (1973), <i>Tet. Lett.</i> , 1001
Pashanone		<i>Didymocarpus pedicellantus</i>	Agarwal <i>et al.</i> , (1973), <i>Indian J. Chem.</i> 11, 9.
Licochalcone-B		<i>Glycyrrhiza glabra</i>	Saitoh <i>et al.</i> , (1975), <i>Tet. Lett.</i> , 50, 4461
3', 6'- dihydroxyl- 2', 4', 5', 4 - tetramethoxy - chalcone		<i>Flemingia strobilifera</i>	Bhatt (1975), <i>Indian J. Chem.</i> 13, 1105.

Kamakugiol		<i>Lindera erythrocarpa</i>	Liu <i>et al.</i> , (1975), <i>J. Pharm. Soc. (Jpn)</i> , 19, 1114
Lassein		<i>Lassea nitida</i>	Pederiva <i>et al.</i> , (1975), <i>An. Assoc. Quim. Argent</i> ; 63, 85.
Aurentiacin		<i>Didymopanax pedicellantus</i>	Chaudhury, <i>et al.</i> , (1976), <i>Phytochemistry</i> , 15, 224.
Cerasidin		<i>Prunus cerasus</i>	Parmer and Nagaranjan, (1977), <i>Phytochemistry</i> , 16, 1317
Cerasin		-do-	-do-
2'-hydroxy - 4', 6'-dimethoxy-3'-methylchalcone		<i>Myrica gale</i>	Malterud, K. E., <i>et al.</i> , <i>Phytochemistry</i> , 16, 1805

2', 4' - dihydroxy - 6'-methoxy-3', 5'- dimethyl chalcone		-do-	-do-
2', 3', 4', 3, 4 - penta- hydroxychalcone		<i>Albilizia adianthfolia</i>	Candy <i>et al.</i> , (1978) <i>Phytochem.</i> , 17, 1807.
Helilandin-B		<i>Helichrysum sutherlandii</i>	Bohlmann <i>et al.</i> , (1978), <i>Phytochem.</i> , 17, 1935.
Isodidymocarpin		<i>Didymo carpus pedicellata</i>	Bose and Chaudhury, (1978), <i>J. Indian Chem. Soc.</i> , 55, 1198
2'-hydroxy - 3', 4', 6'-trimethoxy- chalcone		<i>Popowia cauliflora</i>	Panichpol and Watermann, (1978), <i>Phytochem.</i> , 17, 1363.
2', 3', 4', 6'- tetra- methoxychalcone		-do-	-do-

2', 4- dihydroxyl- 4', 5', 6'- trimethoxychalcone		<i>Chromo- laena</i>	Barua <i>et al.</i> , (1978), <i>Phytochemistry</i> , 17, 1807.
2'-hydroxy- 4, 4', 5', 6'- tetramethoxychalco ne		-do-	-do-
Helilandin-A		<i>Helichrysum sutherlandii</i>	Bohlmann <i>et al.</i> , (1978), <i>Phytochem.</i> , 17, 1935.
Isonebarachalcone		<i>Psoralea corylifolia</i>	Gupta, <i>et al.</i> , (1980), <i>Phytochem.</i> , 19 (9)
2', 4-dihydroxy-3'- methoxy-5'-C- prenyl chalcone.		-do-	Bhalla, V. K, <i>et al.</i> , (1968) <i>Tett. Lett.</i> , 2401.
2', 4', 4-trihydroxy- 3'-C- prenylchalcone.		-do-	-do-

2', 4'-dihydroxy-6'-methoxy-3'-C-prenylchalcone		Sophora angustifolia (Leguminosae)	Komatsu, M. <i>et al.</i> , (1970) Yakugaku. Zasshi, 90 , 463
2', 4', 6'-trihydroxy-3'-C-prenylchalcone		Glycyrrhizagl abra (Leguminosae)	Kattav, N. S. <i>et al.</i> , (1972) Khim. Prir. Soedin, 8 , 805.
2'-hydroxy-4'-methoxy-3'-prenyl chalcone.		Derris Sericea (Leguminosae)	Nascimento do, <i>et al.</i> , (1972) <i>Phytochem</i> , 11 , 3023.
2', 6'-dihydroxy-4', 3, 4, 5-tetramethoxy-5'-C-prenylchalcone		Artocarpus Heterophyllus (Moraceae)	Dayal, R., <i>et al.</i> , (1974). <i>Indian J. Chem.</i> , 12 (8) 895.
2', 4', 6'-trihydroxy-5'-C-prenylchalcone		Deriris rariflora. (Leguminosae)	Filho, R. B., <i>et al.</i> , (1975). <i>Phytochem</i> , 14 , 261.
2', 6'-dihydroxy-4'-methoxy-5'-C-prenylchalcone		Deriris rariflora. (Leguminosae)	Filho, R. B., <i>et al.</i> , (1975). <i>Phytochem</i> , 14 , 261.

2', 4', 6', 3-tetrahydroxy-5'-C-prenylchalcone		Sophora Tomentosa	Komatsu, M. <i>et al.</i> , (1978). <i>Chem. Pharm. Bull.</i> , 26 (12) 3863.
--	---	----------------------	--

1.1.2 Synthetic chalcones and their importance

To improve the biological activities and to better understand the relation between chalcones intake and health outcomes, researches have been done continuously. Many analogues of naturally occurring chalcones introducing new substituent groups and their derivatives have been synthesized and reported. Chalcones possess conjugated double bonds and a completely delocalized π -electron system on both benzene rings. Molecules possessing such system have relatively low redox potentials and have a greater probability of undergoing electron transfer reactions.^[8] Furthermore chalcones are important intermediate in many addition reactions of nucleophiles owing to inductive polarization of carbonyl group at the β position.^[6] Besides these compounds have broad industrial application as artificial sweeteners, scintillator polymerization catalyst, fluorescent whitening agent, organic brightening agent, stabilizer against heat, visible light, ultraviolet light. So the chemistry of chalcones has generated intensive scientific studies throughout the world specially interest has been focused on the synthesis and biodynamic activities of chalcones. There are several methods for the syntheses of chalcones have been investigated. The major synthetic schemes are including Claisen-Schmidt condensation, Suzuki coupling reaction, Wittig reaction, Friedel-Craft acylation, photo-Fries rearrangement, carbonylative Heck reaction also unconventional method via

microwave irradiation. [6] On the Suzuki coupling reaction, benzoyl chloride was reacted with phenyl vinyl boronic acid using anhydrous toluene and catalyzed by tetrakis (triphenylphosphine) and cesium carbonate as base, gave 3', 4', 4'-trimethoxychalcone. In photo-Fries rearrangement, phenyl cinnamate undergoes rearrangement and provided two chalcones molecules, ortho and para-hydroxychalcones. In carbonylative Heck reaction, chalcone was produced by reaction of arylhalide and styrene that added by carbon monoxide and catalyzed by palladium and a phosphine-amine ligand, N-heterocyclic carbene. [9]

1.1.3 Prenylated chalcones and their biological importance


3-methyl-but-2-en-1-yl substituent is named as prenyl functional group. The prenylchalcones are a group of natural products found in fruits, vegetables, nuts, seeds and flowers as well as in teas and wines are important constituent of human diet. Prenyl group is lipophilic by nature and assume to facilitate attachment to cell membrane. Thirty seven naturally occurring prenyl compounds were investigated for their inhibitory activities on the mouse brain monoamino oxidase (MAO) *in vitro*. Three prenylchalcones Lupinifolinol (IC₅₀, 1 μmol), Prenylnaringenin (IC₅₀, 4 μmol) and Topazolin (IC₅₀, 5 μmol) were judged to be very active. [5] Thus in the present study we are interested to extend our studies by the introduction of hydrophobic functionality like prenyl (3-methyl-but-2-en-1yl) group at benzene nucleus moiety of chalcone which may give an important information about positional effect of prenyl group and its role as auxochromes to show enhance and stabilized color effect also to study its bioactivity.

1.2 Synthesis of chalcones through various methods

1.2.1 The Claisen-Schmidt reaction^[10]

The Claisen-Schmidt condensation is a classical method to synthesize chalcones which employs cross aldol condensation of appropriate aldehyde and ketone by base catalyzed (such as NaOH, KOH, Ba(OH)₂, NaCO₃, hydrotalcities, etc.) or acid catalyzed (such as HCl, BF₃, B₂O₃, *p*-toluenesulfonic acid, SOCl₂, etc.) and then followed by dehydration.^[8]


The Claisen-Schmidt condensation (scheme 1) between acetophenone and benzaldehyde derivatives is an important C-C bond forming reaction which allows α , β -unsaturated ketone such as chalcone to be obtained. It is the most frequently used means of establishing the C₆-C₃-C₆ flavonoids nucleus owing to the availability of starting materials such as 2-hydroxyacetophenone and benzaldehyde derivatives to obtain a 2'-hydroxychalcone. Chalcone (Figure 1) bears A-ring substituent provided by the acetophenone (indicated as R₁) and B-ring substituent provided by benzaldehyde (indicated as R₂). The classical Claisen-Schmidt reaction is routinely carried out using aqueous sodium or potassium hydroxide, otherwise ethanolic potassium hydroxide at 50 °C over a period of several hrs. The benzaldehyde derivative is often used in slightly more than equivalent amounts. The extensive conjugation of the products causing absorbance of light in the visible region, leading this yellow color.



Scheme 1: The Claisen-Schmidt condensation reaction scheme.

1.2.1.1 Mechanism of chalcone formation ^[6, 11, 12]

Sodium hydroxide in an alcoholic solvent which used in this experiment leads to fast and reversible formation of intermediate compound. For that reason, it was used to increase the reaction rate of Claisen-Schmidt condensation. Base removed acidic alpha hydrogen from acetophenone, producing a resonance-stabilized enolate ion. This enolate ion then attacked aldehyde molecule, yielding a neutral condensation product and followed by dehydration to generate chalcones in good yields.




Scheme 2: Mechanism for the base-catalyzed Claisen-Schmidt condensation between acetophenone and benzaldehyde

1.2.2 Synthesis of chalcone via Suzuki coupling reaction ^[8]

An efficient synthesis of chalcones was carried out based on the Suzuki coupling reaction between benzoyl chlorides and phenyl vinyl boronic acid.


Phenyl vinyl boronic acid was prepared by dehydrogenative borylation of para-methoxystryne by pinacoleborane oxidative addition-dehydrogenation, catalyzed by the rodium complex, $\text{RhCl}(\text{cod})_2$ to give paramethoxy phenyl ethenylboronic acid pinacol ester (Scheme 3)



Scheme 3: Synthesis of chalcone via Suzuki coupling reaction

1.2.3 Synthesis of chalcone Using Borontrifluoride-etherate


Narender and Reddy ^[13] developed a new methodology by using $\text{BF}_3 \cdot \text{Et}_2\text{O}$ to synthesize several substituted chalcones. The advantages of this method over the existing methods are high yield, simple work-up, short reaction times, no side reactions and separation is needed to get the products. This method is applicable for reactions involving liquid reactants which have base sensitive functional groups such as esters and amides. A condensation between 2-acylated acetophenone and the respective aromatic aldehyde produce 2-acylated chalcone in high yields by using $\text{BF}_3 \cdot \text{Et}_2\text{O}$ as shown in Scheme 4



Scheme 4: Synthesis of chalcone using Borontrifluoride-etherate

1.2.4 Synthesis of chalcone via Friedel-Crafts acylation ^[8]

Besides the Claisen-Schmidt reaction, chalcone can also be synthesized by direct Friedel-Crafts acylation of a phenol. In this approach the phenol becomes the A-ring while the acylation agent provides both the B-ring and the three carbon bridge to form C₆-C₃-C₆ unit. Friedel-Crafts acylation of 2,4-dimethyl-1,3,5-triolbenzene with 3-phenylpropionyl chloride gave 2',4',6'-trihydroxy-3',5'-dimethylchalcone as shown in Scheme 5.




Scheme 5: Synthesis of chalcone via Friedel-Crafts acylation

1.2.5 Synthesis of chalcones by using some modified methods

The substituted benzylidene acetophenones have likewise been obtained by condensing the appropriately substituted acetophenones with substituted benzaldehydes in the presence of alkali. The 2-hydroxyacetophenones on acid or base catalyzed condensation with aldehydes yield either a chalcone or a flavone or a mixture of both. In Kostanecki's method ^[4], the starting materials are suitably substituted 2-hydroxyacetophenone (furnishing A-ring) and benzaldehyde (furnishing B-ring) which are condensed in the

presence of alcoholic caustic potash to give chalcones. If only 2'-hydroxyl group of the chalcone is free, the condensation proceeds smoothly in 10% alcoholic caustic potash in as high yield as 85% but in the case of compounds with a large number of free hydroxyl groups, higher concentration of alcoholic caustic potash (50%-60%) is required. The yields are highest when alcoholic caustic potash (50%-60%) is used at 0-20 °C for 15-72 hours and when starting materials are methoxylated or benzylated. But for polyhydroxy acetophenones good yields are obtained only when the condensation is performed at 0 °C. Chaudhury *et al.*^[14] synthesized flemichapparin by the base condensation of 2, 4-dihydroxy-5-methoxyacetophenone with benzaldehyde at 0-5 °C for 8 days.



Scheme 6: Base catalyzed condensation for synthesis of chalcone.

A modification of the Kostanecki's method was worked out by Russel and Todd.^[15] It involves the condensation of benzoylated derivatives of 2-hydroxyacetophenone with benzoylated derivatives of 2-hydroxybenzaldehyde in the presence of dry hydrogenchloride at 0 °C followed by hydrolysis with alkali to the free chalcone. Thus 2, 4-dibenzoyloxyacetophenone condenses with dibenzoyloxy protocatechuic aldehyde to give the tetrabenzoyloxy chalcone which on hydrolysis yields 2', 4', 3, 4-tetrahydroxy chalcone. The synthesis of some chalcones has been accomplished by using sodium hydride as the base for the condensation. This method gives good yields even if the hydroxyl groups are not protected. Thus 2', 3, 4-trihydroxy-3', 4', 5'-trimethoxychalcone has been synthesized by

Stout and Stouf^[6] by using sodium hydroxide as the base. Chalcone has also been synthesized by irradiating a solution of phenyl- cinnamate in methanol, with UV light (253 nm) for 1.5 hrs. under nitrogen atmosphere. Synthesized two isomeric chalcones were reported by this method.^[17] Synthesis of chalcones using organo phosphorus activated reagent^[18] has been reported. Lupi and co-workers^[19] while synthesizing various prenylated and chromeno chalcones under alkaline conditions have noticed the yield variation by changing the temperature, solvent and using different bases for condensation. Sodium hydroxide, sodium ethoxide and piperidine were used as bases in solvents like absolute alcohol, dry DMSO, anhydrous dioxane at a temperature varying between 25-100 °C for a period ranging from 1-30 hrs. The best yield was obtained when piperidine was used as a base in absolute alcohol for a period of 6 hrs. at a temperature between 60-70 °C using a little excess of aldehyde than the stoichiometric amount. Obera *et al.*^[20] have reported the synthesis of 2', 3', 4', 6', 4-pentahydroxy chalcone is very good yield by the base catalyzed condensation of 2, 3, 4, 6-tetramethoxy acetophenone and methoxybenzaldehyde followed by demethoxy methylation using methanolic hydrochloric acid of the many methods available for protection of hydroxyl groups alkylation with methoxy methyl chloride appeared to be attractive since excellent yields were obtained by heating in acid medium.

Nevertheless, many of these methods require prolonged reaction times, gave poor yields, low selectivity and also suffered from harsh reaction condition for instance toxic reagent, strong acidic and strong basic condition. From many literatures, Claisen-Schmidt condensation still occupies leading positions for synthesizing chalcone, since that method is very simple, inexpensive and easy to conduct.


1.3 Aim of the present study ^[5, 21, 22]

Natural chalcones and their derivatives, plant pigments are abundantly present in fern to higher plants especially in different parts of edible plants. They are belonging to flavonoid family and also considered to be main precursors in the biosynthesis of flavones, isoflavones and other biologically essential heterocycles. Flavonoids are the most common naturally occurring antioxidants and found ubiquitously in plants as pigments for flower coloration, in fruits and vegetables and play an ecological role in nature. A series of 2'-hydroxychalcones and their derivatives are found to have potential therapeutic agents against bacterial diseases. Chalcones are 1,3-diphenyl-2-propene-1-one, in which two aromatic rings are linked by a three carbon α , β -unsaturated carbonyl system (-CO-CH=CH-). These possess conjugated double bonds and a completely delocalized π -electron system on both benzene rings. Molecules possessing such system have relatively low redox potentials and have a greater probability of undergoing electron transfer reactions. During the last sixty years synthetic as well as isolation works on the chalcones are being done throughout the world. They have been demonstrated to possess unique templates associated with very diverse application mainly biological and pharmacological activities such as antibacterial, antiviral, antioxidant, anti-inflammatory, antimutagenic, antiallergic, antiviral, antineoplastic, anti-thrombotic and vasodilatory activities and inhibitory activities in several enzymes. Structure activity relationship (SAR) is also an important factor which has aroused considerable interest to the chemist. Different functional groups which are attached to the benzene ring of chalcones can be varied to enhance activity. Chalcones and its derivatives are still an object of sustained interest. Besides chalcones are isolated and separated from natural resources or dye yielding plants and is known to be used as a natural dye, a good alternative to

nitrogenous toxic azo dye. Synthesis of new naturally occurring organic compounds with basic skeleton of chalcones, flavones and oxygenated flavones and their biological activity were reported by this research groups for long.^[5, 21, 22] In the present study we are interested to extend our studies by the introduction of hydrophilic functionality like hydroxy, methoxy and lipophilic functionality like prenyl (3-methyl-but-2-en-1yl) group at benzene nucleus moiety of chalcone which may give an important information about positional effect of hydroxy, methoxy and prenyl group and its role as auxochromes to show enhance and stabilized color effect also to study its bioactivity.

1.4 Statement of the present study

In the present study eight substituted chalcones were synthesized: 2'-hydroxychalcone (**3a**); 2'-hydroxy-4-methoxychalcone (**3b**); 2'-hydroxy-2,4,5-trimethoxychalcone (**3c**); 2',5'-dihydroxy-2,4,6-trimethoxychalcone (**3d**); 4-hydroxy-3',4',5'-trimethoxychalcone (**3e**); 2'-hydroxy-3'-C-prenylchalcone (**4a**); 2'-hydroxy-5'-C-prenylchalcone (**4b**); 2',5'-dihydroxy-2,4,6-trimethoxy-3'-C-prenylchalcone (**4c**) using Claisen-Schmidt condensation method, a classical, very simple, inexpensive and easy to conduct synthetic strategy. The chalcones (**3a-3e** and **4a-4c**) were obtained from their corresponding acetophenones (**1a-1c**) as well as aldehydes (**2a-2e**) in presence of NaOH/EtOH as shown in scheme 7. The structures of the above compounds were assigned on the basis of spectral data; UV, IR, ¹H and ¹³C NMR together with their elemental analysis.



- | | | |
|--|---|--|
| 1a) $R_1 = -OH,$
$R_2 = R_3 = R_4 = R_5 = H$ | 2a) $R_6 = R_7 = R_8 = R_9 = R_{10} = H$ | 3a) $R_1 = -OH, R_2 = R_3 = R_4 = R_5 = R_6 =$
$R_7 = R_8 = R_9 = R_{10} = H$ |
| 1b) $R_1 = R_4 = -OH,$
$R_2 = R_3 = R_5 = H$ | $R_8 = -OCH_3$ | 3b) $R_1 = -OH, R_2 = R_3 = R_4 = R_5 =$
$R_6 = R_7 = R_9 = R_{10} = H, R_8 = -OCH_3$ |
| 1c) $R_1 = R_5 = H, R_2 = R_3 =$
$R_4 = -OCH_3$ | $R_7 = R_{10} = H$ | 3c) $R_1 = -OH, R_2 = R_3 = R_4 = R_5 = R_7 =$
$R_{10} = H, R_6 = R_8 = R_9 = -OCH_3$ |
| | 2d) $R_6 = R_8 = R_{10} = -OCH_3,$
$R_7 = R_9 = H$ | 3d) $R_1 = R_4 = -OH, R_2 = R_3 = R_5 = R_7 = R_9 =$
$H, R_6 = R_8 = R_{10} = -OCH_3$ |
| | 2e) $R_6 = R_7 = R_9 = R_{10} = H$
$R_8 = -OH$ | 3e) $R_1 = R_5 = R_6 = R_7 = R_9 = R_{10} = H,$
$R_2 = R_3 = R_4 = -OCH_3, R_8 = -OH$ |

Scheme 7: Synthetic route for five synthesized chalcones (**3a-3e**) from their corresponding acetophenones (**1a-1c**) and aldehydes (**2a-2e**)

2-methyl-but-3-en-2-ol in dry dioxane was treated with chalcone (**3a, 3d**) followed by gradual addition of boron trifluoride-etherate to get prenylated chalcone (**4a, 4b, 4c**).

The potency of the development of new antibiotic or chemotherapeutic agents is highly based on *in vitro* antimicrobial screening. The synthesized chalcones (**3a-3e, 4a-4c**) were screened *in vitro* for their antibacterial activity against four bacteria viz. *Bacillus caerius* (G^+ , B₁), *Staphylococcus aureus* (G^+ , B₂), *Eschericia coli* (G^- , B₃), *Agrobacterium Species* (G^- , B₄). The primary assay was performed by disc diffusion technique to classify the microorganism susceptible as well as resistant towards particular compounds. The bioactivity is expressed by the diameter of zone of inhibition in mm. Synthesized chalcones **3d, 3b, 3b, 4c** in 250 μ g/disc showed very high activity against B₁ (38), B₂ (40), B₃ (30), B₄ (40) respectively comparing to those of the standard drug (ciprofloxacin, C-50). The rest of the chalcones showed moderately good activity in different

condition. In addition the synthesized chalcones (**3a-3e**, **4a-4c**) were evaluated for *in vitro* antioxidant activity using 1, 1-diphenyl-2-picrylhydrazyl (DPPH) model. Observation for antioxidant activity is expressed in terms of percent scavenging of DPPH radical and the inhibitory concentration 50% (IC_{50}) is lowest for structure **3d**, 0.923 ($\mu\text{g/mL}$) at DPPH conc. 0.02% indicating highest activity. The compound **3d** contains two phenolic -OH group and suppose to produce phenoxide free radical easily and stabilized by electronic group methoxy occupied ortho and para position of aromatic ring attached to benzaldehyde part. Other structures showed good antioxidant activity as compared to the standard, ascorbic acid, IC_{50} 0.08 ($\mu\text{g/mL}$). The antioxidant activity was also measured in another method using reducing power capacity. All the chalcones especially prenylated chalcone showed appreciable reducing activity and **3c** showed the highest value. Cytotoxic activities of the same compounds were undertaken *in vivo* by brine shrimp lethality test (BST) and expressed by lethal concentration 50%, (LC_{50}). The compound **3d** showed the least toxicity having LC_{50} 71.75 ($\mu\text{g/mL}$) whereas LC_{50} values less than 30 ppm for pure compounds were considered toxic. Increment the no. of methoxy and hydroxy group causes higher LC_{50} values among the compounds (**3a-3d**). As steric hindrance effect causes better tolerance by reducing their killing ability. On the contrary prenylation causes cell membrane permeability and enhance toxicity. In **3e** three methoxy groups are occupied ring A and in three adjacent carbon. It shows poor activities in any sort of biological screening undertaken.



CHAPTER TWO
EXPERIMENTAL

CHAPTER TWO

Experimental

2.1 Section A; General Methods ^[5, 21, 22]

2.1.1 Melting Point Apparatus

Melting points are uncorrected and all solid samples were recorded on an electro-thermal melting point apparatus (EDU-LAB SRL MLTP 06018020).

2.1.2 Spectral techniques (UV, IR, NMR)

(i) **UV:** Chemical reactions were monitored by thin-layer chromatography and spots were visualized with UV light (in nm) using UV-180, SHIMADZU, the spectrophotometer SHIMADZU UV-160, was used to measure absorbance maxima and antioxidant activity both by DPPH method and reducing capacity.

(ii) **IR:** Infrared spectra were recorded on a SHIMADZU “FTIR-8400 spectrophotometer using KBr pellets for solid samples in the Central Science Laboratory, University of Rajshahi, Rajshahi, Bangladesh and the characteristics peaks are expressed in cm^{-1} .

(iii) **NMR:** ^1H NMR and ^{13}C -NMR spectrum was recorded on a BRUKER 400 MHz NMR Spectrometer in CDCl_3 and d_6 -DMSO solution at the Laboratories of Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka and BRUKER 500 MHz SPOECTROMETER University Malaysia Pahang. Chemical shifts were quoted on the δ scale relative to tetra methyl silane (TMS, $\delta = 0$) as an internal standard. The signals are

abbreviated s, d, t, q, m, dd, refer to singlet, doublet, triplet, quartet, multiplet, doublet of doublet respectively. The coupling constant is indicated by J and is given in Hz.

2.1.3 Thin Layer Chromatography (TLC)

The technique of thin layer chromatography (TLC) was extensively used to monitor the progress of the reaction. TLC plates (0.25 mm thickness) were prepared by spreading a layer of silica gel (60 GF₂₅₄, E. MERCK) on clean and thoroughly dried glass plates (2g gel was mixed with 4 mL of distilled water for each plate of size (5cm×20 cm). These were activated by drying at 110 °C before use.

2.1.3.1 Development of TLC plates

i) Using iodine tank: The chromatograms were developed in iodine vapor by placing the plate in the iodine tank.

ii) Using UV light: The presence of the pure compound as a single spot or a mixture can be identified by several spots in UV light.

iii) Using spraying agent: The spots were appeared if the plate was spread with the following spraying reagents.

(i) 2% ferric chloride solution in water is used as a spraying reagent.

(ii) H₂SO₄ (50%) is also used as a spraying reagent and heating the plate.

R_f value is measured from the TLC plate. R_f means retardation factor which measures the velocity of movement of the solute zone relative to that of the solvent front.

$$R_f = \frac{\text{Distance (cm) travelled by solute}}{\text{Distance (cm) travelled by solvent}}$$




Figure 2: A plate for a calculation of R_f value

Usually, the R_f value is constant for any given compound and it corresponds to a physical property of that compound.

2.1.4 Preparative Thin Layer Chromatography (PTLC)

The preparative glass plates (20 cm × 20 cm) were cleaned and dried. The plates were coated with silica gel slurry (silica gel 60 mesh, E. MERCK, 16 g silica gel was mixed with 32 mL of distilled water) with the help of spreader to yield a coating of 0.5 mm thickness. The plates were left at room temperature until their surface become completely dry, the plates were then heated at 120 °C in an electrical oven before use to increase activation. The solutions of compounds to be purified were applied with special type of thin glass capillaries at about 3 cm from bottom of the plates. After drying, the plates were then placed vertically with the spotted end placed downward in chromatographic tanks so that the spotted end of the compounds remained above the solvent. The plates were removed from the tanks when the solvent front reached almost to the upper edge of the plates (1cm far from the upper edge). The plates were usually developed in appropriate solvent and were subsequently dried in the air. The plates were then viewed under UV light or the sides of the plates were exposed to iodine vapor to locate the position of

the compounds. The zones bearing the compounds were scratched off from the plates and subsequently extracted with suitable solvent.

2.1.5 Column Chromatography

Column Chromatography is a very useful technique for the separation of pure compounds from its mixture. For column chromatography silica gel (Kiesel gel-66, 70-230 mesh, ASTM MERCK) used as adsorbent and solvent like n- hexane, petroleum ether, benzene, acetone, ethyl acetate etc. at different proportion was used as eluent. The column was prepared by slurry method silica gel being the stationary phase. The column was thoroughly cleaned and rinsed with acetone and dried. It was then clamped properly and rinsed with solvent used in the preparation of silica gel slurry and cotton plug was fitted at the bottom. The column was half filled with appropriate solvent and the slurry was then poured into it, so that the packing was compact and uniform. Air bubble was avoided by packing the column as quickly as possible. The column was allowed to settle for an hr. The mixture of the compound was completely dried and then was dissolved in eluting solvent and was carefully placed on the surface of the column, and eluted with desired solvent system. In case of polar compounds, the mixture was dissolved in suitable polar solvents and absorbed in small amount of silica gel in a flask. The solvent was removed completely under vacuum by rotary evaporator. From silica gel and this silica gel with absorbed material was put on the surface of the silica gel column. The compounds were then collected as distinct bands. The purity of each band was further checked by TLC examination.

2.1.6 Reagent and Solvent and their Purification

Starting materials, substituted acetophenone (1a, 1c), substituted benzaldehyde (2a-2e), resorsinol, BF_3 -etherate were purchased from Sigma Aldrich and used without further purification.

The reagents and solvents were purified and dried before use where necessary and the rest were used as such from the bottle. All solutions in water, immiscible solvents which has been contact with water were dried over anhydrous sodium sulphate and magnesium sulphate, prior to evaporation. Solvent was usually removed with the help of a vacuum rotary evaporator. Dry hydrogen chloride gas (HCl) was produced by pouring conc. H_2SO_4 over a mixture of sodium chloride and ammonium chloride (5:1) moistened with conc. HCl gas thus produced were dried by passing them through the troughs containing good quality of conc. H_2SO_4 . Pet. ether and benzene were dried by sodium wire and or phosphorus pentoxide (P_2O_5). Acetone and dioxane were dried by keeping over anhydrous K_2CO_3 and distilling the solvents. The liquid reagents were sometimes distilled by following standard method before use.

Acetone: To 500 mL of acetone anhydrous K_2CO_3 (150 g) was added and it was kept overnight. Then acetone was decanted off from K_2CO_3 and distilled to get pure acetone and preserved in an air tight container.

Acetic Acid: About 100 mL of acetic acid was taken in a round bottomed flask and 50 g of CaCl_2 (anhydrous) was added to it. After 6 hrs. the acetic acid was decanted and it was distilled and collected at reduced pressure by maintaining its b.p.

Diethyl ether (Et_2O): Metallic sodium wire was added in diethyl ether and the mixture was kept overnight and was distilled over P_2O_5 at 35°C .

Chloroform (CHCl₃): At first chloroform was shaken with concentrated sulfuric acid after this it was again shaken with distilled water to remove acid. The solvent was then kept in calcium chloride for two hours then it was distilled (b.p. 61 °C).

Methanol: To 500 mL of methanol CaO (50 g) was added and it was kept overnight. Then methanol was decanted off from CaO and refluxed it for 5-6 hrs. using CaCl₂ guard tube. After reflux it was distilled and collected to a round bottom flask. A white cake of Mg-turning was formed in another round bottom flask and the whole mass of methanol was added to it. Again it was refluxed for 4 hrs., distilled out and collected it into an air tight container.

Benzene: To 500 mL of benzene anhydrous CaCl₂ (20 g) was added and it was kept overnight. Then benzene was decanted off from CaCl₂, distilled and collected at 80 °C into an air tight container.

Petroleum ether: To 500 mL of petroleum ether anhydrous CaCl₂ (20 g) was added and it was kept overnight. Then petroleum ether was decanted off from CaCl₂, distilled and collected at 40-60 °C into an air tight container.

Dioxane: To 500 mL of dioxane anhydrous K₂CO₃ (100 g) was added and it was kept overnight. Then dioxane was decanted off from K₂CO₃ and distilled to get pure dioxane and preserved in an air tight container.

2.1.7 Cleaning and drying of glassware

All glassware was cleaned and for most purposes, dried before employed in preparative work in the laboratory. The glassware was washed with a commercial house hold washing powder, which does not scratch glass e.g. "Vim, Trix etc." The washing was either introduced directly into the apparatus or moistened with a little water or it was applied to the dirty surface with a wet test-tube brush, which has been dipped into the powder.

The operation was repeated if necessary. Finally, the apparatus was thoroughly rinsed with distilled water. The most widely used cleaning agent is the chromic acid, cleaning mixture. It is a mixture of chromic acid and concentrated sulphuric acid. It was prepared by the way 5 g of potassium dichromate were dissolved in 5 mL of water in a 250 mL beaker of 100 mL of concentrated sulphuric acid were then added slowly with constant stirring. The temperature will rise of 70-80 °C. The mixture was allowed to cool to about 40 °C and then transferred to a dry glass stopper bottle. Before using chromic acid mixture for cleaning the vessel was rinsed with water to remove organic matter. Sometimes chloroform, acetone, rectified spirit was used as a cleansing agent. Small and bulky both types of glass apparatus were dried by leaving it in an electrically heated oven maintained at 100-120 °C for one to two hrs. Sometimes the glassware was also dried with a commercial hair drier.


2.2 Section B; Synthesis and Characterization of eight 2'-hydroxychalcones (3a-3e, 4a-4c).

2.2.1 Preparation of β -resacetophenone (1b)

Anhydrous zinc chloride (16.5 g) was dissolved in glacial acetic acid (15.8 mL) by warming. To the hot solution at 140 °C, resorcinol (11.0 g) was added with constant stirring. The solution was heated in an oil-bath and the reaction mixture was allowed to complete itself at a temperature not more than 160 °C. After standing for 20 minutes the solution was poured over crushed ice, the precipitate, thus separated was filtered, washed with dil. HCl to remove zinc salts and then with water. It was crystallized from hot dil. HCl (10%, 150 mL), yield 10 g, m. p. 142-143 °C.

2.2.2 Synthesis of 2' - hydroxychalcone (**3a**) and its structural determination

A mixture of 2 - hydroxyacetophenone (**1a**, 0.01 mole) and benzaldehyde (**2a**, 0.01 mole) in ethanolic solution of NaOH (20%, 10 mL) with 30 mL ethanol was stirred at room temperature for about 12 hours. The reaction mixture was kept overnight at room temperature. Then it was diluted with ice cold water and acidified with ice cold dil. HCl. lemon yellow solid (**3a**); yield 61%; melting point 84-85 °C; R_f 0.97 (n-hexane : acetone, 10 : 1) was obtained.




Scheme 8: Synthesis of 2' - hydroxychalcone (**3a**) by condensation of 2 - hydroxyacetophenone (**1a**) with benzaldehyde (**2a**) under alkaline condition

$C_{15}H_{12}O_2$; Solid and lemon yellow, yield 61%; m. p. 84-85 °C, UV: λ_{max} (CH₃OH): 196 and 233 nm, IR (KBr, cm⁻¹): 3414.06 (-OH), 3090 (C=C-H, aromatic, str.), 3046.91 (C=C-H, olifinic, str.), 1959.40 (C=C, olifinic str.), 1640.01 (C=O, conjugated keto group), 1618.00 (C=C, aromatic), 1572.86, 1486.21, 1447.81, 1438.22, 1383.84, 1370.31, 1341.10 (CH₃, bending), 1267.72, 1236.07, 1205.71, 1181.52, 1153.06 (-C-O-C- str.), 1072.50, 1030.80 (C-O, str.), 994.70, 976.52, 864.79, 837.54 (C-H, bending aromatic) [See Appendix, Fig.1], ¹H NMR, δ_H (400 MHz, CDCl₃, 12H); 12.803 (s, 1H, C₂-OH), 7.923 (d, $J = 8.4$ Hz, 1H, C_{6'}-H), 7.921 (d, $J = 14.8$ Hz, 1H, C _{β} -H), 7.668 -7.661 (m, 2H, C₂ and C₆-H), 7.657 (d, $J = 14.8$ Hz, 1H, C _{α} -H), 7.498 (t, 1H, C_{4'}-H), 7.441-7.436 (bs, 3H, C₃, C₄ and C₅-H) 7.029 (d, 1H, $J = 8.4$ Hz,

C_3-H), 6.944 (t, 1H, $C_{5'-H}$), [See Appendix, Fig.2], ^{13}C -NMR (100 MHz, $CDCl_3$, 15C); δ_c 193.77 (1C, $>C=O$), 163.62 (1C, C-2'), 145.49 (1C, C_β), 136.42 (1C, C-4'), 134.63 (1C, C-1'), 130.94 (1C, C-1), 129.67 (1C, C-5'), 129.06 (2C, C-2, C-6), 128.68 (3C, C-3, C-4, C-5), 120.17 (1C, C_α), 120.4 (1C, C-6'), 118.87 (1C, C-3') [See Appendix, Fig.3]

2.2.3 Synthesis of 2' - hydroxy - 4 - methoxychalcone (**3b**) and its structural determination

A mixture of 2 - hydroxyacetophenone (**1a**, 0.01 mole) and 4 - methoxybenzaldehyde (**2b**, 0.01 mole) was treated similar as procedure in 2.2.2 which yielded pineapple yellow solid (**3b**, 61%); m. p. 87-88 °C; R_f 0.88 (n-hexane : acetone, 10:1) was obtained :




Scheme 9: Synthesis of 2' - hydroxy - 4 - methoxychalcone (**3b**) by condensation of 2 - hydroxyacetophenone (**1a**) with 4 - methoxybenzaldehyde (**2b**) under alkaline condition

$C_{16}H_{14}O_3$; Solid and pineapple yellow, yield 61%; m. p. 87-88 °C UV: λ_{max} (CH_3OH): 197 nm, IR (KBr, cm^{-1}): 3414.07 (-OH), 3080 ($C=C-H$, aromatic, str.), 3024.51 ($C=C-H$ olifinic str.), 2969.85 (aliphatic C-H, asym.-str.), 2839.64 (aliphatic C-H, sym-str.), 1922.67 ($C=C$ olifinic str.), 1640.50 ($C=O$, conjugated keto group), 1609.00 ($C=C$, aromatic, str.), 1584.52, 1513.59, 1493.97, 1442.00, 1423.64, 1376.12, 1344.03, 1303.72, (CH_3 , bending), 13275.45, 1261.57, 1211.77, 1177.35, 1164.13, 1130.55 (-C-O-C-

str.), 1032.41(C-O, str.), 985.91, 866.88, 828.34, 804.08 (C-H, bending aromatic) [See Appendix, Fig.4], ^1H NMR, δ_{H} (500 MHz, CDCl_3 , 14H); 12.803 (s, 1H, $\text{C}_{2'}\text{-OH}$), 7.925 (d, $J = 8\text{Hz}$, 1H, $\text{C}_{6'}\text{-H}$), 7.910 (d, $J = 16\text{ Hz}$, 1H, $\text{C}_{\beta}\text{-H}$), 7.640 (dd, $J = 8\text{ Hz}$, 2H, C_2 and $\text{C}_6\text{-H}$), 7.545 (d, $J = 16\text{ Hz}$, 1H, $\text{C}_{\alpha}\text{-H}$), 7.490 (t, 1H, $\text{C}_{4'}\text{-H}$), 7.030 (d, 1H, $J = 8\text{ Hz}$, $\text{C}_{3'}\text{-H}$), 6.960-6.940 (m, 1H, $\text{C}_{5'}\text{-H}$), 6.950 (d, 2H, C_3 and $\text{C}_{5'}\text{-H}$), 3.870 (s, 3H, $\text{C}_4\text{-OCH}_3$) [See Appendix, Fig.5], ^{13}C -NMR (125 MHz, CDCl_3 , 16C); δ_{C} 192.80 (1C, $>\text{C}=\text{O}$), 163.60 (1C, $\text{C}-2'$), 159.80 (1C, $\text{C}-4$), 145.10 (1C, C_{β}), 135.90 (1C, $\text{C}-4'$), 130.20 (2C, $\text{C}-2$ and $\text{C}-6$), 129.60 (1C, $\text{C}-6'$), 127.50 (1C, $\text{C}-1$), 121.80 (1C, $\text{C}-5'$), 121.60 (1C, $\text{C}-1'$), 118.70 (1C, C_{α}), 118.20 (1C, $\text{C}-3'$), 114.20 (2C, $\text{C}-3$ and $\text{C}-5$), 55.80 (1C, $\text{C}_4\text{-OCH}_3$) [See Appendix, Fig.6]

2.2.4 Synthesis of 2'-hydroxy-2,4,5-trimethoxychalcone (3c) and its structural determination

A mixture of 2-hydroxyacetophenone (**1a**, 0.01mole) and 2,4,5-trimethoxybenzaldehyde (**2c**, 0.01mole) was treated similar as procedure in 2.2.2 which yielded a dark yellow solid (**3c**, 61%); R_f 0.84 (benzene: acetone, 5:1) was obtained.




Scheme 10: Synthesis of 2'-hydroxy-2,4,5-trimethoxychalcone (**3c**) by condensation of 2-hydroxyacetophenone (**1a**) with 2,4,5-trimethoxybenzaldehyde (**2c**) under alkaline condition

$C_{18}H_{18}O_5$, Solid and dark yellow, yield 61%; m. p. 120-122 °C, UV: λ_{max} (CH₃OH): 196, 259 and 307 nm, IR (KBr, cm⁻¹): 3416.30 (-OH), 3040 (C=C-H, aromatic, str.), 3009.49 (C=C-H olifinic str.), 2964.99 (aliphatic C-H, asym.-str.), 2864.28 (aliphatic C-H, sym-str.), 2034.08 (C=C, olifinic str.), 1660.01 (C=O, conjugated keto group), 1609.36 (C=C, aromatic), 1520.00, 1479.67, 1412.91, 1358.89 (CH₃, bending), 1292.76, 1218.55, 1187.92 (-C-O-C-, str.), 1128.89, 1026.80, (C-O str.), 995.68, 864.04, 823.76, 757.15 (C-H, bending aromatic) [See Appendix, Fig.7], ¹H NMR, δ_H (400 MHz, CDCl₃, 14H); 13.080 (s, 1H, C_{2'}-OH), 8.170 (d, $J = 16$ Hz, 1H, C _{β} -H), 7.640 (d, $J = 8.6$ Hz, 1H, C_{6'-H}), 7.390 (d, $J = 16$ Hz, 1H, C _{α} -H), 7.370 (m, 1H, C_{4'-H}), 7.010 (m, 1H, C_{5'-H}), 6.920 (d, 1H, $J = 2.6$ Hz, C_{3'-H}), 6.590 (s, 1H, C_{6-H}), 6.100 (s, 1H, C_{3-H}), 3.730 (s, 9H, C₂, C₄, C₅-OCH₃) [See Appendix, Fig.8], ¹³C-NMR (100 MHz, CDCl₃, 16C); δ_c 187.01 (1C, >C=O), 158.50 (1C, C-2'), 153.00 (1C, C-4), 147.80 (1C, C-3), 142.80 (1C, C _{β}), 139.81 (1C, C-4'), 135.71 (1C, C-2), 131.11 (1C, C-6'), 123.90 (1C, C-5'), 123.30 (1C, C-1'), 121.61 (1C, C _{α}), 116.20 (1C, C-3'), 113.81 (1C, C-6), 108.20 (1C, C-1), 100.61 (1C, C-3), 56.40 (C₂-OCH₃), 56.30 (2C, C₄ and C₅-OCH₃) [See Appendix, Fig.9]

2. 2. 5 Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (3d) and its structural determination

A mixture of 2, 5 - dihydroxyacetophenone (**1b**, 0.01 mole) and 2, 4, 6 - trimethoxybenzaldehyde (**2d**, 0.01 mole) was treated similar as procedure in 2.2.2 which yielded orange solid (**3d**, 61%). m. p. 175-176 °C; R_f 0.37 (n-hexane : acetone, 3:1) was obtained.




Scheme 11: Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (**3d**) by condensation of 2, 5 - dihydroxyacetophenone (**1b**) with 2, 4, 6 - trimethoxy benzaldehyde (**2d**) under alkaline condition

$C_{18}H_{18}O_6$; Orange solid, yield 61%; m. p. 175-176 °C, UV: λ_{max} (CH_3OH): 196, 242 and 384 nm, IR (KBr, cm^{-1}): 3415.26 (-OH), 3090.00 (C=C-H, aromatic, str.), 3000.50 (-C=C-H, alifinic, str.), 2940.52 (aliphatic C-H, asym.-str.), 2840.49 (aliphatic C-H, sym-str.), 1605.56 (C=O, conjugated keto group), 1548.81 (C=C, aromatic), 1488.73, 1456.61, 1384.30, 1309.40 (CH_3 , bending), 1269.75, 1213.67, 1159.75, 1120.82 (-C-O-C-,str.), 1062.41, 1022.37 (C-O str.), 992.74, 921.57, 858.89, 828.30 (C-H, bending aromatic) [See Appendix, Fig.10], 1H NMR, δ_H (400 MHz, $CDCl_3$, 18H); 12.769 (s, 2H, $C_{2'}$ and $C_{5'}$ -OH), 8.363 (d, $J=15.6$ Hz, 1H, C_{β} -H), 7.906 (d, $J=16$ Hz, 1H, C_{α} -H), 7.340 (dd, $J=4.0$ & 2.4 Hz, 1H, $C_{6'}$ -H), 7.012-6.982 (m, 1H, $C_{4'}$ -H), 6.894 (t, 1H, $C_{3'}$ -H), 6.877 (bs, 2H, C_3 and C_5 -H), 3.916 (s, 6H, C_2 and C_6 -OCH₃), 3.861 (s, 3H, C_4 -OCH₃) [See Appendix, Fig.11] ^{13}C -NMR (100 MHz, $CDCl_3$, 16C); δ_c 194.89 (1C, >C=O), 163.75 (1C, C-4), 162.09 (2C, C-2 and C-6), 159.00 (1C, $C_{2'}$ -OH), 157.69 (1C, $C_{5'}$ -OH), 147.18 (1C, C_{β}), 137.05 (1C, C-1'), 123.93 (1C, C_{α}), 120.44 (1C, C-6'), 119.48 (1C, C-4'), 119.05 (1C, C-3'), 114.81 (1C, C-1'), 106.55 (1C, C-1), 90.61 (2C, C-3 and C-5), 55.85 (2C, C_2 and C_6 -OCH₃), 55.40 (1C, C_4 -OCH₃) [See Appendix, Fig.12]

2.2.6 Synthesis of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (3e) and its structural determination

A mixture of 3, 4, 5 - trimethoxyacetophenone (**1c**, 0.01mole) and 4 - hydroxy benzaldehyde (**2e**, 0.01 mole) was treated similar as procedure in 2.2.2 which yielded brownish yellow solid (**3e**, 62%). m. p. 177-178 °C; R_f 0.55 (petroleum ether: acetone, 2:1) was obtained and characterized.




Scheme 12: Synthesis of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (**3e**) by condensation of 3, 4, 5 - trimethoxyacetophenone (**1c**) with 4 - hydroxybenzaldehyde (**2e**) under alkaline condition

$C_{18}H_{18}O_5$; brownish yellow, yield 62%; m. p. 177-178 °C, UV: λ_{max} (CH₃OH): 197 nm, IR (KBr, cm^{-1}): 3413.88 (-OH), 3232.28 (C=C-H, aromatic, str.), 3100.00 (-C=C-H, olefinic, str.), 2944.52 (aliphatic C-H, asym.-str.), 2834.07 (aliphatic C-H, sym-str.), 1964 (C-C, C=C), 1640.58 (C=O, conjugated keto group), 1610.44 (C=C, aromatic), 1594.25, 1555.80 (-CH₂ bending), 1464.29, 1439.66, 1413.71, 1384.32, 1349.04 (CH₃, bending), 1277.00, 1233.62, 1203.18 (-C-O-C, str), 1171.84, 1126.91, 1007.65 (C-O, str.), 986.16, 831.80, 720.69, 670.07 (C-H, bending aromatic). [See Appendix, Fig.13], ¹H NMR, δ_H (400 MHz, CDCl₃, 18H) ; 9.850 (s, 1H, C₄-OH), 7.770 (d, $J = 16$ Hz, 1H, C $_{\beta}$ -H), 7.550 (d, $J = 8.4$ Hz, 2H, C₂ and C₆-H), 7.340 (d, $J = 15.6$ Hz, 1H, C $_{\alpha}$ - H), 7.220 (d, 2H, C_{2'} and C_{6'}-H), 6.890 (d, $J = 8.4$ Hz, 2H, C₃ and C₅-H), 3.900 (s, 9H, C_{3'}, C_{4'} and C_{5'}-OCH₃) [See Appendix, Fig.14] ¹³C-NMR (100 MHz, CDCl₃, 18C); δ_c 189.67 (1C,

>C=O), 158.35 (1C, C₄-OH), 153.15 (2C, C-3', and C-5'), 144.94 (1C, C_β), 142.41 (1C, C-4') 133.78 (1C, C-1'), 130.52 (2C, C-2 and C-6), 127.57 (1C, C-1), 119.34 (1C, C_α), 116.07 (2C, C-3 and C-5), 105.95 (2C, C-2' and C-6'), 60.97 (1C, C_{4'}-OCH₃), 56.38 (2C, C_{3'} and C_{5'}-OCH₃) [See Appendix, Fig.15]

2.2.7 Synthesis of 2' - hydroxy - 3' - C - prenylchalcone (4a) and 2' - hydroxy - 5' - C - prenylchalcone (4b):

To a stirred solution of chalcone (2.0 g) in dry dioxane (20 mL) was added gradually borontrifluoride-etherate (2.6 mL) at room temperature during the course of 30 minutes this was added a solution of 2-methyl-but-3-en-2-ol (2.50 mL) in dry dioxane (2.50 mL) and the solution stirred for 6 hrs., kept at room temperature overnight and diluted with moist ether (150 mL). The ethereal layer was washed with water and dried over anhydrous NaSO₂. It was evaporated to dryness and the residue on column chromatography over silica gel and elution successively with n-hexane or a mixture of n-hexane-acetone (3:1). The n-hexane fraction was a light yellow colour gummy substance, 2' - hydroxy - 3' - C - prenylchalcone (4a).



Scheme 13: Synthesis of 2' - hydroxy - 3' - C - prenylchalcone (4a) and 2' - hydroxy - 5' - C - prenylchalcone (4b) by prenylation of 2' -hydroxychalcone (3a)

$C_{20}H_{20}O_2$, gummy substance light, yellow, yield 62%; R_f 0.90, UV: λ_{max} (CH₃OH): 197 nm, 1H NMR, δ_H (500 MHz, CDCl₃, 20 H), 7.935 (dd, $J = 8.0$ Hz, 1H, C_{6'}-H), 7.830 (d, $J = 16.0$ Hz, 1H, C _{β} -H), 7.680 (d, $J = 16.0$ Hz, 1H, C _{α} -H), 7.670-7.650 (m, 1H, C_{5'}-H), 7.460-7.420 (m, 2H, C₂ and C₅-H), 7.200-7.160 (m, 3H, C₃, C₄ and C₅-H), 7.030 (d, $J = 8.0$ Hz, 1H, C_{3'}-H), 5.500 (s, 1H, OH), 4.050-3.900 [m, 1H, -CH₂-CH=C(CH₃)₂], 3.40-3.20 [m, 2H, (-CH₂-CH=C(CH₃)₂), 1.80 (s, CH₃, 6H) [See Appendix, Fig.16], ^{13}C -NMR (125 MHz, CDCl₃, 20 C); δ_c 192.80 (1C, >C=O), 161.50 (1C, C_{2'}-OH), 145.10 (1C, C _{β}), 138.20 (1C, C-4'), 135.20 (1C, C-1), 131.80 [1C, -CH₂-CH=C(CH₃)₂], 129.90 (1C, C-3'), 129.06 (2C, C-2 and C-6), 128.68 (3C, C-3, C-4 and C-5), 123.50 (1C, C-6'), 123.10 [1C, -CH₂-CH=C(CH₃)₂], 122.50 (1C, C-1'), 121.70 (1C, C-5'), 118.70 (1C, C _{α}), 24.60 (1C, -CH₂-CH=C(CH₃)₂), 22.60 [-CH₂-CH=C(CH₃)₂], 18.60 (1C, -CH₂-CH=C(CH₃)₂) [See Appendix, Fig.17]

The n-hexane: acetone (3:1) fraction was a yellow colored gummy substance, 2'-hydroxy-5'-C-prenylchalcone (**4b**).

$C_{20}H_{20}O_2$, Gummy and yellow, yield 62%, R_f 0.90, UV: λ_{max} (CH₃OH): 196 and 241 nm 1H NMR, δ_H (500 MHz, CDCl₃, 20 H), 7.930 (d, $J = 15$ Hz, 1H, C _{β} -H), 7.720-7.700 (m, 1H, C_{6'}-H), 7.640 (d, $J = 10$ Hz, 2H, C₂ and C₆), 7.550 (d, $J = 15$ Hz, 1H, C _{α} -H), 7.450 (bs, 3H, C₃, C₄ and C₅-H), 7.190-7.130 (m, 1H, C_{4'}-H), 6.980-6.960 (m, 1H, C_{3'}-H), 5.500 (s, 1H, OH), 5.200 [s, 1H, -CH₂-CH=C(CH₃)₂], 3.700-3.600 [m, 2H, -CH₂-CH=C(CH₃)₂],


1.820 (3H, -CH₂-CH=C(CH₃)₂), 1.740 (3H, -CH₂-CH=C(CH₃)₂) [See Appendix, Fig.18],

^{13}C -NMR (125 MHz, CDCl₃, 20 C); δ_c 192.80 (1C, >C=O), 160.60 (1C, C_{2'}-OH), 145.10 (1C, C _{β}), 136.20 (1C, C-4'), 135.20 (1C, C-1), 131.80 [1C, -CH₂-CH=C(CH₃)₂], 131.50 (1C, C-6'), 130.50 (1C, C-5'), 128.60 (2C, C-3 and C-5), 128.50 (2C, C-2 and C-6), 127.90 (1C, C-4), 123.10 [1C, -CH₂-CH=C(CH₃)₂], 120.90 (1C, C-1'), 117.40 (1C, C-3'), 118.70 (1C, C _{α}), 34.00

$[-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2]$, 24.60 (1C, $-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2$), 18.60 (1C, $-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2$)
 [See Appendix, Fig.19]

2.2.8 Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (4c):


To a stirred solution of chalcone (2.0 g) in dry dioxane (20 mL) was added gradually borontrifluoride-etherate (2.6 mL) at room temperature during the course of 30 minutes this was added a solution of 2-methyl-but-3-en-2-ol (2.50 mL) in dry dioxane (2.50 mL) and the solution stirred for 6 hrs., kept at room temperature overnight and diluted with moist ether (150 mL). The ethereal layer was washed with water and dried over anhydrous NaSO_2 . It was evaporated to dryness and the residue on column chromatography over silica gel and elution successively with n-hexane-acetone (3:1).



Scheme 14: Synthesis of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (4c) by prenylation of 2', 5' - dihydroxy - 2, 4, 6 - trimethoxychalcone (3d)

$\text{C}_{23}\text{H}_{26}\text{O}_6$, wine red semi solid, yield 62%; R_f 0.91, UV: λ_{max} (CH_3OH): 196 nm, ^1H NMR, δ_{H} (400 MHz, CDCl_3 , 26 H), 8.330 (d, $J = 16.0$ Hz, 1H, $\text{C}_\beta\text{-H}$), 7.420 (d, $J = 16.0$ Hz, 1H, $\text{C}_\alpha\text{-H}$), 7.080 ($J = 8.0$ Hz, 1H, $\text{C}_{4'}\text{-H}$), 7.040 (dd, $J = 8.0$ Hz, 1H, $\text{C}_{6'}\text{-H}$), 7.670-7.650 (m, 1H, $\text{C}_{5'}\text{-H}$), 6.090 (2H, C_3 and $\text{C}_5\text{-H}$), 5.750 [1H, $-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2$], 5.350 (2H, $\text{C}_{2'}$ and $\text{C}_{5'}\text{-OH}$), 3.830 (9H, C_2 and C_4 and $\text{C}_6\text{-OCH}_3$), 3.210 [$-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2$], 1.820 (3H, $-\text{CH}_2-\text{CH}=\text{C}(\text{CH}_3)_2$),

1.700 (3H, $-\text{CH}_2-\text{CH}=\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}$) [See Appendix, Fig.20] ^{13}C -NMR (100 MHz, CDCl_3 , 20 C); δ_c 192.80 (1C, $>\text{C}=\text{O}$), 160.60 (1C, C-4), 159.60 (2C, C-2 and C-6), 154.10 (C_{2'}-OH), 149.60 (1C, C-5'), 134.70 (1C, C-3'), 131.80 (1C, $-\text{CH}_2-\text{CH}=\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}$), 123.90 (1C, C-1'), 123.10 [1C, $-\text{CH}_2-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}$], 122.50 (1C, C_β), 121.30 (1C, C_α), 115.40 (1C, C-4'), 107.00 (1C, C-1), 104.20 (1C, C-6'), 90.90 (2C, C-3 and C-5), 56.20 (2C, C₂ and C₆-OCH₃), 55.8 (1C, C-4), 28.10 ($-\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}-\text{CH}_2-$), 24.60 (1C, $-\text{CH}_2-\text{CH}=\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}$),
18.60 (1C, $-\text{CH}_2-\text{CH}=\overset{\text{CH}_3}{\underset{\text{CH}_3}{\text{C}}}$) [See Appendix, Fig.21]



CHAPTER THREE
BIOLOGICAL ACTIVITY
SCREENING

CHAPTER THREE

Biological Activity Screening

3.1 Section A; General discussion on Antimicrobial Activity ^[5, 11, 21, 22]

Over the past 25 years, the incidence of systemic bacterial infections has been rising dramatically due to an increase in the number of immunocompromised hosts. The search is oriented to find new antibacterial drugs, which may selectively attack the bacteria without inhibiting any biochemical system of the host. ^[23] After the discovery of penicillin and other antibiotics have significantly reduced the complications and mortality rate of infectious diseases. With the wide use of antibiotics in the treatment of bacterial infection has lead to the emergence and spared of resistance strains. The continuous spread of multi drug resistance pathogens have become a serious threat to public health and a major concern for infection control practitioners worldwide.

Bangladesh is predominantly an agricultural country depending mainly on crop plants, agricultural, medicinal and forest product for its economic development. Although crops play a vital role in the economy of the country and agro ecological conditions are favorable for the production of various crops, the yields of crops is often poor. Among the various factors responsible for poor yield of crops, plant disease caused by various microorganisms play a significant role. Gradually men gathered sufficient knowledge of chemistry to inhibit or to kill the microorganism i.e., only inhibit the microbial growth are called 'statis'. But the chemicals which have ability to kill microorganisms are called 'cadal'. Various pesticides are classified as fungicides, viricides, bactericides etc.

Many fatal diseases caused by microorganism viz. bacterial, fungal and viral attacks are known. The treatment of diseases due to bacterial, viral and fungal invasion by chemical compounds were studied and used successfully without affecting the tissues of the host and any other side effects. Many compounds e.g. formaldehyde, iodine, phenol etc. are also active in destroying bacteria.

Antibiotics are the chemical substance obtained from certain non-pathogenic microorganisms (bacteria and fungi) and are used for either killing or inhibiting the growth of pathogenic micro organism without affecting the host tissue.

The words bactericide has originated from Latin words: bacteria. Thus literally speaking a bactericide would be any agencies which have the ability to kill bacteria. By common usage the word is restricted to chemicals. Hence the words bactericide means a chemical capable of killing bacteria. It is not enough that a chemical has high bactericidal activity. Such as chemicals may have no utility unless it stands out in the tests and gives proof of significant control of diseases under varied field conditions. There are several factors which influence the performance of a bactericide under different field conditions. They may be either physical or chemical in nature. The number of chemicals available that shows antibacterial activity runs into hundreds although all are not equally safe, effective and popular. Also different types of organic, aromatic, inorganic and heterocyclic compounds are employed as antibacterial agents.

A biochemical screening to identify the proportions in the clinical use is long overdue. With better understanding of the molecular basis of resistance, it can be envisaged the new diagnostic tools will be developed to allow rapid and appropriate changes in treatment. The use of combinations of bactericides is increasing in medicine and should be a route of choice for

overcoming resistance. [23] Chalcones and their heterocyclic analogs are potential therapeutic agents in bacterial diseases. Chalcones with antibacterial properties have been known since the 1940s. For antibacterial activity, the presence of the enone aggregate in the molecule is important.

New chalcone derivatives are under consideration, and in clinical trial, and will be central to therapy in the medium term. The emergence of resistance to these agents can be predicted and it needs advance assessment to provide assistance for their proper integration into drug therapy.

Compounds with electron releasing groups such as hydroxyl and methoxy showed better antibacterial activity than the others not having such groups. [24] From the above discussion it is clear that the chalcone compounds show antimicrobial activity and widely used as drugs. So, we deliberately synthesized a number of hydroxymethoxy chalcone and finally antibacterial activities of these chalcone were determined. These chalcone were used for valuation of antibacterial activities and syntheses of these compounds are already discussed in the experimental section of the dissertation. The result obtained by *in vitro* antibacterial activity against both Gram-positive and Gram-negative microorganisms of these compounds is discussed in this section.

3.1.1 Methods of Antimicrobial activity determination

In order to detect the antimicrobial activity of a new compound for the development as potential new antibiotic or chemotherapeutic agent *in vitro* antimicrobial screening is a useful technique. In general antimicrobial screening is undertaken in two phases; a primary qualitative assay [24, 25] to detect the presence or absence of activity and secondary assay [26, 27]

which quantifies the relative potency expressed as minimum inhibitory concentration (MIC) value of a pure active compound.

In vitro antibacterial activities of the test chemicals were studied against four pathogenic bacteria. The primary assay can be performed *in vitro* by a number of methods one of which is disc diffusion technique [25, 26] by this method we could classify the organism as susceptible as well as resistance towards particular compounds. The secondary assay is the serial broth dilution assay [11, 24] which quantifies the antimicrobial activity of pure compound by providing the MIC value of the compound for specific susceptible organism. This is an important consideration for the screening of a new antibiotic substance.

3.1.2 Disk Diffusion Technique

The disc diffusion technique [11, 24] is widely accepted for preliminary investigations of materials which are suspected to possess antimicrobial properties. Diffusion procedure is normally used in susceptible intermediate of resistant categories.

Diffusion assays are based on the ability of antibiotics to diffuse from a confined source through a PDA gel and create a concentration gradient. If the agar is seeded or streaked with sensitive organism a zone of inhibition will result where the antibiotic concentration exceeds the minimum inhibitory concentration for that particular organism.

In the disc diffusion technique dried filter paper discs containing known amount of test material are placed on agar plates seeded with test organisms. These plates are kept at low temperature (4 °C) for 24 hrs. Initially the dried discs absorb water from the surrounding test medium and the drug is dissolved. The drug migrates through the adjacent test medium

by concentration gradient of the drug according to physical law that governs diffusion of molecules through an agar gel.^[27] As a result there is a gradual change of drug concentration in the agar surrounding each discs. Then the plates are incubated in an incubator at 37.5 °C for 24 hrs.

As the antibiotic diffusion progresses microbial multiplication also proceeds. At that point of time fungal multiplication proceeds more rapidly than the drug can diffuse and fungal cell which are not inhibited by the antimicrobial agents will continue to multiply until a lawn of growth can be visualized. No growth will appear in the area where drug is present in inhibitory concentrations.

Generally more susceptible the test organism the larger is the zone of inhibition. Antimicrobial activities of the test compounds are expressed by measuring the zone of inhibition observed around the area. The diameter of the inhibition zone is usually measured to understand the extent of inhibition in different concentration.

The size of the inhibitory zones depends principally in the following factors:

- (a) Intrinsic antimicrobial sensitivity of the test compounds.
- (b) Growth rate of the test microorganisms.
- (c) Diffusion rate of the drug, which is related to its water solubility.
- (d) Number of concentration of inoculated test organisms.
- (e) Concentration / amount of the test sample.
- (f) Thickness of the test medium in the Petridishes.

3.1.3 Chemicals used:

Eight synthesized chalcones were used as the test chemicals. The chemicals were synthesized, isolated and characterization in the organic research laboratory, Department of chemistry, University of Rajshahi. The names of the tested chemical are listed in Table 2.

Table 2: List of synthesized chalcones used for antibacterial activities:

Compound no.	Test compounds	Molecular formula
3a	2'-hydroxy chalcone	C ₁₅ H ₁₂ O ₂
3b	2'-hydroxy-4 -methoxy chalcone	C ₁₆ H ₁₄ O ₃
3c	2'-hydroxy-2, 4, 5-trimethoxy chalcone	C ₁₈ H ₁₈ O ₅
3d	2', 5'-dihydroxy-2,4,6-trimethoxychalcone	C ₁₈ H ₁₈ O ₆
3e	4-hydroxy-3', 4', 5' - trimethoxy chalcone	C ₁₈ H ₁₈ O ₅
4a	2'-hydroxy-3'-C-prenyl Chalcone	C ₂₀ H ₂₀ O ₂
4b	2'-hydroxy-5'-C-prenyl Chalcone	C ₂₀ H ₂₀ O ₂
4c	2',5'-dihydroxy-2,4,6-trimethoxy-3'-C-prenylchalcone	C ₂₃ H ₂₆ O ₆

*Ciprofloxacin-50 (*C-50, 50µg/disc) was used as standard.

3.1.4 Test organism:

In the present study of screen the antibacterial activities of chemicals (Table 2) and a number of bacterial strains were used as test organisms. Among these four pathogens were tested (two Gram-positive and two Gram-negative).

3.1.5 Collection of organism:

The test tubes cultures of the bacterial pathogens were collected from the pharmaceutical Microbiological laboratory, Department of pharmacy, University of Rajshahi, Rajshahi Bangladesh. Types of test organism have been studied are listed below (Table 3).

Table 3: List of the test microorganisms (bacteria)

Label	Name of microorganism	Gram-positive (G ⁺) / negative (G ⁻)
B ₁	<i>Bacillus caerius</i>	(G ⁺)
B ₂	<i>Staphylococcus aureus</i>	(G ⁺)
B ₃	<i>Escherichia coli</i>	(G ⁻)
B ₄	<i>Agrobacterium Species</i>	(G ⁻)

3.1.6 Description of procedure:

3.1.6.1 Sterilization procedure:

Antimicrobial screening was carried out in Laminar air flow unit and all types of precaution were highly maintained to avoid any contamination during the test. UV light has switched on before working in laminar hood for 1 hour to avoid any accidental contamination. Petri dishes and other glass

wares were sterilized by autoclaving at a temperature of 121 °C and a pressure of 15 lbs/sq inch for 20 minutes. Micropipette tips culture media; cotton forceps blank discs etc. were also sterilized.

3.1.6.2 Culture media:

The following media were used to demonstrate the antimicrobial activity and for subculture of the test organism.

- i) Nutrient agar medium
- ii) Nutrient broth medium

Among these, the first one is most frequently used and we also used the same for the antibacterial screening.

3.1.6.3 Preparation of nutrient agar:

The constituents of nutrient agar was accurately weighed and dispersed in waters. It was then placed in a water bath to dissolve the ingredients until a transparent solution was obtained.

3.1.6.4 Preparation of medium:

The instant Nutrient Agar (NA) media as weighed and then reconstituted with distilled water in a conical flask according to specification (2.8% w/v). It was then heated in a water bath to dissolve the agar until a transparent solution was obtained and it was then autoclaved at 121 °C for 15 minutes at 15 lbs / sq. inch pressure.

3.1.6.5 Preparation of fresh culture:

The media prepared in the above section was dispensed in 5.0 mL amount of clear test tubes to prepare slants. The test tubes were plugged with cotton and sterilized in an autoclave at 121 °C for 15 minutes at 15 lbs/sq. inch pressure. After sterilization the test tubes were kept in an inclined position for solidification. Finally the slants streaked with pure culture of the test organisms under laminar air flow and incubated for 24 hrs.

3.1.6.6 Preparation of test plates:

- i) 15 mL of Nutrient Agar medium which prepared the previous section was poured in clean test tubes and then plugged with cotton.
- ii) The test tubes were sterilized and allowed to cool at about 45 °C to 50 °C.
- iii) The medium in the test tubes were inoculated with fresh culture of the test bacteria by means of a sterile loop and agitated to ensure uniform dispersion of bacteria in the medium.
- iv) Finally, the medium was poured into sterile petridishes and agitated and clockwise anticlockwise right to left and left to right. Thus plates were ready for sensitivity test.

3.1.6.7 Preparation of discs containing sample:

3.1.6.7.1 Sample discs:

- i) Solution of the synthetic compounds was prepared in acetone so that 170 μ L contained 1mg of the compounds.

- ii) Filter paper discs were taken in a petridish and sterilized by oven at 110 °C for 1 hour.
- iii) 40 µL of the solution of each compound was placed in a particular disc with the help of a micropipette.
- iv) These discs were then air dried.

3.1.6.7.2 Standard discs:

These were used as positive control to ensure the activity of standard antibiotic against the test organism as well as for comparison of the response produced by the known antibacterial agent with that produced by test samples. In our investigation, commercially available Ciprofloxacin-50 containing 50 µg/disc of this antibiotic was used as standard disc.

3.1.6.8 Placement of the discs and incubation:

- i) The sample impregnated discs and standard antibiotic discs were placed gently on the solidified agar plates with the help of a sterile forceps to ensure contact to the medium.
- ii) The plates were then kept in a refrigerator at 4 °C for overnight so that the materials that absorbed the discs could get sufficient time to diffuse into the medium.
- iii) Finally the plates were incubated at 37.5 °C for 24 hrs.

3.1.6.9 Precaution:

The discs were placed in such a way that they were not closer than 15 mm to the edge of the plate and far enough apart to prevent overlapping the zones of inhibition.

3.1.6.10 Determination of antibacterial activity of the test agents by measuring the zone of inhibition:

After 24 hrs. incubation the antibacterial activity was carried out by measuring the zone of inhibition in millimeter (mm) by a transparent scale. The zones made by the samples were compared with that of the standard disc. The results are presented in tabular form in Chapter 4.

3.2 Section B; General Discussion on Antioxidant activity ^[11, 12]

In living systems, free radicals are generated as part of the body's normal metabolic process and the free radical chain reactions are usually produced in the mitochondrial respiratory chain, liver mixed function oxidases, through xanthine oxidase activity, atmospheric pollutants and from transitional metal catalysts, drugs and xenobiotics. In addition, chemical mobilization of fat stores under various conditions such as lactation, exercise, fever, infection and even fasting, can result in increased radical activity and damage. Free radicals or oxidative injury now appears the fundamental mechanism underlying a number of human neurologic and other disorders. Oxygen free radical can initiate peroxidation of lipids, which in turn stimulates glycation of protein, inactivation of enzymes and alteration in the structure and function of collagen basement and other membranes, and play a role in the long-term complication of diabetes. Antioxidants may be defined as radical scavengers which protect the human body against free

radicals that may cause pathological conditions such as ischemia, anaemia, asthma, arthritis, inflammation, neurodegeneration, Parkinson's diseases, mongolism, aging process and perhaps dementias. [28]

The main characteristic of an antioxidant is its ability to trap free radicals. Highly reactive free radicals and oxygen species are present in biological system from a wide variety of sources. These free radicals may oxidize nucleic acids, proteins, lipids or DNA and can initiate degenerative disease. Antioxidant compounds like phenolic acids, polyphenols and flavonoids scavenge free radicals such as peroxide, hydro peroxide or lipid peroxy and thus inhibit the oxidative mechanisms that lead to degenerative diseases.

Oxidation is the transfer of electrons from one atom to another. It represents an essential part of metabolism and aerobic life in general, since oxygen is the ultimate electron acceptor in the electron flow systems that transport energy in the form of ATP. [29] Problems may arise, however, when the electron flow generates free radicals, such as O_2^- centered free radicals, known as reactive oxygen species (ROS), and including superoxide (O_2^-), peroxy ($ROO\cdot$), alkoxy ($RO\cdot$), hydroxyl ($HO\cdot$), nitric oxide ($NO\cdot$) radicals. The contribution of free radical-mediated processes to the pathogenesis of human disease is indicated by biomarkers of oxidative damage to lipids, protein and DNA. Such markers have been identified in patients with atherosclerosis, certain cancer, neurodegenerative diseases and lung disorders, especially those with an inflammatory component to their etiology. A range of oxygen species (ROS) and reactive nitrogen species (RNS) have been implicated in the mechanisms of damage associated with disease development, including superoxide radical (O_2^-), hydrogen peroxide ($HOO\cdot$), lipid alkoxy ($RO\cdot$), and peroxy radicals ($ROO\cdot$), peroxy nitrite ($ONOO\cdot$), nitric oxide ($NO\cdot$) and nitrogen dioxide radical ($NOO\cdot$).

Antioxidants are compounds capable of preventing and even counteracting the damage caused in human tissue by the normal effects of physiological oxidation. A lot of research has shown that antioxidant can play a role in preventing the development of some chronic diseases. In addition to those mentioned previously, diseases such as atherosclerosis, emphysema, iron overload, malaria, muscular dystrophy, retinal degeneration and rheumatoid arthritis are but a few examples where research has shown the likelihood of direct links and the possibility of positive dietary and perhaps even nutraceutical intervention. Antioxidant activity and nutritional labeling data including vitamins, fiber, and minerals will aid in the interpretation of clinical results obtained as various food products are tested in biological models for chronic disease. It is reasonable to expect that high antioxidant foods have greater potential to reduce free radicals in the body than do low antioxidant foods. Thus it is important to know the antioxidant content of foods, in addition to knowing the basic nutritional information such as the protein, fiber, fat, mineral and vitamin contents.

Antioxidant compounds in food play an important role as a health protecting factor. Scientific evidence suggests that antioxidants reduce the risk for chronic diseases including cancer and heart disease. Primary sources of naturally occurring antioxidants are whole grains, fruits and vegetables. Plant sourced food antioxidants like vitamin C, vitamin E, carotenes, phenolic acids, phytate and phytoestrogens have been recognized as having the potential to reduce disease risk. Most of the antioxidant compounds in a typical diet are derived from plant sources and belong to various classes of compounds with a wide variety of physical and chemical properties. Some compounds, such as gallates, have strong antioxidant activity, while others, such as mono-phenols are weak antioxidants.


Table 4: List of some foods containing high levels of antioxidants³⁰

Food	Antioxidant activity (TE / 100 gm)
Red Grapes	1350
Red Cabbage	1000
Broccoli Flowers	500
Spinach	500
Green Grapes	400
Tomato	300
Green Bean	175
Red Bean	11459
Green Cabbage	150
Lima Beans	1055

Table 5: List of some naturally occurring antioxidant^[11]

Antioxidant compounds	Food containing high levels of these antioxidant
Vitamin C (ascorbic acid)	Fruits and vegetables
Vitamin E (tocopherols, tocotrienols)	Vegetable oils
Polyphenolic antioxidants (resveratrol, flavonoids)	Tea, coffee, soy, fruit, olive oil, chocolate, cinnamon, oregano and red wine
Carotenoids (lycopene, carotene)	Fruit, vegetables and eggs

Vitamin E is very often used and takes as anti aging antioxidant. It functions in the following way through the formation of phenoxy radical combine with other unwanted radical in living cell.



Scheme 15: Free radical scavenging mechanism of vitamin E

It is known that the antioxidant properties of chalcones are quite dependant on the two aryl structures, that is, the substitution pattern on the two aryl rings of the chalcone moiety. Especially, the hydroxyl substituent is one of the key groups that enhance greatly the antioxidant activity of chalcone mainly due to its easy conversion to phenoxy radicals through the hydrogen atom transfer mechanism. This phenoxy radical formation may be crucial to the antioxidant properties, which are assessed primarily as radical scavenging potential of phenolic chalcones. In fact, the hydroxyl substituent is common among chalcones from natural sources.

Chalcones basic structure includes two aromatic ring bound by α , β -unsaturated carbonyl group, a unique templates associated with diverse application. Due to the presence of the reactive keto, vinylic group, chalcones and their analogues have been reported to be antioxidant. Hydroxyl, methoxy and prenyl substituents are associated with antioxidant properties.

3.2.1 Study of antioxidant property by some common methods: ^[11, 30]

Various antioxidant activity methods have been used to monitor and compare the antioxidant activity of foods. In recent years, oxygen radical absorbance capacity assay and enhanced chemiluminescence assay have been used to evaluate antioxidant activity of foods, serum and other biological fluids. These methods require special equipment and technical skills for the analysis. The different types of methods published in the literature for the determinations of antioxidant activity of food involve electron spin resonance (ESR) and chemiluminescence methods. These analytical methods measure the radical scavenging activity of antioxidants against free radicals like the 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical, the superoxide anion (O_2^-), the hydroxyl radical (HO^\cdot) or peroxy radicals (ROO^\cdot). The various methods used to measure antioxidant activity of food products can give varying results depending on the specific free radical being used as a reactant. There are other methods which determine the resistance of lipid or lipid emulsions to oxidation in the presence of antioxidant being tested. The malondialdehyde (MDA) or thiobarbituric acid-reactive-substances (TBARS) assay have been used extensively since the 1950's to estimate the peroxidation of lipids in membrane and biological system. These methods can be time consuming because they depend on oxidation of a substrate which is influenced by temperature, pressure, matrix etc. and may not be practical when large numbers of samples are involved. Antioxidant activity methods using free radical traps are relatively straightforward to perform. The ABTS [2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid)] radical cation has been used to screen the relative radical- scavenging abilities of flavonoids and phenolics have used the Oxygen Radical Absorbance Capacity (ORAC) produce to determine antioxidant capacities of fruits and vegetables^[34] In the ORAC method, a sample is added to the peroxy radical generator, 2, 2'-

azinobis (2- aminopropane) dihydrochloride (AAPH) and inhibition of the free radical action is measured using the fluorescent compound, B-phycoerythrin necessary to react with one half of the DPPH is expressed in terms of micromole equivalents of Trolox (TE) per 100 grams of sample, or simply Trolox units per 100 gm or TE/100g.

Table 6: List of some commonly used antioxidant used as standard

Standard antioxidants	Antioxidant activity (TE/100 Grams)
Ascorbate	442,000
Trolox	400,000
Vitamin E	201,000
BHT	395,000

3.2.1.1 DPPH Radical Scavenging Activity

The radical scavenging activity was measured using the method of Hatano *et al.* with some modifications. Summarily, 0.2 mL solution of DPPH in methanol was prepared and 1.5 mL of this solution was added to the equal volume of each test samples dissolved in methanol at different concentrations. After shaking, the mixture was maintained in dark for 30 mins. Then, the absorbance was measured at 517 nm against a blank. Ascorbic acid and butylated hydroxyanisole (BHA) were used as standard references. The scavenging activity was calculated using the formula:

$$\text{Scavenging Activity (\%)} = [(A_{517} \text{ of control} - A_{517} \text{ of sample}) / A_{517} \text{ of control}] \times 100.$$

3.2.1.2 Ferrous ion Chelating Activity:

Ferrous iron-ferrozine complex method with some modification was used for determination of chelating activity of samples for ferrous ions Fe^{2+} . Briefly, 25 μL of FeCl_2 solution (2 mM) was added to a mixture containing 1.5 mL of H_2O and 2 mL of the test samples in methanol at different concentrations. The reaction was preceded by adding 50 μL ferrozine solution (5mM) after 30 seconds. The mixture was shaken well and incubated for 10 mins. at room temperature. Absorbance of the solution was then measured at 562 nm Quercetin was used as positive control. The ability of the extracts and fractions to chelate ferrous ion was calculated using the equation described above for DPPH.

3.2.1.3 Inhibition of β -carotene Bleaching:

Antioxidant activity of the samples was determined according to a slightly modified version of the β -carotene bleaching method. In this study 5mg of β -carotene was dissolved in 10 mL of chloroform. 33 μL of linoleic acid, 750 μL of β -carotene solution and 225 mg of Tween 40 were mixed. The solvent was completely removed using a rotary evaporator. Then 75 mL of oxygenated distilled water was added and the mixture was emulsified for 15 mins. in a sonicator to form an emulsion A. Aliquots of 3.5 mL of this were transferred into a series of stopper test tubes containing 1 mL of samples dissolved in ethanol in various concentrations. Optical density (OD) at 470 nm was determined for all samples immediately ($t=0$) and at the end of the time period ($t=120$). A second emulsion was also prepared and used as blank to zero the spectrophotometer. This emulsion consisted of 50 mL of oxygenated water, 22 μL of linoleic acid and 150 mg of Tween 40. The percentage inhibition was calculated according to the following formula:

$$\text{Inhibition (\%)} = [(AA (120) - AC (120)) / (AC (0) - AC (120))] \times 10$$

In which AA (120) is the absorbance of the sample at $t = 120$ mins. AC (120) is the absorbance of the control at $t = 120$ mins. and AC (0) is the absorbance of the control at $t = 0$ mins.

3.2.1.4 Trolox Equivalent Antioxidant Capacity (TEAC):

Antioxidant capacity of the samples were measured with some modifications. In this procedure ABTS (54.2 mg) was dissolved in phosphate buffer (pH = 7) and activated to ABTS + radical by addition of 100 mg MnO_2 with stirring for 30 mins. Then, the solution was centrifuged (5 min, 10000 rpm), filtered (0.22 μ m) and diluted with phosphate buffer so that $A_0 = 0.7$. Then, 2 mL of ABTS + radical was added to test tubes containing 1 mL of samples dissolved in ethanol in various concentrations. Time of reaction was 20 mins. Absorbance of the solution were measured at a wavelength of 734 nm and antioxidant capacity of the samples were calculated according to the following formula:

$$\text{Inhibition (\%)} = [(A_0 - A_1)/A_0] \times 100.$$

Where A_0 = the absorbance of the control, A_1 = the absorbance of the sample. Trolox was used as standard.

3.3 Radical Scavenging Activity Determination of the Present Study by DPPH Method:

DPPH assay has been extensively used for screening antioxidant activity because it can accommodate many samples in a short period and is sensitive enough to detect active ingredients at low concentration. When DPPH radicals encounter a proton donating substance such as an antioxidant,

it would be scavenged and the absorbance is reduced. Thus, the DPPH radicals were widely used to investigate the scavenging activity of some natural compounds.

3.3.1 Principle:

The 1, 1-diphenyl -2- picrylhydrazyl radical has been widely used to evaluate the free radical scavenging capacity of antioxidants. DPPH free radical is reduced to the corresponding hydrazine when it reacts with hydrogen donors. DPPH can make stable free radicals in mucous or methanol solution. With this method it was possible to determine the antiradical power of an antioxidant activity by measurement of the decrease in the absorbance of DPPH at 517 nm. Resulting from a color change from purple to yellow the absorbance decreased when the DPPH was scavenged by an antioxidant, through donation of hydrogen to form a stable DPPH molecule. In the radical form this molecule had an absorbance at 517 nm which disappeared after acceptance of an electron or hydrogen radical from an antioxidant compound to become a stable diamagnetic molecule




Figure 3: Structure of 1, 1-diphenyl-2-picrylhydrazyl

Strong absorption maximum at 517 nm and is purple in color. The color turns from purple to yellow as the molar absorptivity of the DPPH radical at 517 nm reduces from 9660 to 1660 when the odd electron of DPPH

radical becomes paired with hydrogen from a free radical scavenging antioxidant to form the reduced DPPH-H. The resulting decolorization of stoichiometric with respect to number of electrons captured.

3.3.2 Materials:

i) Apparatus used:

- ✓ Micropipette (1-10 mL)
- ✓ UV spectrophotometer
- ✓ Electronic balance
- ✓ Test tube
- ✓ Beaker

ii) Chemical used:

- ✓ Ascorbic acid
- ✓ DPPH (1, 1-diphenyl-2-picrylhydrazyl)
- ✓ Methanol

3.3.3 Experimental procedure:

Firstly, 5 test tubes were taken to make aliquots of five kinds (2, 5, 10, 15, 20 $\mu\text{g/mL}$) of concentrated solution. Test sample and ascorbic acid were weighed 3 times and dissolved in methanol to make the required concentration by dilution technique. DPPH was weighed and dissolved in methanol to make .02 % solution. To dissolve homogeneously, a magnetic stirrer was used. After making the desired concentration 2 μL of DPPH solution was applied on each test tube by pipette. DPPH was also applied on the blank test tube at the same time where only methanol was taken as blank. The test tube was incubated at room temperature for 30 mins. in dark place to

complete the reaction. Then the absorbance of the solution was measured at 517 nm using a spectrophotometer against blank. Ascorbic acid was used as positive control.

The percentage (%) of scavenging was calculated from the following equation.

$$\% \text{ of scavenging} = [(A_0 - A_1) / A_0] \times 100.$$


Where,

A_0 = the absorbance of the control, and

A_1 = the absorbance of the sample.

Then percent of scavenging were plotted against concentration using MS Excel software. Inhibitory concentration IC_{50} value was calculated from the graph which indicates value conc. ($\mu\text{g/mL}$) at 50% inhibition. Percent inhibition, IC_{50} values are presented in tabular form in Chapter 4 and activity is discussed.

A Flowchart of Determination of DPPH radical scavenging activity




Scheme 16: Determination of DPPH radical scavenging activity

3.4 Reducing power capacity:

The reducing power of chalcones was determined by Oyaizu method. [31] It has been reported that there is a direct correlation between antioxidant activities and reducing power of certain compounds. The reducing properties

A Flowchart of Determination of DPPH radical scavenging activity




Scheme 16: Determination of DPPH radical scavenging activity

3.4 Reducing power capacity:

The reducing power of chalcones was determined by Oyaizu method.
^[31] It has been reported that there is a direct correlation between antioxidant activities and reducing power of certain compounds. The reducing properties

are generally associated with the presence of reductants, which have been shown to exert antioxidant action by breaking the free radical chain by donating a hydrogen atom. The presence of reductants such as antioxidant substances in the samples causes the reduction of the Fe^{2+} - ferricyanide complex to the ferrous form by donating an electron. The amount of Fe^{2+} complex can then be monitored by measuring the formation of Perl's Prussian blue at 700 nm.



3.4.1 Reagents


- i. Potassium ferricyanide (MERCK, Germany)
- ii. Trichloro acetic acid (MERCK, Germany)
- iii. Ferric Chloride (sigma chemical company, USA)
- iv. Ascorbic acid (sigma chemical company, USA)
- v. Phosphate Buffer (sigma chemical company, USA)

3.4.2 Procedure

- i. 0.25 mL of samples and standard of different concentrations solution were taken in a test tube.
- ii. 0.625 mL of sodium buffer (2M) and 0.625 mL of potassium ferricyanide were added to them and vortexed well.
- iii. The reaction mixture was incubated at 50 °C for 10 mins. and 0.625 mL of trichloro acetic acid was added.
- iv. The reaction mixture was centrifuged at 3000 rpm for 10 mins.
- v. 1.8 mL supernatant solution was withdrawn and mixed with 1.8 mL of distilled water.
- vi. 0.36 mL of ferric chloride was added to dilute the reaction mixture.

- vii. Then the absorbance of the solution was measured at 700 nm using a spectrophotometer against a blank.
- viii. A typical blank solution containing the same solution mixture without sample or standard was prepared and incubated under the same condition as the rest of the sample solution.

A Flowchart of Determination Reducing power capacity




Scheme 17: Determination of Reducing power capacity

3.5 Cytotoxicity Bioassay ^[32]


In vivo lethality assay in a simple zoological organism, such as Brine shrimp lethality test (BST) has been applied as a simple and useful tool for preliminary screening^[35, 36] of toxicity of physiologically active plant extract or synthesized compounds, detection of fungal toxins, heavy metal, pesticides and also cytotoxicity testing of dental materials.^[35, 37] This general bioassay is rapid, reliable and has been used for over thirty years in toxicological studies. However, it has been demonstrated that a positive relationship exists between brine shrimp lethality and human carcinoma. Thus, BST can also be extrapolated for cell line toxicity and anti tumor activity. Principal of this method was based on the ability of certain compounds to kill laboratory cultured *Artemia* nauplii brine shrimp. BST is one of the simplest biological responses to monitor is lethality, since there is only one criterion: either dead or alive. It has been shown that *Artemia* is highly vulnerable to toxins at early development stages and assumed to exhibit their greatest sensitivity to test compounds.^[35] Subsequently, in this study we used 24 hrs. nauplii as object experimental. Brine shrimp lethality bioassay was carried out to investigate cytotoxicity of the synthesized chalcones. Here *in vivo* lethality test were carried out using brine shrimp nauplii eggs (*Artemia salina* Lech). Eggs were placed in a small tank containing 3.8% NaCl solution for hatching. After 48 hours of hatching period the nauplii were ready for the experiment. After hatching, active nauplii free from egg shells were collected with Pasteur pipette from tank and ready to be used for the assay. Then 5mg of each synthesized compound was accurately measured and dissolved in 1 mL of DMSO to get a concentration of 5mg/mL. From the stock solutions, a variety of solution conc. were prepared as 2.5, 5, 10, 20, 40 and 80 $\mu\text{g/mL}$ (ppm) respectively. 10 Brine shrimp nauplii were then placed in each test tube using Pasteur pipette. For the control test of each test tube containing the same

volume of DMSO plus sea water up to 5 mL was used. After 24 hrs. of incubation the test tubes were observed using a magnifying glass and the number of survivors in each test tube were counted, noted and calculating percentages of death. Larvae were considered dead if they did not show any movement during several seconds of observation. ^[35] Lethal dose 50% (LC₅₀) values for the tested compounds were calculated from the graph. LC₅₀ value greater than 1000 ppm for plant extracts was considered inactive, whereas LC₅₀ values less than 30 ppm for pure compounds were considered toxic.

A Flowchart of Determination of cytotoxicity of chalcone



Scheme 18: Determination of cytotoxicity of chalcone



CHAPTER FOUR
RESULT AND DISCUSSION

CHAPTER FOUR


Results and Discussion

4.1 Section A; Synthesis and Structure Elucidation

Part- I of this dissertation is concerned with the synthesis and structural elucidation of some hydroxyl, methoxy and prenyl chalcones e.g. 2' - hydroxychalcone (**3a**), 2' - hydroxy - 4 - methoxychalcone (**3b**), 2' - hydroxy - 2, 4, 5 - trimethoxychalcone (**3c**), 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy chalcone (**3d**), 4 - hydroxy - 3', 4', 5' - trimethoxychalcone (**3e**), 2' - hydroxy - 3' - C - prenylchalcone (**4a**), 2' - hydroxy - 5' - C - prenylchalcone (**4b**), 2', 5' - dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (**4c**).

One of the starting materials β -resacetophenone (**1b**) has been prepared and confirmed by literature melting point, R_f value and IR spectrum.

Among the above 2' - hydroxychalcones **3a-3e** were synthesized by Claisen-Schmidt condensation according to scheme 7 using substituted acetophenone (**1a -1c**) and substituted benzaldehyde (**2a-2e**). The completion of reaction was monitored by using silica gel thin layer chromatography (TLC). The compounds were purified by recrystallation, PTLC or column chromatography.



Scheme 7: Synthetic route for five synthesized chalcones (**3a-3e**) from their corresponding acetophenone (**1a-1c**) and aldehyde (**2a-2e**)

3a and **3d** were also undertaken by prenylation to produce their prenylated derivatives according to scheme 13 and scheme 14

Chalcones, a group of compounds with two aromatic rings connected by a keto-vinyl chain, $-\text{CO}-\text{CH}=\text{CH}-$ so these have some unique structural features in spectroscopy.

The oxygenated chalcones usually possess UV absorption maxima in the range of 340-390 nm and chalcones lacking B-ring oxygenation may have their absorption at considerably shorter wavelengths and a minor peak usually appears in the range of 220-270 nm.

The infrared spectra of chalcones show usually a band near 1625-1650 cm^{-1} , characteristic of an α , β -unsaturated carbonyl group.

The α -H and β -H of chalcones resonate at δ 6.7 -7.4 and δ 7.3 -7.7 as two doublets ($J = 16-17$ Hz) respectively in the ^1H NMR spectra. This large J value shows that the olefinic bond has *trans* geometry. In the ^{13}C NMR

spectra of chalcones, the carbonyl carbon appears between δ 188.6 and 194.6. The α and β carbon atoms give rise to signals in between δ 116.1-128.1 and δ 136.9-145.4 respectively and can be readily identified by their characteristic appearance as a six line multiplet in the off resonance. The important spectral values are presented in tabular form (Table 7) and characteristics features were noticed.

Table 7: List of spectral data for some important functionality for compounds (3a-3e, 4a-4c)

Comp no.	UV λ_{\max} (CH ₃ OH) nm	IR (KBr, cm ⁻¹)	¹ H and ¹³ C NMR (δ ppm)
3a	196, 233	3414.06 (-OH), 3046.91 (C=C-H, olifinic, str.), 1640.01 (C=O conj. keto), 1618.00 (C=C, arm.)	12.800 (s, 1H, C ₂ OH), 7.920 (d, J = 14.8 Hz, 1H, C _{β} -H), 7.660 (d, J =14.8 Hz, 1H, C _{α} -H) 193.77 (1C, >C=O), 145.49 (1C, C _{β}), 120.17 (1C, C _{α})
3b	197	3414.07 (-OH), 3024.51 (C=C-H, str.), 1640.50 (C=O), 1609.00 (C=C, arm. str.)	12.800 (s, 1H, C _{2'} -OH), 7.910 (d, J =16 Hz, 1H, C _{β} -H), 7.540 (d, J =16 Hz, 1H, C _{α} -H), 3.870 (s, 3H, C ₄ -OCH ₃) 192.80 (1C, >C=O), 145.10 (1C, C _{β}), 118.70 (1C, C _{α}), 55.80 (1C, C ₄ -OCH ₃)
3c	196, 259, 307	3416.30 (-OH), 3009.49 (C=C-H str.), 1660.01 (C=O), 1609.36 (C=C, arm.)	13.080 (s, 1H, C ₂ -OH), 8.170 (d, J =16 Hz, 1H, C _{β} -H), 7.390 (d, J =16 Hz, 1H, C _{α} -H), 3.730 (s, 9H, C ₂ , C ₄ , C ₅ -OCH ₃) 187.01 (1C, >C=O), 142.80 (1C, C _{β}), 121.61 (1C, C _{α}), 56.40 (C ₂ -OCH ₃), 56.30 (2C, C ₄ and C ₅ -OCH ₃)

3d	196, 242, 384	3415.26 (-OH), 3000.50 (-C=C-H, str.), 1605.56 (C=O), 1548.81 (C=C, arm.)	12.770 (s, 2H, C _{2'} and C _{5'} -OH), 8.360 (d, J =15.6 Hz, 1H, C _β -H), 7.910 (d, J =16 Hz, 1H, C _α - H), 3.920 (s, 6H, C ₂ and C ₆ - OCH ₃), 3.860 (s, 3H, C ₄ -OCH ₃) 194.89 (1C, >C=O), 159.00 (1C, C ₂ -OH), 157.69 (1C, C ₅ -OH), 147.18 (1C, C _β), 123.93 (1C, C _α), 55.85 (2C, C ₂ and C ₆ - OCH ₃), 55.40 (1C, C ₄ -OCH ₃)
3e	197	3413.88 (-OH), 3232.28 (-C=C-H, olifinic str.), 1640.58 (C=O, conj. keto), 1610.44 (C=C, arm.)	9.850 (s, 1H, C ₄ -OH), 7.770(d, J = 14.8 Hz, 1H, C _β -H), 7.340 (d, J =15.6 Hz, 1H, C _α - H), 3.90 0(s, 9H, C _{3'} , C _{4'} and C _{5'} - OCH ₃) 189.67 (1C, >C=O), 158.35 (1C, C ₄ - OH), 144.94 (1C, C _β), 119.34 (1C, C _α), 60.97(1C, C ₄ -OCH ₃), 56.38 (2C, C _{3'} and C _{5'} -OCH ₃)
4a	197		7.830 (d, J =16.0 Hz, 1H, C _β -H), 7.680 (d, J =16.0 Hz, 1H, C _α -H), 5.500 (s, 1H, OH), 4.050-3.900 (m, 1H, CH ₂ -CH), 3.400-3.200 (m, 2H, CH ₂) 1.800 (s, CH ₃ , 6H) 192.80(1C, >C=O), 161.50(1C, C ₂ OH), 145 .10 (1C, C _β), 118.70 (1C, C _α), 24.60 (1C, -CH ₂ -CH=C ^{CH₃} _{CH₃}) 22.60 (1C, CH ₂ -CH) 18.60 (1C, -CH ₂ -CH=C ^{CH₃} _{CH₃})

4b	196, 241		<p>7.930 (d, $J=15$ Hz, 1H, C$_{\beta}$-H), 7.550 (d, $J=15$ Hz, 1H, C$_{\alpha}$-H), 5.500 (s, 1H, OH), 5.200 (s, 1H, CH$_2$-CH), 3.700-3.600 (m, 2H, CH$_2$), 1.800 (s, 3H, -CH$_2$-CH=C(CH$_3$)), 1.740 (3H, -CH$_2$-CH=C(CH$_3$))</p> <p>192.80(1C,$>$C=O), 160.60(1C, C$_2$-OH), 145.10(1C, C$_{\beta}$), 123.00(CH$_2$-CH), 118.700(1C, C$_{\alpha}$), 24.60 (1C, -CH$_2$-CH=C(CH$_3$)) 22.60(1C, CH$_2$), 18.60(1C, -CH$_2$-CH=C(CH$_3$))</p>
4c	196		<p>8.330 (d, $J=16.0$ Hz, 1H, C$_{\beta}$-H), 7.420 (d, $J=16.0$ Hz, 1H, C$_{\alpha}$-H), 5.350 (s, 2H, C$_2'$ and C$_5'$-OH), 5.750 (1H, CH$_2$-CH), 3.210 (2H, CH$_2$), 1.820 (s, CH$_3$, 6H), 1.820 (s, 3H, -CH$_2$-CH=C(CH$_3$)), 1.700 (s, 3H, -CH$_2$-CH=C(CH$_3$))</p> <p>192.80(1C,$>$C=O), 154.10(1C, C$_2$-OH), 122.50(1C, C$_{\beta}$), 123.00(CH$_2$-CH), 121.30(1C, C$_{\alpha}$), 24.60 (1C, -CH$_2$-CH=C(CH$_3$)) 28.10(1C, CH$_2$), 18.60(1C, -CH$_2$-CH=C(CH$_3$))</p>

4.2 Section B; Results for Antimicrobial Screening

Chalcones are a group of natural products found in fruits, vegetables, nuts, seeds and flowers as well as in teas and wines are important constituents of human diet. They have been demonstrated to possess many biological and pharmacological activities such as antibacterial, antifungal, antiviral, antioxidant, anti-inflammatory, antimutagenic and antiallergic activities and inhibitory activities on several enzymes. [5,33]

In the present study to screen the antibacterial activities all the compounds (**3a-3e**, **4a-4c**) were employed as test against four bacteria *Bacillus caerius* (G^+ , B₁), *Staphylococcus aureus* (G^+ , B₂), *Escherichia coli* (G^- , B₃), *Agrobacterium Species* (G^- , B₄). The primary assay was performed by disc diffusion technique to classify the microorganism susceptible as well as resistant towards particular compounds. The bioactivity is expressed by the diameter of zone of inhibition in mm. Ciprofloxacin is used as standard drug. The result of Antibacterial screening is presented as tabular form in Table 8

Table 8: Result of the antibacterial activity of the compounds (**3a-3e**, **4a-4c**) against *Bacillus caerius*. (G^+ , B₁) at 250 ($\mu\text{g disc}^{-1}$)

Comp. no	Molecular formula	Diameter of inhibition zone (mm)	*C-50
3a	$\text{C}_{15}\text{H}_{12}\text{O}_2$	24	40
3b	$\text{C}_{16}\text{H}_{14}\text{O}_3$	28	
3c	$\text{C}_{18}\text{H}_{18}\text{O}_5$	30	
3d	$\text{C}_{18}\text{H}_{18}\text{O}_6$	38	
3e	$\text{C}_{18}\text{H}_{18}\text{O}_5$	12	
4a	$\text{C}_{20}\text{H}_{20}\text{O}_2$	08	
4b	$\text{C}_{20}\text{H}_{20}\text{O}_2$	12	
4c	$\text{C}_{23}\text{H}_{26}\text{O}_6$	20	

* Ciprofloxacin-50

The trend for antibacterial (*Bacillus caerius*, G^+ , B_1) activity at conc. $250\mu\text{g disc}^{-1}$ C-50 (40 mm) > 3d (38 mm) > 3c (30 mm) > 3b (28 mm) > 3a (24 mm) > 4c (20 mm) > 3e, 4b (12 mm) > 4a (08 mm)

a) *Bacillus caerius*. (G^+ , B_1) [Figure-4]

The inhibition zones of *Bacillus caerius* due to the treatment of different compounds at the different concentrations are presented in Table 8. All the compounds (3a-3e, 4a-4c) showed inhibition zones at concentration $250 (\mu\text{g disc}^{-1})$. It was found that the inhibition zones of compound 3c and 3d were more effective than that of the other compounds and similar to the standard.

Table 9: Results of the antibacterial activity of the compounds against *Staphylococcus aureus* (G^+ , B_2) at $250 (\mu\text{g disc}^{-1})$

Comp. no	Molecular formula	Diameter of inhibition zone (mm)	*C-50
3a	$\text{C}_{15}\text{H}_{12}\text{O}_2$	20	42
3b	$\text{C}_{16}\text{H}_{14}\text{O}_3$	40	
3c	$\text{C}_{18}\text{H}_{18}\text{O}_5$	20	
3d	$\text{C}_{18}\text{H}_{18}\text{O}_6$	25	
3e	$\text{C}_{18}\text{H}_{18}\text{O}_5$	12	
4a	$\text{C}_{20}\text{H}_{20}\text{O}_2$	08	
4b	$\text{C}_{20}\text{H}_{20}\text{O}_2$	12	
4c	$\text{C}_{23}\text{H}_{26}\text{O}_6$	30	

* Ciprofloxacin-50

The trend for antibacterial (*Staphylococcus aureus*, G^+ , B_2) activity at conc. $250 \mu\text{g disc}^{-1}$

C-50 (42 mm) > 3b (40 mm) > 4c (30 mm) > 3d (25 mm) > 3a, 3c (20 mm) > 3e, 4b (12 mm) > 4a (08 mm)

b) *Staphylococcus aureus* (G^+ , B_2) [Figure-5]

The inhibition zones of *Staphylococcus aureus* due to the treatment of different compounds at the different concentrations are presented in Table 9. It was found that the inhibition zone of compound **3b** was extremely high 40 mm as compared to that of standard (42 mm) compound. Other compounds show significant antibacterial activity.

Table 10: Results of the antibacterial activity of the compounds against *Escherichia coli*. (G^- , B_3) at $250 (\mu\text{g disc}^{-1})$

Comp. no	Molecular formula	Diameter of inhibition zone (mm)	*C-50
3a	$\text{C}_{15}\text{H}_{12}\text{O}_2$	18	40
3b	$\text{C}_{16}\text{H}_{14}\text{O}_3$	30	
3c	$\text{C}_{18}\text{H}_{18}\text{O}_5$	30	
3d	$\text{C}_{18}\text{H}_{18}\text{O}_6$	20	
3e	$\text{C}_{18}\text{H}_{18}\text{O}_5$	06	
4a	$\text{C}_{20}\text{H}_{20}\text{O}_2$	06	
4b	$\text{C}_{20}\text{H}_{20}\text{O}_2$	06	
4c	$\text{C}_{23}\text{H}_{26}\text{O}_6$	26	

* Ciprofloxacin-50

The trend for antibacterial (*Escherichia coli*, G^- , B_3) activity at conc. 250 μg disc⁻¹

C-50 (40 mm) > 3b, 3c (30 mm) > 4c (26 mm) > 3d (20 mm) > 3a (18 mm) > 3e, 4a, 4b (06 mm)

c) *Escherichia coli*. (G^- , B_3) [Figure-6]

The inhibition zones of *Escherichia coli* due to the treatment of different compounds at the different concentrations are presented in Table 10. It was found that the inhibition zone of compound **3b, 3c** (30 mm) show fairly good activity as compared to that of standard (40 mm) compound. Other compounds relatively less susceptible towards this microorganism.

Table 11: Results of the antibacterial activity of the compounds against *Agrobacterium Species* (G^- , B_4) at 250 (μg disc⁻¹)

Comp. no	Molecular formula	Diameter of inhibition zone (mm)	*C-50
3a	$\text{C}_{15}\text{H}_{12}\text{O}_2$	15	42
3b	$\text{C}_{16}\text{H}_{14}\text{O}_3$	20	
3c	$\text{C}_{18}\text{H}_{18}\text{O}_5$	15	
3d	$\text{C}_{18}\text{H}_{18}\text{O}_6$	24	
3e	$\text{C}_{18}\text{H}_{18}\text{O}_5$	07	
4a	$\text{C}_{20}\text{H}_{20}\text{O}_2$	08	
4b	$\text{C}_{20}\text{H}_{20}\text{O}_2$	08	
4c	$\text{C}_{23}\text{H}_{26}\text{O}_6$	40	

* Ciprofloxacin-50

The trend for antibacterial (*Agrobacterium Species*, G^- , B_4) activity at conc. $250 \mu\text{g disc}^{-1}$

C-50 (42 mm) > 4c (40 mm) > 3d (24 mm) > 3b (20 mm) > 3a, 3c (15 mm) > 4a, 4b (08 mm) > 3e (07 mm)

d) *Agrobacterium Species* (G^- , B_4) [Figure-7]

The inhibition zones of *Agrobacterium Species* due to the treatment of different compounds at the different concentrations are presented in Table 11. All the compounds show better activity specially **4c** (40 mm) against the microorganism.



Figure 4: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against *Bacillus caerius* (G^+ , B_1)



Figure 5: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against *Staphylococcus aureus* (G^+ , B_2)




Figure 6: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against *Escherichia coli* (G^- , B₃).



Figure 7: Photographic representation of the zone of inhibition at the concentration of $250 \mu\text{g disc}^{-1}$ against *Agrobacterium Species* (G^- , B₄)

3b, **3c** and **3d** exhibited fairly good potentialities against both G^+ and G^- bacteria and in some cases about near to that of the standard drugs. Prenyl derivative of compound **3d**, compound **4c** shows very high activity against (G^-) pathogen than other prenyl derivatives. This led us to conclude that the presence of electron releasing hydroxyl ($-\text{OH}$) and methoxy groups ($-\text{OCH}_3$) are responsible for the better antimicrobial effects. **4c** contains lipophilic prenyl group and enhance cell membrane permeability thus more susceptible to (G^-). In **3e** three methoxy groups are occupied ring A and in three adjacent carbons. So steric effect may be a cause of poor activity.

4.3 Antioxidant Activity Screening by DPPH method

A simple method that has been developed to determine the antioxidant activity of foods utilizes the stable 1, 1-diphenyl-2 picrylhydrazyl (DPPH) radical. The electron in the DPPH free radical gives a strong absorption maximum at 517 nm and purple in color in methanolic solution. The color turns purple to yellow as the molar absorptivity of the DPPH radical at 517 nm reduces from 9660 to 1640 when odd electron of DPPH radical becomes paired with hydrogen from a free radical scavenging antioxidant to form the reduced DPPH-H. The resulting decolorization is stoichiometric with respect to no. of electrons captured. The activity is measured in terms of % inhibition and minimum inhibitory concentration IC_{50} from graph. The % scavenging and IC_{50} values are shown in Table 12.

Table 12: DPPH radical scavenging data of synthesized chalcones (**3a-3e**, **4a**, **4c**) and their corresponding IC_{50} values

Compound no.	Conc. $\mu\text{g}/\text{mL}$	Absorbance at 517 nm	% Inhibition	* IC_{50}
3a	2	0.210	23.36	54.94
	5	0.203	25.91	
	10	0.191	30.29	
	15	0.189	31.02	
	20	0.186	32.12	
3b	2	0.184	32.85	36.80
	5	0.180	34.31	
	10	0.178	35.04	
	15	0.165	39.78	
	20	0.160	41.61	

3c	2	0.109	10.60	74.73
	5	0.233	14.96	
	10	0.223	18.61	
	15	0.216	21.17	
	20	0.220	19.71	
3d	2	0.120	56.20	0.923
	5	0.110	59.65	
	10	0.093	66.06	
	15	0.082	70.07	
	20	0.076	72.26	
3e	2	0.222	18.98	93.01
	5	0.219	20.07	
	10	0.216	21.17	
	15	0.210	23.36	
	20	0.205	25.18	
4a	2	0.221	19.34	65.67
	5	0.213	22.26	
	10	0.208	24.08	
	15	0.201	26.64	
	20	0.197	28.10	
4c	2	0.252	8.030	103.31
	5	0.240	12.41	
	10	0.235	14.23	
	15	0.232	15.33	
	20	0.230	16.06	

*Ascorbic acid is the standard and IC_{50} 0.08 ($\mu\text{g}/\text{mL}$). The Table excludes the value of **4b**.




Figure 8: Graph showing DPPH radical scavenging of synthesized chalcones (3a-3e, 4a, 4c) without standard Ascorbic acid.




Figure 9: Comparison of $(IC_{50} + 5)$ value of Ascorbic acid and synthesized chalcones (3a-3e, 4a, 4c). 5 is added with original IC_{50} to display the smaller values in the graph.

Generally smaller the IC_{50} value higher the antioxidant activity thus the compound **3d** showed highest activity (**0.923**). As it contains two Phenolic -OH group and suppose to produce phenoxide free radical easily and stabilized by electronic group methoxy occupied *ortho* and *para* position of Ring B. Presence of prenyl group reduce this activity.

4.4 Reducing power capacity

It has been reported that there is a direct correlation between antioxidant activities and reducing power of chalcones. The reducing properties are generally associated with the presence of reductants, which have been shown to exert antioxidant action by breaking the free radical chain by donating a hydrogen atom. In this test the yellow color of the test solution changed to various shades of green and blue, depending on the reducing power of each sample. The presence of reductants i.e.-antioxidants caused the concentration of the Fe^{3+} /ferric reducing complex to the ferrous form. Therefore, by measuring the formation of Purl's Prussian blue at 700 nm we can monitor the Fe^{2+} concentration; a higher absorbance at 700 nm indicates a higher reducing power. The result is shown in Table 13. The reducing power of all the samples increased gradually with the increase in concentrations of the samples.

Table 13: Reducing power of chalcones (3a-3e, 4a-4c) with standard Ascorbic acid

Compound no.	Conc.($\mu\text{g/mL}$)	Absorbance
3a	5	0.008
	10	0.014
	15	0.063
	20	0.078
3b	5	0.079
	10	0.088
	15	0.099
	20	0.136
3c	5	0.290
	10	0.561
	15	0.869
	20	1.106
3d	5	0.044
	10	0.059
	15	0.063
	20	0.065
3e	5	0.016
	10	0.060
	15	0.129
	20	0.155
4a	5	0.004
	10	0.017
	15	0.037
	20	0.048

4b	5	0.200
	10	0.304
	15	0.342
	20	0.355
4c	5	0.070
	10	0.219
	15	0.247
	20	0.260
Ascorbic acid	5	0.659
	10	0.914
	15	2.232
	20	2.395




Figure 10: Reducing power of chalcones (3a-3e, 4a-4c) with standard Ascorbic acid




Figure 11: Reducing power of chalcones (3a-3e, 4a-4c) without standard




Figure 12: Reducing power of chalcones (3a-3e, 4a-4c) compare to standard Ascorbic acid in 10 µg/mL concentration

All the chalcones show appreciable reducing activity. Insertion of prenyl functionality increase reducing capacity.

4.5 Cytotoxic activity ^[6]

Brine shrimp lethality test (BST) has been applied as a simple and useful tool for preliminary screening of toxicity of physiologically active plant extract or synthesized compounds, detection of fungal toxins, heavy metal, pesticides and also cytotoxicity testing of dental materials. This general bioassay is rapid, reliable and has been used for over thirty years in toxicological studies. However, it has been demonstrated that a positive relationship exists between brine shrimp lethality and human carcinoma. Thus, BST can also be extrapolated for cell line toxicity and anti tumor activity. Principal of this method was based on the ability of certain compounds to kill laboratory cultured *Artemia nauplii* brine shrimp. BST is one of the simplest biological responses to monitor is lethality, since there is only one criterion: either dead or alive. It has been shown that *Artemia* is highly vulnerable to toxins at the early developmental stages and assumed to exhibit their greatest sensitivity to test compounds. Subsequently, in this study we used nauplii as object experimental.

Table 14: Brine Shrimp Lethality Assay of Chalcones (3c-3e, 4a-4c) in DMSO

Comp. no	Conc. µg /mL	% of mortality	*LC ₅₀
3c	2.5	11.11	54.22
	5	12.50	

3d	2.5	11.11	71.75
	5	12.50	
	10	14.28	
	20	14.28	
	40	28.57	
	80	57.14	
3e	2.5	33.33	10.10
	5	37.50	
	10	62.50	
	20	71.43	
	40	71.43	
4a	2.5	55.55	0.27
	5	62.50	
	10	75.00	
4b	2.5	44.44	1.61
	5	50.50	
	10	57.14	
	20	57.14	
	40	85.71	
	80	85.71	
4c	2.5	33.33	29.38
	5	37.50	
	10	37.50	
	20	42.86	
	40	57.14	

*3a and 3b do not show any presentable data and excluded from this table. Only those conc. of 3c-4c are included at which nauplii remain alive.




Figure 13: Bar diagram of LC₅₀

From literature review [6] it is known LC₅₀ value greater than 1000 ppm for plant extracts was considered inactive, whereas LC₅₀ values less than 30 ppm for pure compounds were considered toxic. All compounds (**3c-3e**, **4a-4c**) showed a dose dependent cytotoxic activity at the tested concentrations. The LC₅₀ results of chalcone derivatives evaluated in this screening are listed in Table 14. Compound **3d** was the most active than other compounds, presenting the highest LC₅₀ of 71.75 ppm. **3c** also showed better activity LC₅₀ value 54.22 ppm. Whereas from the result **3e**, **4a**, **4b**, **4c** showed poor activity. It is also known from the literature [6] chalcone containing *o*-methoxy substitution in ring-B reducing its ability to kill *Artemia nauplii* due to steric hindrance to the receptor of specimen. In **3c** and **3d** there are three methoxy functionality in ring B and in **3d** both the *ortho* position are occupied by methoxy group so **3d** is least toxic. In **3c** only one *ortho* position is occupied by methoxy group so it has lower LC₅₀ value than that of **3d** and slightly toxic compared to **3d**. **4a**, **4b**, **4c** these three compounds contain lipophilic prenyl functionality which enhance cell membrane permeability and increase toxicity. Among these three prenylated chalcones **4c** contain *ortho*, *para* methoxy functionality and having the higher LC₅₀ value 29.38 ppm. The present study also revealed that **3a**, **3b**, **3e**, **4a**, **4b** synthesized compounds are toxic against *Artemia sp.* Therefore, these compounds should be studied furthermore for getting antitumor compounds.



CHAPTER FIVE

CONCLUSION

CHAPTER FIVE

Conclusion

Bangladesh is predominantly an agricultural country depending mainly on crop plants, agricultural, medicinal and forest product for its economic development. Although crops play a vital role in the economy of the country and agro ecological conditions are favorable for the production of various crops, the yields of crops is often poor. Among the various factors responsible for poor yield of crops, plant disease caused by various microorganisms play a significant role. Gradually men gathered sufficient knowledge of chemistry to inhibit or to kill the microorganism i.e., only inhibit the microbial growth are called 'statis'. But the chemicals which have ability to kill microorganisms are called 'cadal'. Various pesticides are classified as fungicides, viricides, bactericides etc.

Our previous study on synthesis of chalcone, hydroxychalcone, chalcone epoxide, flavone and oxygenated flavones which exhibiting a good antimicrobial activity have been reported for long. ^[21, 22] In the present study we are interested to extend our studies by the introduction of hydroxyl, methoxy and prenyl group at different position of benzene nucleus moiety of chalcone (Figure 1) which may give an important information about positional effect of these substituents and its role as biofunctionality.

Chalcones are the important constituent of many natural sources and have variety of biological activities. The synthetic technique is based on Claisen-Smith base catalyzed condensation of substituted acetophenone (**1a-1c**) with substituted benzaldehyde (**2a-2e**) shown in scheme 7. Thus we have successfully synthesized feight substituted chalcones. The parent chalcone is 2' - hydroxychalcone (**3a**) and its methoxy derivatives ;2' - hydroxy - 4 - methoxychalcone (**3b**), 2' - hydroxy - 2, 4, 5 - trimethoxychalcone (**3c**), 2', 5'

- dihydroxy - 2, 4, 6 - trimethoxychalcone (**3d**), 4 - hydroxy - 3', 4', 5'-trimethoxychalcone (**3e**). **3a** and **3d** were also undertaken by prenylation to produce their prenylated derivatives; 2' - hydroxy - 3' - C - prenylchalcone (**4a**), 2' - hydroxy - 5' - C - prenylchalcone (**4b**), 2', 5'- dihydroxy - 2, 4, 6 - trimethoxy - 3' - C - prenylchalcone (**4c**). The compounds were purified by recrystallation, PTLC or column chromatography. All the compounds provided a single spot in TLC analysis and possess a very sharp melting range, therefore it can be concluded that the synthesized compounds were pure. The structures of the synthesized compounds were confirmed by UV, IR, ^1H and ^{13}C NMR spectroscopic data. Chalcones, a group of compounds with two aromatic rings connected by a keto-vinyl chain, $-\text{CO}-\text{CH}=\text{CH}-$ so these have some unique structural features which were noticed by analysis and important spectral values are presented in tabular form in Table 7.

Once the structure of a biological active compound is known the medicinal chemist is ready to move on to study the structure activity relationship by varying slightly from the original molecule and studying what effect this has on biological activity. The present study includes design a suitable method for the synthesis of chalcones, with varying hydroxy, methoxy and prenyl substituents and to provide SAR through antibacterial, antioxidant and cytotoxicity screening.

Compounds with electron releasing groups such as hydroxyl and methoxy showed better antibacterial activity than the others not having such groups. All the compounds were employed as test chemicals for *in vitro* antibacterial potentiality test against four pathogenic bacteria viz. *Bacillus caerius* (G^+ , B₁), *Staphylococcus aureus* (G^+ , B₂), *Eschericia coli* (G^- , B₃), *Agrobacterium Species* (G^- , B₄). The primary assay was performed by disc diffusion technique to classify the microorganism susceptible as well as resistant towards particular compounds. The bioactivity is expressed by the

diameter of zone of inhibition in mm and presented in Table 8-Table 11. Most of the compounds show moderately good antibacterial activity against both G^+ and G^- bacteria.

The trend for antibacterial (*Bacillus caerius*, G^+ , B_1) activity at conc. $250\mu\text{g disc}^{-1}$

C-50 (40 mm) > 3d (38 mm) > 3c (30 mm) > 3b (28 mm) > 3a (24 mm) > 4c (20 mm) > 3e, 4b (12 mm) > 4a (08 mm)

The trend for antibacterial (*Staphylococcus aureus*, G^+ , B_2) activity at conc. $250\mu\text{g disc}^{-1}$

C-50 (42 mm) > 3b (40 mm) > 4c (30 mm) > 3d (25 mm) > 3a, 3c (20 mm) > 3e, 4b (12 mm) > 4a (08 mm)


The trend for antibacterial (*Escherichia coli*, G^- , B_3) activity at conc. $250\mu\text{g disc}^{-1}$

C-50 (42 mm) > 3b, 3c (30mm) > 4c (26 mm) > 3d (20mm) > 3a (18 mm) > 3e, 4a, 4b (06mm)

3b, 3c, 3d exhibited fairly good potentialities against both bacteria and in some cases about near to that of the standard drugs. Prenyl derivative of compound **3d**, compound **4c** show very high activity against (G^-) than other prenyl derivatives. This led us to conclude that the presence of electron releasing hydroxyl (-OH) and methoxy group (-OCH₃) are responsible for the antimicrobial effects. **4c** contains lipophilic prenyl group and more susceptible to (G^-). In **3e** three methoxy groups are occupied ring A and in three adjacent carbon. So steric effect may be a cause of poor activity.

In addition the synthesized chalcones (**3a-3e, 4a-4c**) were evaluated for *in vitro* antioxidant activity using diphenylpicrylhydrazyl (DPPH) model. Observation for antioxidant activity is expressed in terms of percent

scavenging of DPPH radical and the inhibitory concentration 50% (IC_{50}) is lowest for structure **3d**, 0.923 ($\mu\text{g/mL}$) at DPPH conc. 0.02% indicating highest activity and presented in Table 12. The compound **3d** contains two phenolic -OH group and suppose to produce phenoxide free radical easily and stabilized by electronic group methoxy occupied *ortho* and *para* position of aromatic ring attached to benzaldehyde part. Other structures showed good antioxidant activity as compared to the standard, ascorbic acid, IC_{50} 0.08 ($\mu\text{g/mL}$). The antioxidant activity was also measured in another method using reducing power capacity. All the chalcones especially prenylated chalcone showed appreciable reducing activity and presented in Table 13, **3c** showed the highest value. Cytotoxic activities of the same compounds were undertaken *in vivo* by brine shrimp lethality test (BST) and expressed by lethal concentration 50%, (LC_{50}) and presented in Table 14. The compound **3d** showed the least toxicity having LC_{50} 71.75 ($\mu\text{g/mL}$) whereas LC_{50} values less than 30 ppm for pure compounds were considered toxic. Increment the no. of methoxy and hydroxy group causes higher LC_{50} values among the compounds (**3a-3d**). As steric hindrance effect causes better tolerance by reducing their killing ability. On the contrary prenylation causes cell membrane permeability and enhance toxicity. In **3e** three methoxy groups are occupied ring A and in three adjacent carbon. It shows poor activities in any sort of biological screening undertaken. All the examined compounds can be regarded as prosperous materials to introduce a safe, cheap and easy to prepare natural coloring agent for food and beverage industry, textile replacing costly natural dye or harmful azo dyes as well as antioxidant in food preservation and in cosmetic industry.



REFERENCES


References

1. Carlo, G. Di., Mascolo, N., Izzo, A. A. and Capasso, F. (1999), Flavonoids; Old and New Aspects of A Class of Natural Therapeutic Drugs, *Life Sci.*, **65**, 337-353.
2. Patolia, P.U., Gohel, V. P., Purohit, D. M. and Patolia, V. N. (2014), Synthesis and Biological Evaluation of Some New Chalcone and Isoxazole Derivatives, *Int. J. Sci and Tech.*, **2** (5), 137.
3. Dodia, J. V., Dangar, V. R. and Shah, V. R. (2014), Synthesis and Characterization of Some Isoxazole Derivatives of Vanillin as Antimicrobial Agent, *Int. J. Sci and Tech.*, **2** (4), 64.
4. Kostanecki, S.V. and Tambor, J. (1899), *Chem. Ber.*, **32**, 1921.
5. Lumbiny, B. J., Islam, M. A., Hossain, M. J., and Khatun, M. M., (2014), Synthesis and Bioactivity Screening of Some Natural Colouring Agents, Substituted Chalcones; An Environment-Friendly Alternative to Harmful Azo Dyes. *Raj. Univ. J. Env. Sci.*, **3**, 65-74.
6. Sulistyotywyaty, M. I., Nofianti, K. A., Suzana and Budiati, T. (2013), Synthesis and Brine Shrimp Bioassay of Chalcone and Its Two Methoxy Derivatives. *Int. J. Phar. & Chem. Sci.* **2** (2), 719-724.
7. Mabry, T. J., Markbam, K. R. and Thomas, M. B. (1970), *The Systematic Identification of Flavonoids*. Springer-Verlag, New York, Heidelberg, Berlin.
8. Patil, C. B., Mahajan, S. K. and Katti, S. A. (2009), Chalcone: A Versatile Molecule. *J. Pharm. Sci. & Res.* **1**(3), 11-22.

9. Dyrager C. (2012), Design and synthesis of Chalcone and Chromone Derivatives as Novel Anticancer Agents. Doctoral thesis. Department of Chemistry University of Gothenburg.
10. Claisen, L. and Claparede, A. (1881), *Chem. Ber.*, **14**, 2463.
11. M. Sc. thesis (Roll no. 11063, 2011) An Eco-Friendly Synthesis of Some Non-Natural Microwave Condition and the Screening of Their Antimicrobial/Antioxidant Activity against *E.Coli* and DPPH, Jahanginagar University, Bangladesh.
12. M. Sc. thesis (Roll no. 08224572, 2012) Effect of Hydroxy, Methoxy and Benzoyloxy Substituents on Structure Activity Relationship of Some Synthesized Chalcones; Determined by Antimicrobial and Antioxidant activity, Rajshahi University, Bangladesh.
13. Narender, T. and Papi Reddy, K. (2007), A Simple and Highly Efficient Method for the Synthesis of Chalcones by Using Borontrifluoride-Etherate. *Tet Letts*, **48**:3177-3180
14. Chaudhury, N. A., Kirtaniya, C. L. and Mukharjee, B. (1971), *Tetrahedron*, **27**, 21.
15. Russel, A. and Todd, J. (1934), *J. Chem. Soc.*, 1506.
16. Stout, G. H. and Stouf, V. F. (1961), *Tetrahedron*, **14**, 296.
17. Bhatia, V. K. and Kagon, (1970), *Indian J. Chem.*, 1203.
18. Grinor, G. V., Chervenuk, G. I. and Dombrovskii, A. V. (1968), *U. Obsh. Kim.*, **18** (2), 225.
19. Lupi, A., Monache, G. D., Monache, F. D., Masini, Bettelo, G. B., Lima, O. G. D. and Mello (1975), *J. F., I. L. Farmco, Ed. Sc.* **30**, 450.
20. Obera, G., Ondera, J. and Kurikhara, V. (1978), *Bull. Chem. Soc.*, **51**, 362.

21. Islam, M. A. (1981), Chemical Investigation of *Milletia Ovalifolia* Seeds and Synthesis of Desmodol and Sieisocordin, Ph. D Thesis, New Delhi University, India.
22. (a) Alam, S., Miah, M. A. J. and Islam, M. A. (2007), Synthesis of Maximaflavanone-A and Its Derivative. *Pak. Journal of Science Res.*, vol. **50** (2), 80-84. (b) Islam, A., Gupta, R. K. and Krishnamurti, M. (1980), *Phytochemistry*, **19**, 1558-1559. (c) Islam, A., Gupta, R. K. and Krishnamurti, M. *Indian J. Chem.* (1981), **20 B**, 21-22. (d) Alam, S., Miah, M. A. J. and Islam M. A. (2004), *J. Biol. Sci.*, **4** (4), 527-531. (e) Morshed M. H. (2005) *et al.*, *Pak. Journal of Biological Science*, vol. **8** (7), 1025-1029. (f) Mostahar, S. Alam, S. & Islam, A. (2006), *Ind. J. Chem.*, **45 B**, 478-486. (g) Mostahar, S., Khatun, P. and Islam, A. (2007), *J. Biol. Sci.*, **7** (3), 514-519. (h) Lumbiny, B. J., Rahman, M., Khatun, M. M., Akter, R., Khatun, M. R. and Islam, M. A. (2015), An Environmentally Safe Green Approach of Synthesis of Some New 2'-Hydroxychalcone Epoxides and Their Anti-Infective Activity Screening, *Raj. Univ. J. Env. Sci.*, **4**, 67-74.
23. Gupta, D., Jain D. K. and Trivedi, P. (2010), Recent Advances in Chalcones as Anti-infective Agents, *Int. J. Chem. Sci.*: **8** (1), 649-654
24. Bauer, A. W., Kirby, W. M. M., Sherris, J. C. and Turck, M. (1966), *Am. J. Clin. Pathol.* **44**, 493.
25. Gnananickam, S. S., Smith, D. A. (1980), Selective Toxicity of Isoflavonoid Phytoalexins to Grampositive bacteria, *Phytopathology*, **70**, 894
26. Reiner, R. (1982) Detection of Antibiotic Activity. In, Antibiotics an introduction, Roche scientific services, Switzerland, p-70.
27. Nishina, C. *et al.*, (1987), *Agric. Biol. Chem.*, **51**, 139.

28. Sreedhar, V., Nath, LK. R., Gopal, N. M. and Nath, M. S. (2010) *In-vitro* Antioxidant Activity and Free Radical Scavenging Potential of Roots of Vitex Trifoliata, *RJPBS*, **1** (4).
29. Bors *et al.*, (1990), *Methods Enzymol.*, **186**, 343.
30. Prakash, A., Regelhof, F. and Miller, E., Antioxidant Activity, Analytical Progress, (A Publication of Medallion Labs)
31. Singleton, V. L. and Rossi, J. A. (1965), Colorimetry of Totalphenolics With Phosphomolybdic Phosphotungstic Acid Reagents. *American Journal of Enology and viticulture*, **16**, 144-158
32. Mostahar, S. M. (2002) Synthesis of some 2'-Oxygenated Flavones and Their Antimicrobial and Cytotoxic Activity, M. Sc. thesis, Rajshahi University, Bangladesh.
33. Prashad, K. and Mohan, S. (2012) Synthesis, Characterization and Antimicrobial Activity of Some Chalcones, *Int. J. Pharm. & Tech.*, **4** (1), 4060-4066.
34. Prior, R. L. Wu, X and Schaich, K. (2005), standardized Methods for the Determination of Antioxidant Capacity and Phenolics in Foods and Dietary Supplements, *J. Agric. Food Chem.*, **53**, 4290-4302
35. Carballo JL. (2002) A Comparison Between Two Brine Shrimp Assay to Detect *In Vitro* Toxicity in Marine Natural Products, *BMC Biotechnology.*, **2** (17), 1472-6750.
36. Kesavan SS., (2011), Application of Brine Shrimp Bioassay for Screening Cytotoxic Actinomycetes, *International Journal of Pharmacy and Pharmaceutical Science Research.*, **1** (3), 104-107.
37. Meyer, (1982), Brine Shrimp A Convenient General Bioassay for Active Plant Constituents, *Planta Medical*, **45**:31-34.



Appendix-A: List of Spectra

1. **Figure-1:** IR spectrum of 2' - hydroxychalcone, **3a** in KBr cm^{-1}
2. **Figure-2:** ^1H NMR spectrum of 2' - hydroxychalcone, **3a**, in CDCl_3 , 400 MHz
3. **Figure-3:** ^{13}C NMR spectrum of 2' - hydroxychalcone, **3a**, in CDCl_3 , 100 MHz
4. **Figure-4:** IR spectrum of 2' - hydroxy - 4 - methoxychalcone, **3b** in KBr cm^{-1}
5. **Figure-5:** ^1H NMR spectrum of 2' - hydroxy - 4 - methoxychalcone, **3b** in CDCl_3 , 500 MHz
6. **Figure-6:** ^{13}C NMR spectrum of 2' - hydroxy - 4 - methoxychalcone, **3b** in CDCl_3 , 125 MHz
7. **Figure-7:** IR spectrum of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone, **3c** in KBr cm^{-1}
8. **Figure-8:** ^1H NMR spectrum of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone, **3c** in CDCl_3 , 400 MHz
9. **Figure-9:** ^{13}C NMR spectrum of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone, **3c** in CDCl_3 , 100 MHz
10. **Figure-10:** IR spectrum of 2', 5' -dihydroxy - 2, 4, 6 - trimethoxychalcone, **3d** in KBr cm^{-1}
11. **Figure-11:** ^1H NMR spectrum of 2', 5' -dihydroxy - 2, 4, 6- trimethoxy chalcone, **3d** in CDCl_3 , 400 MHz
12. **Figure-12:** ^{13}C NMR spectrum of 2', 5' -dihydroxy - 2, 4, 6- trimethoxy chalcone, **3d** in CDCl_3 , 100 MHz
13. **Figure-13:** IR spectrum of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone, **3e** in KBr cm^{-1}

14. **Figure-14:** ^1H NMR spectrum of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone, **3e** in CDCl_3 , 400 MHz
15. **Figure-15:** ^{13}C NMR spectrum of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone, **3e** in CDCl_3 , 100 MHz
16. **Figure-16:** ^1H NMR spectrum of 2' - hydroxy - 3' - C - prenylchalcone, **4a** in CDCl_3 , 500 MHz
17. **Figure-17:** ^{13}C NMR spectrum of 2' - hydroxy - 3' - C - prenylchalcone, **4a** in CDCl_3 , 125 MHz
18. **Figure-18:** ^1H NMR spectrum of 2' - hydroxy - 5' - C - prenylchalcone, **4b** in CDCl_3 , 500 MHz
19. **Figure-19:** ^{13}C NMR spectrum of 2' - hydroxy - 5' - C - prenylchalcone, **4b** in CDCl_3 , 125 MHz
20. **Figure-20:** ^1H NMR spectrum of 2', 5' - dihydroxy -2, 4, 6 - trimethoxy - 3' - C - prenylchalcone, **4c** in CDCl_3 , 400 MHz
21. **Figure-21:** ^{13}C NMR spectrum of 2', 5' - dihydroxy -2, 4, 6 - trimethoxy - 3' - C - prenylchalcone, **4c** in CDCl_3 , 100 MHz




Figure-1: IR spectrum of 2' - hydroxychalcone, **3a** in KBr cm⁻¹




Figure-2: ¹H NMR spectrum of 2' - hydroxychalcone, 3a, in CDCl₃, 400 MHz


INARS, BCSIR, ¹³C spectrum, 01 in CDCl₃, Marjina Khatun, RU

NMR



Current Data Parameters
NAME: A-347_D-254_Marjina Khatun_RU
PROG: 1

F2 - Acquisition Parameters
Date_ 20150427
Time 10.48
INSTRUM spect
PROBHD 5 mm PABD BE7
PULPROG zgpg30
SOLVENT CDCl₃
NS 2
DS 4
SWH 29761.974 Hz
FIDRES 0.164133 Hz
AQ 1.117133 sec
RG 1638
DM 16.00 usec
DE 1.00 usec
TE 300.2 K
D1 1.00000000 sec
D11 0.05000000 sec
D12 0.05000000 sec
D13 0.05000000 sec
D14 0.05000000 sec
D15 0.05000000 sec
D16 0.05000000 sec
D17 0.05000000 sec
D18 0.05000000 sec
D19 0.05000000 sec
D20 0.05000000 sec
D21 0.05000000 sec
D22 0.05000000 sec
D23 0.05000000 sec
D24 0.05000000 sec
D25 0.05000000 sec
D26 0.05000000 sec
D27 0.05000000 sec
D28 0.05000000 sec
D29 0.05000000 sec
D30 0.05000000 sec
D31 0.05000000 sec
D32 0.05000000 sec
D33 0.05000000 sec
D34 0.05000000 sec
D35 0.05000000 sec
D36 0.05000000 sec
D37 0.05000000 sec
D38 0.05000000 sec
D39 0.05000000 sec
D40 0.05000000 sec
D41 0.05000000 sec
D42 0.05000000 sec
D43 0.05000000 sec
D44 0.05000000 sec
D45 0.05000000 sec
D46 0.05000000 sec
D47 0.05000000 sec
D48 0.05000000 sec
D49 0.05000000 sec
D50 0.05000000 sec
D51 0.05000000 sec
D52 0.05000000 sec
D53 0.05000000 sec
D54 0.05000000 sec
D55 0.05000000 sec
D56 0.05000000 sec
D57 0.05000000 sec
D58 0.05000000 sec
D59 0.05000000 sec
D60 0.05000000 sec
D61 0.05000000 sec
D62 0.05000000 sec
D63 0.05000000 sec
D64 0.05000000 sec
D65 0.05000000 sec
D66 0.05000000 sec
D67 0.05000000 sec
D68 0.05000000 sec
D69 0.05000000 sec
D70 0.05000000 sec
D71 0.05000000 sec
D72 0.05000000 sec
D73 0.05000000 sec
D74 0.05000000 sec
D75 0.05000000 sec
D76 0.05000000 sec
D77 0.05000000 sec
D78 0.05000000 sec
D79 0.05000000 sec
D80 0.05000000 sec
D81 0.05000000 sec
D82 0.05000000 sec
D83 0.05000000 sec
D84 0.05000000 sec
D85 0.05000000 sec
D86 0.05000000 sec
D87 0.05000000 sec
D88 0.05000000 sec
D89 0.05000000 sec
D90 0.05000000 sec
D91 0.05000000 sec
D92 0.05000000 sec
D93 0.05000000 sec
D94 0.05000000 sec
D95 0.05000000 sec
D96 0.05000000 sec
D97 0.05000000 sec
D98 0.05000000 sec
D99 0.05000000 sec
D100 0.05000000 sec



3a




Figure-3: ¹³C NMR spectrum of 2' - hydroxychalcone, 3a, in CDCl₃, 100 MHz




Figure-4: IR spectrum of 2' - hydroxy - 4 - methoxychalcone, **3b** in KBr cm⁻¹






Figure-5: ¹H NMR spectrum of 2' - hydroxy - 4 - methoxychalcone, 3b in CDCl₃, 500 MHz



F2 - Acquisition Parameters
 Date_ 2026-05-03
 Time 12:44
 INSTRUM spect
 PROBEID 5 mm QNP1 13
 PULPROG zgpg30
 TD 65536
 SOLVENT CDCl3
 NS 4
 DS 4
 SWH 18841.270 Hz
 FIDRES 0.336754 Hz
 AQ 1.0515572 sec
 RG 6584
 DN 25.000 MHz
 DE 5.00 MHz
 TE 300.2 K
 D12 0.0000000 sec
 PL1 22.00 dB
 PL5 22.00 dB
 D1 2.0000000 sec
 D11 0.0500000 sec
 D12 0.0500000 sec
 SFO2 250.132358 MHz
 NUC1 13C
 NUC2 1H
 P1 6.00 sec
 PL1 22.00 dB
 P1 10.00 sec
 SF01 62.8931573 MHz
 NUC1 1H
 PL1 0.00 dB
 D11 1.0000000 sec
 F2 - Processing parameters
 SI 32768
 SF 62.8931573 MHz
 WCN EN
 SSB 0
 LB 1.00 Hz
 GB 0
 PC 1.00
 F2 - INE-plot parameters
 DI 20.00 cm
 F1 200.701 MHz
 F2 125.013 MHz
 F3 5.621 MHz
 F4 365.411 MHz
 PPM0 0.24888 ppm/Hz
 MLEV 612: 80083 Hz/Hz

Figure-6: ¹³C NMR spectrum of 2'-hydroxy - 4 - methoxychalcone, 3b in CDCl₃, 125 MHz



Sam_2_155

Figure-7: IR spectrum of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone, **3c** in KBr cm⁻¹




Figure-8: ¹H NMR spectrum of 2' - hydroxy - 2, 4, 5 - trimethoxychalcone, **3c** in CDCl₃, 400 MHz




Figure-9: ¹³C NMR spectrum of 2'-hydroxy-2,4,5-trimethoxychalcone, **3c** in CDCl₃, 100 MHz




Figure-10: IR spectrum of 2', 5' -dihydroxy - 2, 4, 6 - trimethoxychalcone, 3d in KBr cm⁻¹




Figure-11: ¹H NMR spectrum of 2', 5' -dihydroxy - 2, 4, 6- trimethoxy chalcone, **3d** in CDCl₃, 400 MHz




Figure-13: IR spectrum of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone, **3e** in KBr cm⁻¹



INARS, BCSIR, ¹³C spectrum, Chloroform in CDCl₃, Margina, RU

Current Data Parameters
NAME A-887_D-678_Margina_RU
EXPER 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20141218
Time 15:15
INSTRUM spect
PROBHD 5 mm PABBO BB7
PULPROG zgpg30
AQ 25.16
RG 25516
SOLVENT CDCl₃
DS 9542
NS 1
SMH 29761.904 Hz
FIDRES 0.454131 Hz
AQ 1.101048 sec
RG 197.33
OR 16.800 MHz
DE 9.21 MHz
TE 300.2 K
TD 1.8000000 sec
D11 0.31000000 sec
TD0 1

CHANNEL F1
SFO1 100.6254023 MHz
NUC1 13C
P1 31.00 usec
PLW1 70.00000000 W

CHANNEL F2
SFO2 400.1720005 MHz
NUC2 1H
P2 18.00000000 usec
PLW2 17.00000000 W
SFO3 0.23139500 MHz
PLW3 0.38742000 W

F2 - Processing Parameters
SI 32748
SF 100.6252533 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40







Figure-15: ¹³C NMR spectrum of 4 - hydroxy - 3', 4', 5' - trimethoxychalcone, **3e** in CDCl₃, 100 MHz





F2 - Acquisition Parameters
 Date_: 2026-09-03
 Time: 12:44
 INSTRUM: spect
 PROBRG: 5 in. Out 1.1
 PULPROG: zgpg30
 TL: 60000
 SOLVENT: DMSO
 NS: 328
 DS: 4
 SWH: 18041.270 Hz
 FIDRES: 0.2007854 Hz
 AQ: 1.12515577 sec
 RG: 10268
 DA: 221.200 mm
 DE: 1.00 mm
 TE: 300.2 K
 D12: 0.00002000 sec
 P-13: 22.10 us
 D1: 2.00000000 sec
 D11: 0.05000000 sec
 D12: 0.05000000 sec
 SFO2: 250.1325346 MHz
 NUC2: 13C
 P-12: 0.20 us
 P-11: 22.10 us
 SFO1: 125.7611533 MHz
 NUC1: 13C
 P-1: 0.05 us
 D11: 1.05000000 sec
F2 - Processing parameters
 SI: 32768
 SF: 62.8250000 MHz
 MD: 64
 SSB: 0
 LB: 1.00 Hz
 GB: 0
 PC: 1.4
13C NMR parameters
 CD: 20.00 us
 CF: 200.701 MHz
 F2: 125.7611533 MHz
 F2F: 5.00 MHz
 F2: 365.11 Hz
 PPHO: 5.74354 ppm/ct
 HEDY: 61.00000000 Hz/ct

Figure-17: ¹³C NMR spectrum of 2'-hydroxy-3'-C-prenylchalcone, 4a in CDCl₃, 125 MHz

sample 5
cdci3

7.85
7.83
7.81
7.80
7.79
7.78
7.77
7.76
7.75
7.74
7.73
7.72
7.71
7.70
7.70
7.70
7.69
7.68
7.67
7.66
7.66
7.66
7.66
7.65
7.64
7.63
7.62
7.61
7.60
7.59
7.58
7.57
7.56
7.55
7.54
7.53
7.52
7.51
7.50
7.49
7.48
7.47
7.46
7.45
7.44
7.43
7.42
7.41
7.40
7.39
7.38
7.37
7.36
7.35
7.34
7.33
7.32
7.31
7.30
7.29
7.28
7.27
7.26
7.25
7.24
7.23
7.22
7.21
7.20
7.19
7.18
7.18
7.17
7.16
7.15
7.14
7.13
7.12
7.11
7.10
7.09
7.08
7.07
7.06
7.05
7.04
7.03
7.02
7.01
7.00
6.99
6.98
6.97
6.96
6.96
6.96
6.95
6.94
6.93
6.92
6.91
6.90
6.89





Figure-18: ^1H NMR spectrum of 2' - hydroxy - 5' - C - prenylchalcone, **4b** in CDCl_3 , 500 MHz

FE - Acquisition Parameters
Date: 20/05/2013
Time: 12:44
INSTRUM: spect
PROBHD: 5 mm QNP113
PULPROG: zgpg30
TD: 65536
SOLVENT: DMSO
NS: 328
DS: 4
SWH: 18844.270 MHz
FIDRES: 0.300754 Hz
AQ: 1.8515570 sec
RG: 14224
IN: 22.700 mmol
DE: 0.00 mmol
TE: 300.2 K
D12: 0.00000000 sec
P1: 22.00 sec
D1: 2.00000000 sec
CPDPRG2: waltz16
PCPD2: 90.00 MHz
NUC1: 13C
PL1: 0.00 sec
P12: 22.00 sec
PL12: 1E: 0.00 sec
SFO: 62.501975 MHz
AQC: 1E: 0.00 sec
P1: 0.00 sec
D1: 1.00000000 sec

FE - Processing parameters
SI: 32768
SF: 62.501975 MHz
WDW: EM
SSB: 0
LB: 1.00 Hz
GB: 0
PC: 1.4

SC - 13C: parameters
D1: 0.00 sec
SFO: 200.700 MHz
F1: 12820.13 MHz
F2: 5.000 MHz
F3: 30.111 MHz
PPM0: 1.74265 ppm/CF
PPM1: 6.121805 ppm/CF




Figure-19: ¹³C NMR spectrum of 2' - hydroxy - 5' - C -prenylchalcone, 4b in CDCl₃, 125 MHz

INARS,BCSIF,1H spectrum, Chloroform 30.11.14 in CDCl₃,Margina,RU



Current Data Parameters
 NAME A-887_D-678_Margina_RU
 EXPNO 5
 PROCNO 1

F2 - Acquisition Parameters
 Date_ 20141222
 Time 9.44
 INSTRUM spect
 PROBHD 5 mm F4BBO BB/
 PULPROG zg30
 TD 65536
 SOLVENT CDCl₃
 NS 64
 DS 1
 SWH 6002.401 Hz
 FIDRES 0.091589 Hz
 AQ 5.4591489 sec
 RG 127.57
 DR 83.300 usec
 DE 13.63 usec
 TE 300.0 K
 FL 1.00000000 sec
 TD0 1

===== CHANNEL f1 =====
 SFO1 400.1724008 MHz
 NUCL1 1H
 FL1 10.50 usec
 PLW1 17.00000000 W

F2 - Processing Parameters
 SI 32768
 SF 400.1700121 MHz
 MDW EM
 SSB 0
 LB 0
 GB 0
 PC 5.00






Figure-20: ¹H NMR spectrum of 2', 5' - dihydroxy -2, 4, 6 - trimethoxy - 3'- C -prenylchalcone, **4c** in CDCl₃, 400 MHz



F2 - Acquisition Parameters
Date_ 2008-06-03
Time 12:44
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zgpg30
TD 65536
SOLVENT CDCl3
RG 327.5
DS 4
SWH 19841.200 MHz
FIDRES 0.357754 MHz
AQ 1.0519571 sec
RG 16384
IN 25.000 MHz
DE 15.00 MHz
TE 300.2 K
D12 0.00002000 sec
PL12 22.00 dB
D1 2.00000000 sec
CPDPRG2 waltz16
PCPD2 90.00 MHz
SFO2 250.1312348 MHz
NUC2 13
PL2 0.00 dB
PL12 22.00 dB
P1 1.00 sec
SFO1 62.9031575 MHz
NUC1 13
PL1 0.00 dB
D11 1.05000000 sec

F2 - Processing Parameters
SI 32768
SF 62.9031575 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.00

1D NMR file parameters
SI 20.00 MHz
SF 200.701 MHz
PC 1.052013 sec
PZ 5.00 sec
F2 360.11 MHz
P1 1.74289 sec/cf
F2CF 6.128063 Hz/cf

Figure-21: ¹³C NMR spectrum of 2', 5' - dihydroxy -2, 4, 6 - trimethoxy - 3'- C - prenylchalcone, 4c in CDCl₃, 100 MHz